

E 184

*Printed or made in Great Britain
by Watts & Co., Johnson's Court, Fleet Street,
London, E.C. 4*

CONTENTS OF VOL. 1.

	PAGE
LIST OF ILLUSTRATIONS	v
GLOSSARY	vii
TRANSLATOR'S PREFACE	ix
TABLE - CLASSIFICATION OF THE ANIMAL WORLD	xiv

PREFACE	
I. THE FUNDAMENTAL LAW OF ORGANIC EVOLUTION	1
II. THE OLDER EMBRYOLOGY	9
III. MODERN EMBRYOLOGY	15
IV. THE OLDER PHYLOGENY	22
V. THE MODERN SCIENCE OF EVOLUTION	28
VI. THE OYSTER AND THE ANGEL	36
VII. CLONATION	51
VIII. THE GASTREA THEORY	59
IX. THE GASTRULATION OF THE VERTEBRATE	71
X. THE COLON THEORY	90
XI. THE VERTEBRATE CHARACTER OF MAN	103
XII. EMBRYONIC SHIELD AND GERMINATIVE AREA	115
XIII. DORSAL BODY AND VENTRAL BODY	125
XIV. THE ARTICULATION OF THE BODY	141
XV. FETAL MEMBRANES AND CIRCULATION	156

LIST OF ILLUSTRATIONS

	PAGE		PAGE
1. The human ovum	36	40. Cleavage of the frog's ovum	72
2. Stem-cell of an echinoderm	37	41-44. Sections of fertilised toad ovum	73
3. Three epithelial cells	38	45-48. Gastrulation of the salamander	74
4. Five spiny or grooved cells	38	49. Segmentation of the lamprey	76
5. Ten liver-cells	38	50. Gastrulation of the lamprey	76
6. Nine star-shaped bone-cells	39	51. Gastrulation of <i>Ceratodus</i>	77
7. Eleven star-shaped cells	39	52. Ovum of a deep-sea bony fish	78
8. Unfertilised ovum of an echino- derm	40	53. Segmentation of a bony fish	79
9. A large branching nerve-cell	41	54. Discord gastrula of a bony fish	80
10. Blood-cells	42	55 and 56. Sections of blastula of shark	80
11. Indirect or mitotic cell-division	43	57. Discord segmentation of bird's ovum	82
12. Mobile cells	44	58-61. Gastrulation of the bird	83
13. Ova of various animals	45	62. Germinal disk of the lizard	84
14. The human ovum	46	63 and 64. Gastrulation of the opossum	85
15. Fertilised ovum of hen	47	65-67. Gastrulation of the opossum	86
16. A creeping amoeba	47	68-71. Gastrulation of the rabbit	88
17. Division of an amoeba	48	72. Gastrula of the placental mammal	89
18. Ovum of a sponge	49	73. Gastrula of the rabbit	90
19. Blood-cells, or phagocytes	50	74 and 75. Diagram of the four sec- ondary germinal layers	91
20. Spermia or spermatozoa	52	76 and 77. Coelomula of <i>Sagitta</i>	92
21. Spermatozoa of various animals	53	78. Section of young <i>Sagitta</i>	93
22. A single human spermatozoon	53	79 and 80. Section of amphioxus- larva	94
23. Fertilisation of the ovum	54	81 and 82. Section of amphioxus- larva	95
24. Impregnated echinoderm ovum	55	83 and 84. Chordula of the amphioxus	96
25. Impregnation of the star-fish ovum	56	85 and 86. Chordula of the amphibia	96
26 and 27. Impregnation of sea-urchin ovum	57	87 and 88. Section of coelomula-emb- ryos of vertebrates	97
28. Stem-cell of a rabbit	58	89 and 90. Section of coelomula-emb- ryo of Triton	97
29. Gastrulation of a coral	61	91. Dorsal part of three Triton-emb- ryos	98
30. Gastrula of a gastropod	63	92. Chordula-embryo of a bird	99
31. Gastrula of a worm	63	93. Vertebrate-embryo of a bird	100
32. Gastrula of an echinoderm	63	94 and 95. Section of the primitive streak of a chick	101
33. Gastrula of an arthropod	63		
34. Gastrula of a mollusc	63		
35. Gastrula of a vertebrate	63		
36. Gastrula of a house sponge	64		
37. Cells from the primary germinal layers	65		
38. Gastrulation of the amphioxus	66		
39. Gastrula of the amphioxus	67		

	PAGE		PAGE
96. Section of the primitive groove of a rabbit	101	158 and 159. Embryo of the amphioxus	143
97. Section of primitive mouth of a human embryo	102	161 and 162. Sections of shark-embryos	146
98-102. Three-fifths primitive vertebrate	106	163. Section of a Triton-embryo	147
103. Redundant mammary glands	112	164-166. Vertebræ	147
104. A Greek gynecomast	113	167. Head of a shark-embryo	148
105. Severance of the dorsal mammal embryo	116	168 and 169. Head of a chick embryo	149
106 and 107. The visceral embryonic vesicle	118	170. Head of a dog-embryo	149
108. Four ectodermic cells	119	171. Human embryo of fourth week	150
109. Two ectodermic cells	119	172. Section of shoulder of a chick-embryo	151
110-113. Ovary of a rabbit	120	173. Section of pelvic region of chick embryo	152
115-118. Embryonic vesicle of a rabbit	121	174. Development of the lizard's legs	153
119. Section of the genitalia of four vertebrates	123	175. Human embryo five weeks old	154
120-123. Embryonic shield of a rabbit	125	176-178. Embryos of the cat	155
124. Lateral view of the amphioxus	127	179. Human embryos	156
125. 6. Hordula of a frog	127	180. Human embryo of fourth week	157
126. Section of frog-embryo	128	181. Human embryo of fifth week	158
127 and 128. Dorsal shield of a chick	128	182. Section of tail of human embryo	159
129. Section of hind end of a chick	129	183 and 184. Human embryo dissected	160
130. Genital area of the rabbit	129	185. Miss Julia Estlin	161
131. Embryo of the opossum	130	186-190. Human embryos	161
132. Embryonic shield of the rabbit	130	191. Human embryo of sixteen to eighteen days	162
133. Human embryo at the nuchal ridge	131	192 and 193. Human embryo of fourth week	162
134. Embryonic shield of rabbit	131	194. Human embryo with its membranes	164
135. Embryonic shield of opossum	132	195. Diagram of the embryonic organs	165
136. Embryonic disk of a chick	133	196. Section of the pregnant womb	166
137. Embryonic disk of a higher vertebrate	134	197. Embryo of Senecio-jeribon	167
138-142. Section of maturing mammal embryo	135	198. Section of pregnant womb	168
143-146. Sections of embryonic chicks	136	199 and 200. Human fetus and placenta	169
147. Section of embryonic chick	136	201. Vitelline vessels in germinative area	170
148. Section of fore-half of chick-embryo	139	202. Boat-shaped embryo of the dog	171
149 and 150. Sections of human embryo	140	203. Lar or white-headed gullion	172
151. Section of a shark-embryo	140	204. Young orange	173
152. Section of a duck-embryo	141	205. Wild orange	174
153-155. Sole-shaped embryonic disk of chick	143	206. Bald-headed chimpanzee	175
156 and 157. Embryo of the amphioxus	144	207. Fetal membranes and umbilical cord	176
		208. Female gorilla	177
		209. Male giant-gorilla	178

GLOSSARY

AKRANIA: animals without skull (*anencephal*)
ANTHROPOGENESIS: the evolution (*genesis*) of man (*anthropos*)

ANTHROPOLOGY: the science of man

ARLITE: (in compounds) the first or typical
 —as, archi-ectula, archi-gastrula, etc.

BIOGENESIS, the science of the genesis of life (*bios*)

BILATE: (in compounds) pertaining to the early embryo (*bilateral* = a bud); hence —

Blastoderm, skin (*derma*) or enclosing layer of the embryo

Blastosphere: the embryo in the hollow sphere stage

Blastula, same as preceding

Epiblast, The outer layer of the embryo (ectoderm)

Hypoblast, the inner layer of the embryo (entoderm)

BRANCHIAL, pertaining to the gills (*branchia*)

CARYO: (in compounds) pertaining to the nucleus (*carion*); hence, —

Caryokinemes, the movement of the nucleus

Caryolysis, dissolution of the nucleus

Caryoplasm, the matter of the nucleus

CENTROBLASTICAL, see under **LEUCITH**

CHORDATA and **CHORDOATA**, animals with a dorsal chord or back-bone

COELOM or **COELOMA**, the body-cavity in the embryo; hence, —

Coelenterata: animals without a body-cavity

Celomata: animals with a body-cavity

Celomatous, formation of the body-cavity

CYTO: (in compounds) pertaining to the cell (*cytos*); hence, —

Cyblast: the nucleus of the cell

Cytodes: cell-like bodies, imperfect cells

Cytoplasm, the matter of the body of the cell

Cytosoma: the body (*soma*) of the cell

CRYPTOCHISM: abnormal retention of the twitches in the body

DATOPHISM: see **PLASM**

DUALISM: the belief in the existence of two entirely distinct principles (such as matter and spirit)

DYSTROLOGY, the science of those features in organisms which refute the "designs-argument"

ECTODERM, the outer (*ecto*) layer of the embryo

ENTODERM, the inner (*ento*) layer of the embryo

EPIDERM: the outer layer of the skin

EPIGENESIS, the theory of gradual development of organs in the embryo

ENTOPHYTES, the third or central eye in the early vertebrates

EPISOMA, see **SOMA**

EPITHELIA, tissues covering the surface of parts of the body (such as the mouth, etc.)

GOVINA, the sexual glands

GOVACHORDISM, separation of the male and female sexes

GOVOTOMES: sections of the sexual glands

GYNAMAST: a male with the breasts (*mastis*) of a woman (*gynae*)

HEPATIC, pertaining to the liver (*hepar*)

HOLOBLASTIC, embryos in which the animal and vegetal cells divide equally (*holos* = whole)

HYPERTASTISM: the possession of more than the normal breasts (*mastis*)

HYPOTACHIAL: underneath (*hypo*) the gills

HYPOMYRIS: sensitive offshoot from the brain in the primitive vertebrate

HYPOMYRIS: see **SOMA**

LECITH-: pertaining to the yolk (*lecithus*); hence:—

Comelecithal: eggs with the yolk in the centre

Lecithoma: the yolk-sac

Telolecithal: eggs with the yolk at one end

METABOLYTIC: cleaving in part (*metabol*) only

META-: (in compounds) the "after" or secondary stage; hence:—

Metagaster: the secondary or permanent gut (*gaster*)

Metoplasia: secondary or differentiated plasma

Metastoma: the secondary or permanent mouth (*stoma*)

Metazoa: the higher or later animals, made up of many cells

Metosum: the mature or advanced stage

METAZOEA: the segments into which the embryo breaks up

METAMORPH: the regeneration of the embryo

MEYER: the most primitive of the unicellular organisms

MEYER: belief in the fundamental unity of all things

MONOTHOUS: the science of organs, forms (generally equivalent to anatomy)

MYOTOMES: segments into which the muscles break up

NEPHRA: the kidneys; hence:—

Nephridia: the rudimentary kidney-organs

Nephrotomes: the segments of the developing kidneys

ONTOGENESIS: the science of the development of the individual (generally equivalent to embryology)

PHAGOCYTES: the genera of the movements in the vital particles

PHAGOCYTIC: cells that absorb food (*phagocyt*—to eat)

PHYLOGENY: the science of the evolution of species (*phyl*)

PLACOTYL: cells that move about (*placotyl*)

PLASMA: the colour or jelly-like matter of which organisms are composed; hence:—

Caryoplasma: the matter of the nucleus (*caryon*)

Cytoplasm: the matter of the body of the cell

Daukoplasm: secondary or differentiated plasma

Metoplasma: same as preceding

Protoplasma: primitive or undifferentiated plasma

PLASMA: the simplest form of plasma

PLASTIDULE: small particles of plasma

POLYSPERMISM: the penetration of more than one sperm-cell into the ovum

PRO- or **PROT-** (in compounds) (in earlier form (opposed to *meta*); hence:—

Prochorion: the first form of the chorion

Progoner: the first reproductive woman

Pronephridia: the earlier form of the kidneys

Prozona: same as preceding

Prostoma: the first or primitive mouth

Protista: the earliest or unicellular organisms

Proterotheca: the earliest phase of the vertebrate

Protophyta: the primitive or unicellular plants

Protoplasma: undifferentiated plasma

Protosoma: the primitive or unicellular animals

RENA: pertaining to the kidneys (*renes*)

SCATULATION: packing or boxing up (*scatula* = a box)

SELFOTOMES: segments into which the primitive skeleton falls

SOMA: the body; hence:—

Cytosoma: the body of the cell (*cyto*)

Epi-soma: the upper or back-half of the embryonic body

Soma: segments of the embryonic body

Hypo-soma: the under or belly half of the embryonic body

TETALOGY: the belief in design and purpose (*telos*) in nature

TETALOGICAL: see *TETALOGY*

UTERINE: pertaining to the uterus (*uterus*)

VITELLIN: pertaining to the yolk (*vitellus*)

PREFACE

[BY JOSEPH MCCABE]

THE work which we now place within the reach of every reader of the English tongue is one of the finest productions of its distinguished author. The first edition appeared in 1874. At that time the conviction of man's natural evolution was even less advanced in Germany than in England, and the work raised a storm of controversy. Theologians—forgetting the commonest facts of our individual development—spoke with the most profound disdain of the theory that a Luther or a Goethe could be the outcome of development from a tiny speck of protoplasm. The work, one of the most distinguished of them said, was “a fleck of shame on the escutcheon of Germany.” To-day its conclusion is accepted by influential clerics, such as the Dean of Westminster, and by almost every biologist and anthropologist of distinction in Europe. Evolution is not a laboriously reached conclusion, but a guiding truth, in biological literature to-day.

There was ample evidence to substantiate the conclusion even in the first edition of the book. But fresh facts have come to light in each decade, always enforcing the general truth of man's evolution, and at times making clearer the line of development. Professor Haeckel embodied these in successive editions of his work. In the fifth edition, of which this is a translation, reference will be found to the very latest facts bearing on the evolution of man, such as the discovery of the remarkable effect of mixing human blood with that of the anthropoid ape. Moreover, the ample series of illustrations has been considerably improved and enlarged; there is no scientific work published, at a price remotely approaching that of the present edition, with so abundant and excellent a supply of illustrations. When it was issued in Germany, a few years ago, a distinguished biologist wrote in the *Frankfurter Zeitung* that it would secure immortality for its author, the most notable critic of the idea of immortality. And the *Daily Telegraph* reviewer described the English version as a “handsome edition of Haeckel's monumental work,” and “an issue worthy of the subject and the author.”

The influence of such a work, one of the most constructive that Haeckel has ever written, should extend to more than the few hundred readers who are able to purchase the expensive volumes of the original issue. Few pages in the story of science are more arresting and generally instructive than this great picture of “mankind in the making.” The horizon of the mind is healthily expanded as we follow the search-light of science down the vast avenues of past time, and gaze on the uncouth forms that enter

into, or illustrate, the line of our ancestry. And if the imagination recoils from the strange and remote figures that are lit up by our search-light, and hesitates to accept them as ancestral forms, science draws aside another veil and reveals another picture to us. It shows us that each of us passes, in our embryonic development, through a series of forms hardly less uncouth and unfamiliar. Nay, it traces a parallel between the two series of forms. It shows us man beginning his existence, in the ovary of the female infant, as a minute and simple speck of jelly-like plasm. It shows us (from analogy) the fertilised ovum breaking into a cluster of cohering cells, and folding and curving, until the limb-less, head-less, long-tailed *fœtus* looks like a worm-shaped body. It then points out how gill-slits and corresponding blood-vessels appear, as in a lowly fish, and the fin-like extremities bud out and grow into limbs, and so on; until, after a very clear ape-stage, the definite human form emerges from the series of transformations.

It is with this *embryological* evidence for our evolution that the present volume is concerned. There are illustrations in the work that will make the point clear at a glance. Possibly too clear; for the simplicity of the idea and the eagerness to apply it at every point have carried many, who borrow hastily from Haeckel, out of their scientific depth. Haeckel has never shared their errors, nor encouraged their superficiality. He insists from the outset that a complete parallel could not possibly be expected. Embryonic life itself is subject to evolution. Though there is a general and substantial law—as most of our English and American authorities admit—that the embryonic series of forms recalls the ancestral series of forms, the parallel is blurred throughout and often distorted. It is not the obvious resemblance of the embryos of different animals, and their general similarity to our extinct ancestors in this or that organ, on which we must rest our case. A careful study must be made of the various stages through which all embryos pass, and an effort made to prove their real identity and therefore genealogical relation.

This is a task of great subtlety and delicacy. Many scientists have worked at it together with Professor Haeckel—I need only name our own Professor Balfour and Professor Ray Lankester—and the scheme is fairly complete. But the general reader must not expect that even so clear a writer as Haeckel can describe these intricate processes without demanding his very careful attention. Most of the chapters in the present volume (and the second volume will be less difficult) are easily intelligible to all; but there are points at which the line of argument is necessarily subtle and complex. In the hope that most readers will be induced to master even these more difficult chapters, I will give an outline of the characteristic argument of the work. Haeckel's distinctive services in regard to man's evolution have been: (1) The construction of a complete ancestral tree, though, of course, some of the stages in it are purely conjectural, and not final; (2) The tracing of the remarkable reproduction of ancestral forms in

the embryonic development of the individual. Naturally, he has not worked alone in either department. The second volume of this work will embody the first of these two achievements; the present one is mainly concerned with the latter. It will be useful for the reader to have a synopsis of the argument and an explanation of some of the chief terms invented or employed by the author.

The main theme of the work is that, in the course of their embryonic development, all animals, including man, pass roughly and rapidly through a series of forms which represents the succession of their ancestors in the past. After a severe and extensive study of embryonic phenomena, Haeckel has drawn up a "law" (in the ordinary scientific sense) to this effect, and has called it "the biogenetic law," or the chief law relating to the evolution (*genesis*) of life (*bios*). This law is widely and increasingly accepted by embryologists and zoologists. It is enough to quote a recent declaration of the great American zoologist, President D. Starr Jordan: "It is, of course, true that the life-history of the individual is an epitome of the life-history of the race"; while a distinguished German zoologist (Sarasin) has described it as being of the same use to the biologist as spectrum analysis is to the astronomer.

But the reproduction of ancestral forms in the course of the embryonic development is by no means always clear, or even always present. Many of the embryonic phases do not recall ancestral stages at all. They may have done so originally, but we must remember that the embryonic life itself has been subject to adaptive changes for millions of years. All this is clearly explained by Professor Haeckel. For the moment, I would impress on the reader the vital importance of fixing the distinction from the start. He must thoroughly familiarise himself with the meaning of five terms. *Biogeny* is the development of life in general (both in the individual and the species), or the science describing it. *Ontogeny* is the development (embryonic and post-embryonic) of the individual (*on*), or the science describing it. *Phylogeny* is the development of the race or stem (*phylon*), or the science describing it. Roughly, *ontogeny* may be taken to mean embryology, and *phylogeny* what we generally call evolution. Further, the embryonic phenomena sometimes reproduce ancestral forms, and they are then called *palaogenetic* (from *pala* = again): sometimes they do not recall ancestral forms, but are later modifications due to adaptation, and they are then called *neogenetic* (from *neos* = new or foreign). These terms are now widely used, but the reader of Haeckel must understand them thoroughly.

The first five chapters are an easy account of the history of embryology and evolution. The sixth and seventh give an equally clear account of the sexual elements and the process of conception. But some of the succeeding chapters must deal with embryonic processes so unfamiliar, and pursue them through so wide a range of animals in a brief space,

that, in spite of the 200 illustrations, they will offer difficulty to many a reader. As our aim is to secure, not a superficial acquiescence in conclusions, but a fair comprehension of the truths of science, we have retained these chapters. However, I will give a brief and clear outline of the argument, so that the reader with little leisure may realize their value.

When the animal ovum (egg-cell) has been fertilised, it divides and sub-divides until we have a cluster of cohering cells, externally not unlike a raspberry or mulberry. This is the *morula* (= mulberry) stage. The cluster becomes hollow, or filled with fluid in the centre, all the cells rising to the surface. This is the *blastula* (hollow ball) stage. One half of the cluster then bends or folds in upon the other, as one might do with a thin Indianrubber ball, and we get a vase-shaped body with hollow interior (the first stomach, or "primitive gut"), an open mouth (the first or "primitive mouth"), and a wall composed of two layers of cells (two "germinal layers"). This is the *gastrula* (stomach) stage, and the process of its formation is called *gastrulation*. A glance at the illustration on p. 61 will make this perfectly clear.

So much for the embryonic process in itself. The application to evolution has been a long and laborious task. Briefly, it was necessary to show that *all* the multicellular animals passed through these three stages, so that our biogenetic law would enable us to recognise them as reminiscences of ancestral forms. This is the work of Chaps. VIII. and IX. The difficulty can be realised in this way: As we reach the higher animals the ovum has to take up a large quantity of yolk, on which it may feed in developing. Think of the bird's "egg." The effect of this was to flatten the germ (the *morula* and *blastula*) from the first, and so give, at first sight, a totally different complexion to what it has in the lowest animals. When we pass the reptile and bird stage, the large yolk almost disappears (the germ now being supplied with blood by the mother), but the germ has been permanently altered in shape, and there are now a number of new embryonic processes (membranes, blood-vessel connections, etc.). Thus it was no light task to trace the identity of this process of *gastrulation* in all the animals. It has been done, however; and with this introduction the reader will be able to follow the proof. The conclusion is important. If all animals pass through the curious *gastrula* stage, it must be because they all had a common ancestor of that nature. To this conjectural ancestor (it lived before the period of fossilisation begins) Haeckel gives the name of the *Gastrea*, and in the second volume we shall see a number of living animals of this type ("gastræans").

The line of argument is the same in the next chapter. After laborious and careful research (though this stage is not generally admitted in the same sense as the previous one), a fourth common stage was discovered, and given the name of the *Cœlomeia*. The *blastula* had one layer of cells, the *blastoderm* (*derma* = skin): the *gastrula* two layers, the *ectoderm* ("outer skin") and *entoderm* ("inner skin"). Now a third layer (*mesoderm*

the middle skin) is formed, by the growth inwards of two pouches or folds of the skin. The pouches blend together, and form a single cavity (the body cavity, or *cœlom*), and its two walls are two fresh "germinal layers." Again, the identity of the process has to be proved in all the higher classes of animals, and when this is done we have another ancestral stage, the *Cœlomata*.

The remaining task is to build up the complex frame of the higher animals—always showing the identity of the process (on which the evolutionary argument depends) in enormously different conditions of embryonic life—out of the four "germinal layers." Chap. IX. prepares us for the work by giving us a very clear account of the essential structure of the back-boned (vertebrate) animal, and the probable common ancestor of all the vertebrates (a small fish of the lancelet type). Chaps. XI.-XIV. then carry out the construction step by step. The work is now simpler, in the sense that we leave all the invertebrate animals out of account; but there are so many organs to be fashioned out of the four simple layers that the reader must proceed carefully. In the second volume each of these organs will be dealt with separately, and the parallel will be worked out between its embryonic and its phylogenetic (evolutionary) development. The general reader may wait for this for a full understanding. But in the meantime the wonderful story of the construction of all our organs in the course of a few weeks (the human frame is perfectly formed, though less than two inches in length, by the twelfth week) from so simple a material is full of interest. It would be useless to attempt to summarise the process. The four chapters are themselves but a summary of it, and the eighty fine illustrations of the process will make it sufficiently clear. The last chapter carries the story on to the point where man at last parts company with the anthropoid ape, and gives a full account of the membranes or wrappers that enfold him in the womb, and the connection with the mother.

In conclusion, I would urge the reader to consult, at his free library perhaps, the complete edition of this work, when he has read the present abbreviated edition. Much of the text has had to be condensed in order to bring out the work at our popular price, and the beautiful plates of the complete edition have had to be omitted. The reader will find it an immense assistance if he can consult the library edition.

JOSEPH MCCABE.


Crickham, March, 1906.

HAECKEL'S CLASSIFICATION OF THE ANIMAL WORLD

Unicellular animals (Protozoa)

1. Unnucleated { Bacteria
Protomonas } Monera
2. Nucleated { a. Rhizopoda { Amoebsina
Radiolarians
b. Infusoria { Flagellata
Ciliata
3. Cell-colonies { Catallacta
Blattula

Multicellular animals (Metazoa)



* This classification is given for the purpose of explaining Haeckel's use of terms in this volume. The general reader should bear in mind that it differs very considerably from more recent schemes of classification. He should compare the scheme framed by Professor E. Ray Lankester.

THE EVOLUTION OF MAN

CHAPTER I.

THE FUNDAMENTAL LAW OF ORGANIC EVOLUTION

ral phenomena into which I would introduce my readers in the following chapters has a quite peculiar place in the broad realm of scientific inquiry. There is no object of investigation that touches man more closely, and the knowledge of which should be more acceptable to him, than his own frame. But among all the various branches of the natural history of mankind, or *anthropology*, the story of his development by natural means must excite the most lively interest. It gives us the key of the great world-riddles at which the human mind has been working for thousands of years. The problem of the nature of man, or the question of man's place in nature, and the cognate inquiries as to the past, the earliest history, the present situation, and the future of humanity—all these most important questions are directly and intimately connected with that branch of study which we call the science of the evolution of man, or, in one word, "*Anthropogeny*" (the genesis of man). Yet it is an astonishing fact that the science of the evolution of man does not even yet form part of the scheme of general education. In fact, educated people even in our day are for the most part quite ignorant of the important truths and remarkable phenomena which anthropogeny teaches us.

As an illustration of this curious state of things, it may be pointed out that most of what are considered to be "educated" people do not know that every human being is developed from an egg, or ovum, and that this egg is one simple cell, like any other plant or animal egg. They are equally ignorant that in the course of the development of this tiny, round egg-cell there is first formed a body that is totally

different from the human frame, and has not the remotest resemblance to it. Most of them have never seen such a human embryo in the earlier period of its development, and do not know that it is quite indistinguishable from other animal embryos. At first the embryo is no more than a round cluster of cells, then it becomes a simple hollow sphere, the wall of which is composed of a layer of cells. Later it approaches very closely, at one period, to the anatomic structure of the lancelet, afterwards to that of a fish, and again to the typical build of the amphibian and mammal. As it continues to develop, a form appears which is like those we find at the lowest stage of mammal-life (such as the duck-bills), then a form that resembles the marsupials, and only at a late stage a form that has a resemblance to the ape, until at last the definite human form emerges and closes the series of transformations. These suggestive facts are, as I said, still almost unknown to the general public—so completely unknown that, if one casually mentions them, they are called in question or denied outright as fairy-tales. Everybody knows that the butterfly emerges from the pupa, and the pupa from a quite different thing called a larva, and the larva from the butterfly's egg. But few besides radical men are aware that man, in the course of his individual formation, passes through a series of transformations which are not less surprising and wonderful than the familiar metamorphoses of the butterfly.

The mere description of these remarkable changes through which man passes during his embryonic life should arouse considerable interest. But the mind will experience a far keener satisfaction when

we trace these curious facts to their causes, and when we learn to behold in them natural phenomena which are of the highest importance throughout the whole field of human knowledge. They throw light first of all on the "natural history of creation," then on psychology, or "the science of the soul," and through this on the whole of philosophy. And as the general results of every branch of inquiry are summed up in philosophy, all the sciences come in turn to be touched and influenced more or less by the study of the evolution of man.

But when I say that I propose to present here the most important features of these phenomena and trace them to their causes, I take the term, and I interpret my task, in a very much wider sense than is usual. The lectures which have been delivered on this subject in the universities during the last half-century are almost exclusively adapted to medical men. Certainly, the medical man has the greatest interest in studying the origin of the human body, with which he is daily occupied. But I must not give here this special description of the embryonic processes such as it has hitherto been given, as most of my readers have not studied anatomy, and are not likely to be entrusted with the care of the adult organism. I must content myself with giving some parts of the subject only in general outline, and must not enter upon all the marvellous, but very intricate and not easily described, details that are found in the story of the development of the human frame. To understand these fully a knowledge of anatomy is needed. I will endeavour to be as plain as possible in dealing with this branch of science. Indeed, a sufficient general idea of the course of the embryonic development of man can be obtained without going too closely into the anatomic details. I trust we may be able to arouse the same interest in this delicate field of inquiry as has been excited already in other branches of science; though we shall meet more obstacles here than elsewhere.

The story of the evolution of man, as it has hitherto been expounded to medical students, has usually been confined to embryology—more correctly, *ontogeny*—or the science of the development of the individual human organism. But this is really only the first part of our task, the first half of the story of the evolution of man in that wider sense in which we

understand it here. We must add as the second half—as another and not less important and interesting branch of the science of the evolution of the human stem—*phylogeny*, this may be described as the science of the evolution of the various animal forms from which the human organism has been developed in the course of countless ages. Everybody now knows of the great scientific activity that was roused by the publication of Darwin's *Origin of Species* in 1859. The chief direct consequence of this publication was to provoke a fresh inquiry into the origin of the human race, and this has proved beyond question our gradual evolution from the lower species. We give the name of "Phylogeny" to the science which describes this ascent of man from the lower ranks of the animal world. The chief sources that it draws upon for facts are "Ontogeny," or embryology, the science of the development of the individual organism. Moreover, it derives a good deal of support from *palæontology*, or the science of fossil remains, and even more from comparative anatomy, or *morphology*.

These two branches of our science—on the one side ontogeny or embryology, and on the other phylogeny, or the science of races or evolution—are most vitally connected. The one cannot be understood without the other. It is only when the two branches fully co-operate and supplement each other that "Biogeny" (or the science of the genesis of life in the widest sense) attains to the rank of a philosophic science. The connection between them is not external and superficial, but profound, intrinsic, and causal. This is a discovery made by recent research, and it is most clearly and correctly expressed in the comprehensive law which I have called "the fundamental law of organic evolution," or "the fundamental law of biogeny." This general law, to which we shall find ourselves constantly recurring, and on the recognition of which depends one's whole insight into the story of evolution, may be briefly expressed in the phrase: "The history of the fetus is a recapitulation of the history of the race"; or, in other words, "Ontogeny is a recapitulation of phylogeny." It may be more fully stated as follows: The series of forms through which the individual organism passes during its development from the ovum to the complete bodily structure is a brief, condensed repetition

of the long series of forms which the animal ancestors of the said organism, or the ancestral forms of the species, have passed through from the earliest period of organic life down to the present day.

The causal character of the relation which connects embryology with stem-history is due to the action of heredity and adaptation. When we have rightly understood these, and recognised their great importance in the formation of organisms, we can go a step further and say: Phylogenesis is the mechanical cause of ontogenesis.¹ In other words, the development of the stem, or race, is, in accordance with the laws of heredity and adaptation, the cause of all the changes which appear in a condensed form in the evolution of the species.

The chain of manifold animal forms which represent the ancestry of each higher organism, or even of man, according to the theory of descent, always form a connected whole. We may designate this uninterrupted series of forms with the letters of the alphabet: A, B, C, D, E, etc., to Z. In apparent contradiction to what I have said, the story of the development of the individual, or the ontogeny of most organisms, only offers to the observer a part of these forms, so that the defective series of embryonic forms would run: A, B, D, F, H, K, M, etc., or, in other cases, B, D, H, L, M, N, etc. Here, then, as a rule, several of the evolutionary forms of the original series have fallen out. Moreover, we often find—to continue with our illustration from the alphabet—one or other of the original letters of the ancestral series represented by corresponding letters from a different alphabet. Thus, instead of the Roman B and D, we often have the Greek β and δ . In this case the text of the biogenetic law has been corrupted, just as it had been abbreviated in the preceding case. But, in spite of all this, the series of ancestral forms remains the same, and we are in a position to discover its original complexion.

In reality, there is always a certain parallel between the two evolutionary series. But it is obscured from the fact

that in the embryonic succession much is wanting that certainly existed in the earlier ancestral succession. If the parallel of the two series were complete, and if this great fundamental law affirming the causal connection between ontogeny and phylogeny in the proper sense of the word were directly demonstrable, we should only have to determine, by means of the microscope and the dissecting knife, the series of forms through which the fertilised ovum passes in its development; we should then have before us a complete picture of the remarkable series of forms which our animal ancestors have successively assumed from the dawn of organic life down to the appearance of man. But such a repetition of the ancestral history by the individual in its embryonic life is very rarely complete. We do not often find our full alphabet. In most cases the correspondence is very imperfect, being greatly distorted and falsified by causes which we will consider later. We are thus, for the most part, unable to determine in detail, from the study of its embryology, all the different shapes which an organism's ancestors have assumed; we usually—and especially in the case of the human foetus—encounter many gaps. It is true that we can fill up most of these gaps satisfactorily with the help of comparative anatomy, but we cannot do so from direct embryological observation. Hence it is important that we find a large number of lower animal forms to be still represented in the course of man's embryonic development. In these cases we may draw our conclusions with the utmost security as to the nature of the ancestral form from the features of the form which the embryo momentarily assumes.

To give a few examples, we can infer from the fact that the human ovum is a simple cell, that the first ancestor of our species was a tiny unicellular being, something like the amoeba. In the same way, we know, from the fact that the human foetus consists, at the first, of two simple cell-layers (the *gastrula*), that the *gastrula*, a form with two such layers, was certainly in the line of our ancestry. A later human embryonic form (the *chorda*) points just as clearly to a worm-like ancestor (the *prochordone*), the nearest living relation of which is found among the actual ascidians. To this succeeds a most important embryonic stage (*acrania*), in which our headless forms

¹ The term "genesis" which occurs throughout, means, of course, "birth" or origin. From this we get: Biogeny = the origin of life (*bios*); Autogeny = the origin of this (*auto*); Ontogeny = the origin of the individual (*ontos*); Phylogeny = the origin of the species (*phylon*); and so on. In each case the term may refer to the species itself, or to the science describing the process, — I name.

presents, in the main, the structure of the lancelet. But we can only indirectly and approximately, with the aid of comparative anatomy and ontogeny, conjecture what lower forms enter into the chain of our ancestry between the gastrea and the chordula, and between this and the lancelet. In the course of the historical development many intermediate structures have gradually fallen out, which must certainly have been represented in our ancestry. But, in spite of these many, and sometimes very appreciable, gaps, there is no contradiction between the two successions. In fact, it is the chief purpose of this work to prove the real harmony and the original parallelism of the two. I hope to show, on a substantial basis of facts, that we can draw most important conclusions as to our genealogical tree from the actual and easily demonstrable series of embryonic changes. We shall then be in a position to form a general idea of the wealth of animal forms which have figured in the direct line of our ancestry in the lengthy history of organic life.

In this evolutionary appreciation of the facts of embryology we must, of course, take particular care to distinguish sharply and clearly between the primitive, palinogenetic (or ancestral) evolutionary processes and those due to cenogenesis.¹ By *palinogenetic* processes, or embryonic recapitulations, we understand all those phenomena in the development of the individual which are transmitted from one generation to another by heredity, and which, on that account, allow us to draw direct inferences as to corresponding structures in the development of the species. On the other hand, we give the name of *cenogenetic* processes, or embryonic variations, to all those phenomena in the fetal development that cannot be traced to inheritance from earlier species, but are due to the adaptation of the fetus, or the infant-form, to certain conditions of its embryonic development. These cenogenetic phenomena are foreign or later additions; they allow us to draw no direct inference whatever as to corresponding processes in our ancestral

history, but rather hinder us from doing so.

This careful discrimination between the primary or palinogenetic processes and the secondary or cenogenetic is of great importance for the purposes of the scientific history of a species, which has to draw conclusions from the available facts of embryology, comparative anatomy, and paleontology, as to the processes in the formation of the species in the remote past. It is of the same importance to the student of evolution as the careful distinction between genuine and spurious texts in the works of an ancient writer, or the purging of the real text from interpolations and alterations, is for the student of philology. It is true that this distinction has not yet been fully appreciated by many scientists. For my part, I regard it as the first condition for forming any just idea of the evolutionary process, and I believe that we must, in accordance with it, divide embryology into two sections — palinogenesis, or the science of recapitulated forms, and cenogenesis, or the science of supererogating structures.

To give at once a few examples from the science of man's origin in illustration of this important distinction, I may instance the following processes in the embryology of man, and of all the higher vertebrates, as *palinogenetic*: the formation of the two primary germinal layers and of the primitive gut, the undivided structure of the dorsal nerve-tube, the appearance of a simple slit and between the modulatory tube and the gut, the temporary formation of the gill-slits and arches, the primitive kidneys, and so on.² All these, and many other important structures, have clearly been transmitted by a steady heredity from the early ancestors of the mammal, and are, therefore, direct indications of the presence of similar structures in the history of the stem. On the other hand, this is certainly not the case with the following embryonic forms, which we must describe as *cenogenetic* processes: the formation of the yolk-sac, the allantois, the placenta, the amnion, the septulmenon, and the chorion—or, generally speaking, the various fetal membranes and the corresponding changes in the blood vessels. Further instances are: the dual structure of the heart cavity, the temporary division of the plates of the primitive vertebrae and

¹ *Palinogenesis* = new birth, or re-formation (*palin* = again, genuine or genuine — development), hence its application to the phenomena which are recapitulated by heredity from earlier ancestral forms. *Cenogenesis* = forming or superadded development (*ken* = and *genesis*); hence, those phenomena which exist later in the story of life to displace the inherited structure by a fresh adaptation to environment.—THOM.

² All these, and the following structures, will be fully described in later chapters.—THOM.

lateral plates, the secondary closing of the ventral and intestinal walls, the formation of the navel, and so on. All these and many other phenomena are certainly not traceable to similar structures in any earlier and completely-developed ancestral form, but have arisen simply by adaptation to the peculiar conditions of embryonic life (within the fetal membranes). In view of these facts, we may now give the following more precise expression to our chief law of biogeny.—The evolution of the fetus (or *ontogenesis*) is a condensed and abbreviated recapitulation of the evolution of the stem (or *phylogenesis*); and this recapitulation is the more complete in proportion as the original development (or *palingenesis*) is preserved by a constant heredity, on the other hand, it becomes less complete in proportion as a varying adaptation to new conditions increases the disturbing factors in the development (or *anagenesis*).

The congenetic alterations or distortions of the original palingenetic course of development take the form, as a rule, of a gradual displacement of the phenomena, which is slowly effected by adaptation to the changed conditions of embryonic existence during the course of thousands of years. This displacement may take place as regards either the position or the time of a phenomenon.

The great importance and strict regularity of the times-variations in embryology have been carefully studied recently by Ernest Haeckel, in his *Biogenetik* (Jena, 1888). He contends that our biogenetic Law has not been impaired by the attacks of its opponents, and goes on to say: "Surely any piece of knowledge has contributed so much to the advance of embryology as this, its formulation as one of the most signal services to general biology. It was not until this law passed into the flesh and blood of investigators, and they had accustomed themselves to see a reminiscence of ancestral history in embryonic structures, that we witnessed the great progress which embryological research has made in the last two decades." The best proof of the correctness of this opinion is that now the most fruitful work is done in all branches of embryology with the aid of this biogenetic Law, and that it enables students to attain every year thousands of brilliant results that they would never have reached without it.

It is only when one appreciates the congenetic processes in relation to the

palingenetic, and when one takes careful account of the changes which the latter may suffer from the former, that the radical importance of the biogenetic law is recognised, and it is felt to be the most illuminating principle in the science of evolution. In this task of discrimination it is the silver thread in relation to which we can arrange all the phenomena of this realm of marvels—the "Ariadne thread," which alone enables us to find our way through this labyrinth of forms. Hence the brothers Sarasin, the zoologists, could say with perfect justice, in their study of the evolution of the *Ichthyophis*, that "the great biogenetic law is just as important for the zoologist in tracing long-extinct processes as spectrum analysis is for the astronomer."

Even at an earlier period, when a correct acquaintance with the evolution of the human and animal frame was only just being obtained—and that is scarcely eighty years ago!—the greatest astonishment was felt at the remarkable similarity observed between the embryonic forms, or stages of fetal development, in very different animals; attention was called even then to their close resemblance to certain fully-developed animal forms belonging to some of the lower groups. The older scientists (Oken, Treviranus, and others) knew perfectly well that these lower forms in a sense illustrated and fixed, in the hierarchy of the animal world, a temporary stage in the evolution of higher forms. The famous anatomist Meckel spoke in 1821 of a "similarity between the development of the embryo and the series of animals." Boer raised the question in 1828 how far, within the vertebrate type, the embryonic forms of the higher animals assume the permanent shapes of members of lower groups. But it was impossible fully to understand and appreciate this remarkable resemblance at that time. We owe our capacity to do this to the theory of descent; it is this that puts in their true light the action of *heredity* on the one hand and *adaptation* on the other. It explains to us the vital importance of their constant reciprocal action in the production of organic forms. Darwin was the first to teach us the great part that was played in this by the ceaseless struggle for existence between living things, and to show how, under the influence of this (by natural selection), new species were produced and maintained solely by the interaction of heredity and

adaptation. It was thus Darwinism that first opened our eyes to a true comprehension of the supremely important relations between the two parts of the science of organic evolution—Ontogeny and Phylogeny.

Heredity and adaptation are, in fact, the two constructive physiological functions of living things; unless we understand these properly we can make no headway in the study of evolution. Hence, until the time of Darwin no one had a clear idea of the real nature and causes of embryonic development. It was impossible to explain the curious series of forms through which the human embryo passed; it was quite unintelligible why this strange succession of animal-like forms appeared in the series at all. It had previously been generally assumed that the man was found complete in all his parts in the ovum, and that the development consisted only in an unfolding of the various parts, a simple process of growth. This is by no means the case. On the contrary, the whole process of the development of the individual presents to the observer a connected succession of different animal-forms; and these forms display a great variety of external and internal structure. But *why* each individual human being should pass through this series of forms in the course of his embryonic development it was quite impossible to say until Lamarck and Darwin established the theory of descent. Through this theory we have at last detected the real causes, the *efficient causes*, of the individual development; we have learned that these *mechanical causes* suffice of themselves to effect the formation of the organism, and that there is no need of the *final causes* which were formerly assumed. It is true that in the academic philosophies of our time these final causes still figure very prominently; in the new philosophy of nature we can entirely replace them by *efficient causes*. We shall see, in the course of our inquiry, how the most wonderful and hitherto insoluble enigmas in the human and animal frame have proved amenable to a mechanical explanation, by causes acting without provision, through Darwin's reform of the science of evolution. We have every where been able to substitute unconscious causes, acting from necessity, for conscious, purposive causes.¹

If the new science of evolution had done no more than this, every thoughtful man would have to admit that it had accomplished an immense advance in knowledge. It means that in the whole of philosophy that tendency which we call monistic, in opposition to the dualistic, which has hitherto prevailed, must be accepted.² At this point the science of human evolution has a direct and profound bearing on the foundations of philosophy. Modern anthropology has, by its astounding discoveries during the second half of the nineteenth century, compelled us to take a completely monistic view of life. Our bodily structure and its life, our embryonic development and our evolution as a species, teach us that the same laws of nature rule in the life of man as in the rest of the universe. For this reason, if for no others, it is desirable, nay, indispensable, that every man who wishes to form a serious and philosophic view of life, and, above all, the expert philosopher, should acquaint himself with the chief facts of this branch of science.

The facts of embryology have so great and obvious a significance in this connection that even in recent years dualist and teleological philosophers have tried to rid themselves of them by simply denying them. This was done, for instance, as regards the fact that man is developed from an egg, and that this egg or ovum is a simple cell, as in the case of other animals. When I had explained this pregnant fact and its significance in my *History of Creation*, it was described in many of the theological journals as a dishonest invention of my own. The fact that the embryos of man and the dog are, at a certain stage of their development, almost indistinguishable was also denied. When we examine the human embryo in the third or fourth week of its development, we find it to be quite different in shape and structure from the full-grown human being, but almost identical with that of the ape, the dog, the rabbit, and

as well as in inorganic changes. On the other hand, the dualist or vitalist philosophy of nature affirms that unconscious forces are only at work in the inorganic world, and that we find conscious, purposive, or final causes in organic nature.

¹ Monism is neither purely materialistic nor purely spiritualistic, but a reconciliation of these two principles, since it regards the whole of nature as one, and sees only different phases of work in it. Dualism, on the contrary, holds that nature and spirit, matter and form, the world and God, inorganic and organic nature, are separate and independent substances. Cf. *The Reality of the Universe*, chap. ix.

¹ The dualist or mechanical philosophy of nature holds that only unconscious, purposive, efficient causes are at work in the whole field of nature, in organic life

other mammals, at the same stage of ontogeny. We find a bean-shaped body of very simple construction, with a tail below and a pair of fins at the sides, something like those of a fish, but very different from the limbs of man and the mammals. Nearly the whole front half of the body is taken up by a shapeless head without face, at the sides of which we find gill-clefts and arches as in the fish. At this stage of its development the human embryo does not differ in any essential detail from that of the ape, dog, horse, ox, etc., at a corresponding period. This important fact can easily be verified at any moment by a comparison of the embryos of man, the dog, rabbit, etc. Nevertheless, the theologians and dualist philosophers pronounced it to be a materialistic invention; even scientists, to whom the facts should be known, have sought to deny them.

There could not be a clearer proof of the profound importance of these embryological facts in favour of the monistic philosophy than is afforded by these efforts of its opponents to get rid of them by silence or denial. The truth is that these facts are most inconvenient for them, and are quite irreconcilable with their views. We must be all the more pressing on our side to put them in their proper light. I fully agree with Huxley when he says, in his *Man's Place in Nature*: "Though these facts are ignored by several well-known popular leaders, they are easy to prove, and are accepted by all scientific men; on the other hand, their importance is so great that those who have once mastered them will, in my opinion, find few other biological discoveries to astonish them."

We shall make it our chief task to study the evolution of man's bodily frame and its various organs in their external form and internal structures. But I may observe at once that this is accompanied step by step with a study of the evolution of their functions. These two branches of inquiry are inseparably united in the whole of anthropology, just as in zoology (of which the former is only a section) or general biology. Everywhere the peculiar form of the organism and its structures, internal and external, is directly related to the special physiological functions which the organism or organ has to execute. This intimate connection of structure and function, or of the instrument and the work done by it, is seen in the

science of evolution and all its parts. Hence the story of the evolution of structures, which is our immediate concern, is also the history of the development of functions; and this holds good of the human organism as of any other.

At the same time, I must admit that our knowledge of the evolution of functions is very far from being as complete as our acquaintance with the evolution of structures. One might say, in fact, that the whole science of evolution has almost confined itself to the study of structures; the evolution of functions hardly exists even in name. That is the fault of the physiologists, who have as yet concerned themselves very little about evolution. It is only in recent times that physiologists like W. Engelmann, W. Preyer, M. Verworn, and a few others, have attacked the evolution of functions.

It will be the task of some future physiologist to engage in the study of the evolution of functions with the same zeal and success as has been done for the evolution of structures in morphogeny (the science of the genesis of forms). Let me illustrate the close connection of the two by a couple of examples. The heart in the human embryo has at first a very simple construction, such as we find in permanent form among the ascidians and other low organisms; with this is associated a very simple system of circulation of the blood. Now, when we find that with the full-grown heart there comes a totally different and much more intricate circulation, our inquiry into the development of the heart becomes at once, not only an anatomical, but also a physiological, study. Thus it is clear that the ontogeny of the heart can only be understood in the light of its phylogeny (or development in the past), both as regards function and structure. The same holds true of all the other organs and their functions. For instance, the science of the evolution of the alimentary canal, the lungs, or the sexual organs, gives us at the same time, through the exact comparative investigation of structure-development, most important information with regard to the evolution of the functions of these organs.

This significant connection is very clearly seen in the evolution of the nervous system. This system is in the economy of the human body the medium of sensation, will, and even thought, the highest of the psychic functions; in a word, of

all the various functions which constitute the proper object of psychology. Modern anatomy and physiology have proved that these psychic functions are immediately dependent on the fine structure and the composition of the central nervous system, or the internal texture of the brain and spinal cord. In these we find the elaborate cell-machinery, of which the psychic or soul-life is the physiological function. It is so intricate that most men still look upon the mind as something supernatural that cannot be explained on mechanical principles.

But embryological research into the gradual appearance and the formation of this important system of organs yields the most astounding and significant results. The first sketch of a central nervous system in the human embryo presents the same very simple type as in the other vertebrates. A spinal tube is formed in the external skin of the back, and from this first comes a simple spinal cord without brain, such as we find to be the permanent psychic organ in the lowest type of vertebrate, the amphioxus. Not until a later stage is a brain formed at the anterior end of this cord, and then it is a brain of the most rudimentary kind, such as we find permanently among the lower fishes. This simple brain develops step by step, successively assuming forms which correspond to those of the amphibia, the reptiles, the duck-bills, and the lemurs. Only in the last stage does it reach the highly organized form which distinguishes the apes from the other vertebrates, and which attains its full development in man.

Comparative physiology discovers a precisely similar growth. The function of the brain, the psychic activity, rises step by step with the advancing development of its structure.

Thus we are enabled, by this story of the evolution of the nervous system, to understand at length the *maternal development of the human mind* and its gradual unfolding. It is only with the aid of embryology that we can grasp how these highest and most striking faculties of the animal organism have been historically evolved. In other words, a knowledge of the evolution of the spinal cord and brain in the human embryo leads us directly to a comprehension of the historic development (or phylogeny) of the human mind, that highest of all faculties, which

and supernatural in the adult man. This is certainly one of the greatest and most pregnant results of evolutionary science. Happily our embryological knowledge of man's central nervous system is now so adequate, and agrees so thoroughly with the complementary results of comparative anatomy and physiology, that we are thus enabled to obtain a clear insight into one of the highest problems of philosophy, the phylogeny of the soul, or the anecdotal history of the mind of man. Our chief support in this comes from the embryological study of it, or the ontogeny of the soul. This important section of psychology owes its origin especially to W. Preyer, in his interesting works, such as *The Mind of the Child*, *The Biography of a Baby* (1900), of Millicent Washburn Shinn, also deserves mention [See also Preyer's *Mental Development in the Child* (translation), and Sully's *Studies of Childhood and Children's Writings*].

In this way we follow the only path along which we may hope to reach the solution of this difficult problem.

Thirty-six years have now elapsed since, in my *General Morphology*, I established phylogeny as an independent science and showed its intimate causal connection with ontogeny; thirty years have passed since I gave in my *Gastrea-theory* the proof of the justice of this, and completed it with the theory of germinal layers. When we look back on this period we may ask, What has been accomplished during it by the fundamental law of biogeny? If we are unpartial, we must reply that it has proved its fertility in hundreds of sound results, and that by its aid we have acquired a vast fund of knowledge which we should never have obtained without it.

There has been no dearth of attacks—often violent attacks—on my conception of an intimate causal connection between ontogeny and phylogeny; but no other satisfactory explanation of these important phenomena has yet been offered to us. I say this especially with regard to Wilhelm His's theory of a "mechanical evolution," which questions the truth of phylogeny generally, and would explain the complicated embryonic processes without going beyond by simple physical changes—such as the bending and folding of leaves by electricity, the origin of cavities through unequal strain of the tissues, the formation of processes

fact is that these embryological phenomena themselves demand explanation in turn, and this can only be found, as a rule, in the corresponding changes in the

long ancestral series, or in the physiological functions of heredity and adaptation.

CHAPTER II.

THE OLDER EMBRYOLOGY

It is in many ways useful, on entering upon the study of any science, to cast a glance at its historical development. The saying that "everything is best understood in its growth" has a distinct application to science. While we follow its gradual development we get a clearer insight into its aims and objects. Moreover, we shall see that the present condition of the science of human evolution, with all its characteristics, can only be rightly understood when we examine its historical growth. This task will, however, not detain us long. The study of man's evolution is one of the latest branches of natural science, whether you consider the embryological or the phylogenetic section of it.

Apart from the few germs of our science which we find in classical antiquity, and which we shall notice presently, we may say that it takes its definite rise, as a science, at the year 1759, when one of the greatest German scientists, Caspar Friedrich Wolff, published his *Phorm generis hominis*. That was the foundation-stone of the science of animal embryology. It was not until fifty years later, in 1809, that Jean Lamarck published his *Philosophie Zoologique*—the first effort to provide a base for the theory of evolution; and it was another half-century before Darwin's work appeared (in 1859), which we may regard as the first scientific attainment of this aim. But before we go further into this solid establishment of evolution, we must cast a brief glance at that famous philosopher and scientist of antiquity, who stood alone in this, as in many other branches of science, for more than 2,000 years: the "father of natural history," Aristotle.

The extant scientific works of Aristotle deal with many different sides of biological research, the most comprehensive of them is his famous *History of Animals*. But not less interesting is the smaller work, *On the Generation of Animals* (*Peri soon genesear*). This work treats especially of embryonic development, and it is of great interest as being the earliest of its kind and the only one that has come down to us in any completeness from classical antiquity.

Aristotle studied embryological questions in various classes of animals, and among the lower groups he learned many most remarkable facts which we only rediscovered between 1830 and 1860. It is certain, for instance, that he was acquainted with the very peculiar mode of propagation of the cuttle-fishes, or cephalopoda, in which a yolk-sac hangs out of the mouth of the foetus. He knew, also, that embryos come from the eggs of the bee even when they have not been fertilised. This "parthenogenesis" (or virgin-birth) of the bees has only been established in our time by the distinguished zoologist of Munich, Siebold. He discovered that male bees come from the unfertilised, and female bees only from the fertilised, eggs. Aristotle further states that some kinds of fishes (of the genus *Serranus*) are hermaphrodites, each individual having both male and female organs and being able to fertilise itself; this, also, has been recently confirmed. He knew that the embryo of many fishes of the shark family is attached to the mother's body by a sort of placenta, or nutritive organ very rich in blood; apart from these, such an arrangement is only found among the higher mammals and

man. This placenta of the shark was looked upon as legendary for a long time, until Johannes Müller proved it to be a fact in 1839. Thus a number of remarkable discoveries were found in Aristotle's embryological work, proving a very good acquaintance of the great scientist—possibly helped by his predecessors—with the facts of ontogeny, and a great advance upon succeeding generations in that respect.

In the case of most of these discoveries he did not merely describe the fact, but added a number of observations on its significance. Some of those theoretical remarks are of particular interest, because they show a correct appreciation of the nature of the embryonic processes. He conceives the development of the individual as a new formation, in the course of which the various parts of the body take shape successively. When the human or animal form is developed in the mother's body, or separately in an egg, the heart—which he regards as the starting-point and centre of the organism—must appear first. Once the heart is formed the other organs arise, the internal ones before the external, the upper (those above the diaphragm) before the lower (or those beneath the diaphragm). The brain is formed at an early stage, and the eyes grow out of it. These observations are quite correct. And, if we try to form some idea from these data of Aristotle's general conception of the embryonic process, we find a dim premonition of the theory which Wolff showed 2,000 years afterwards to be the correct view. It is significant, for instance, that Aristotle denied the eternity of the individual in any respect. He said that the species or genus, the group of similar individuals, might be eternal, but the individual itself is temporary. It comes into being in the act of procreation, and passes away at death.

During the 2,000 years after Aristotle no progress whatever was made in general zoology, or in embryology in particular. People were content to read, copy, translate, and comment on Aristotle. Scarcely a single independent effort at research was made in the whole of the period. During the Middle Ages the spread of strong religious beliefs put formidable obstacles in the way of independent scientific investigation. There was no question of resuming the advance of biology. Even when human anatomy began to stir itself

independent research was resumed into the structure of the developed body, anatomists did not dare to extend their inquiries to the unformed body, the embryo, and its development. There were many reasons for the prevailing horror of such studies. It is natural enough, when we remember that a Bull of Boniface VIII. excommunicated every man who ventured to dissect a human corpse. If the dissection of a developed body were a crime to be thus punished, how much more dreadful must it have seemed to deal with the embryonic body still enclosed in the womb, which the Creator himself had decently veiled from the curiosity of the scientist! The Christian Church, then putting many thousands to death for unbelief, had a shrewd presentiment of the menace that science contained against its authority. It was powerful enough to see that its rival did not grow too quickly.

It was not until the Reformation broke the power of the Church, and a refreshing breath of the spirit dissolved the icy chains that bound science, that anatomy and embryology, and all the other branches of research, could begin to advance once more. However, embryology lagged far behind anatomy. The first works on embryology appear at the beginning of the sixteenth century. The Italian anatomist, Fabrizio ab Aquapendente, a professor at Padua, opened the advance. In his two books (*De formato feto*, 1600, and *De formatione fetus*, 1604) he published the older illustrations and descriptions of the embryos of man and other mammals, and of the hen. Similar imperfect illustrations were given by Spiegelius (*De formato feto*, 1631), and by Needham (1697) and his more famous compatriot, Harvey (1652), who discovered the circulation of the blood in the animal body and formulated the important principle, *Omnis vivens ex ovo* (all life comes from pre-existing life). The Dutch scientist, Swammerdam, published in his *Bible of Nature* the earliest observations on the embryology of the frog and the division of its egg-yell. But the most important embryological studies in the sixteenth century were those of the famous Italian, Marcello Malpighi, of Bologna, who led the way both in zoology and botany. His treatises, *De formatione pulli* and *De ovo incubato* (1673), contain the first consistent description of the development of the chick in the fertilized

Here I ought to say a word about the important part played by the chick in the growth of our science. The development of the chick, like that of the young of all other birds, agrees in all its main features with that of the other chief vertebrates, and even of man. The three highest classes of vertebrates—mammals, birds, and reptiles (lizards, serpents, tortoises, etc.)—have from the beginning of their embryonic development so striking a resemblance in all the chief points of structure, and especially in their first forms, that for a long time it is impossible to distinguish between them. We have known now for some time that we need only examine the embryo of a bird, which is the easiest to get at, in order to learn the typical mode of development of a mammal (and therefore of man). As soon as scientists began to study the human embryo, or the mammal-embryo generally, in its earlier stages about the middle and end of the seventeenth century, this important fact was very quickly discovered. It is both theoretically and practically of great value. As regards the theory of evolution, we can draw the most weighty inferences from the similarity between the embryos of widely different classes of animals. But for the practical purposes of embryological research the discovery is invaluable, because we can fill up the gaps in our imperfect knowledge of the embryology of the mammals from the more thoroughly studied embryology of the bird. Hen's eggs are easily to be had in any quantity, and the development of the chick may be followed step by step in artificial incubation. The development of the mammal is much more difficult to follow, because here the embryo is not detached and enclosed in a large egg, but the tiny ovum remains in the womb until the growth is completed. Hence, it is very difficult to keep up sustained observation of the various stages in any great extent, quite apart from such extrinsic considerations as the cost, the technical difficulties, and many other obstacles which we encounter when we would make an extensive study of the fertilized mammal. The chicken has, therefore, always been the chief object of study in this connection. The excellent incubators we now have enable us to observe it in any quantity and at any stage of development, and so follow the whole course of its formation step by step.

By the end of the seventeenth century Malpighi had advanced as far as it was possible to do with the imperfect microscope of his time in the embryological study of the chick. Further progress was arrested until the instrument and the technical methods should be improved. The vertebrate embryos are so small and delicate in their earlier stages that you cannot go very far into the study of them without a good microscope and other technical aid. But this substantial improvement of the microscope and the other apparatus did not take place until the beginning of the nineteenth century.

Embryology made scarcely any advance in the first half of the eighteenth century, when the systematic natural history of plants and animals received so great an impulse through the publication of Linné's famous *Systema Naturæ*. Not until 1759 did the genius arise who was to give it an entirely new character, Caspar Friedrich Wolff. Until then embryology had been occupied almost exclusively in unfortunate and misleading efforts to build up theories on the imperfect empirical material then available.

The theory which then prevailed, and remained in favour throughout nearly the whole of the eighteenth century, was commonly called at that time "the evolution theory"; it is better to describe it as "the preformation theory." Its chief point is this: There is no new formation of structures in the embryonic development of any organism, animal or plant, or even of man; there is only a growth, or unfolding, of parts which have been constructed or pre-formed from all eternity, though on a very small scale and closely packed together. Hence, every living germ contains all the organs and parts of the body, in the form and arrangement they will present later, already within it, and thus the whole embryological process is merely an *evolution* in the literal sense of the word, or an *unfolding*, of parts that were pre-formed and folded up in it. So, for instance, we find in the hen's egg not merely a simple cell, that divides and subdivides and forms germinal layers, and at last, after all kinds of variation and cleavage and reconstruction, brings forth

* This theory is usually known as the "evolution theory" in Germany, in contradistinction to the "epigenesis theory." But as it is the latter that is called the "evolution theory" in England, France, and Italy, and "epigenesis" and "epigenism," etc. (able to be synonymous, it seems better to call the first the "preformation theory."

the body of the chick; but there is in every egg from the first a complete chicken, with all its parts made and neatly packed. These parts are so small or so transparent that the microscope cannot detect them. In the hatching, these parts merely grow larger, and spread out in the normal way.

When this theory is consistently developed it becomes a "scatulation theory." According to its teaching, there was made in the beginning one couple or one individual of each species of animal or plant, but this one individual contained the germs of all the other individuals of the same species who should ever come to life. As the age of the earth was generally believed at that time to be fixed by the Bible at 5,000 or 6,000 years, it seemed possible to calculate how many individuals of each species had lived in the period, and so had been packed inside the first being that was created. The theory was consistently extended to man, and it was affirmed that our common parent Eve had had stored in her ovary the germs of all the children of men.

The theory at first took the form of a belief that it was the *female* who were thus encased in the first being. One couple of each species was created, but the female contained in her ovary all the future individuals of the species, of either sex. However, this had to be altered when the Dutch microscopist, Leeuwenhoek, discovered the male spermatozoa in 1690, and showed that an immense number of these extremely fine and mobile thread-like beings exist in the male sperm (this will be explained in the seventh chapter). This astonishing discovery was further advanced when it was proved that these living bodies, swarming about in the seminal fluid, were real animalcules, and, in fact, were the pre-formed germs of the future generation. When the male and female procreative elements came together at conception, these thread-like spermatozoa ("seed-animal") were supposed to penetrate into the fertile body of the ovum and begin to develop there, as the plant seed does in the fruitful earth. Hence, every spermatozoon was regarded as a *homunculus*, a tiny complete man, all the parts were believed to be pre-formed in it, and merely grew larger when it reached its proper medium in the female

ovum. This theory, also, was consistently developed in the sense that in each of these thread-like bodies the whole of its posterity was supposed to be present in the minutest form. Adam's sexual glands were thought to have contained the germs of the whole of humanity.

This "theory of male scatulation" found itself at once in keen opposition to the prevailing "female" theory. The two rival theories at once opened a very lively campaign, and the physiologists of the eighteenth century were divided into two great camps—the Annalists and the Ovulists—which fought vigorously. The annalists held that the spermatozoa were the true germs, and appealed to the lively movements and the structure of these bodies. The opposing party of the Ovulists, who clung to the older "evolution theory," affirmed that the ovum is the real germ, and that the spermatozoa merely stimulate it at conception to begin its growth, all the future generations are stored in the ovum. This view was held by the great majority of the biologists of the eighteenth century, in spite of the fact that Wolff proved it in 1759 to be without foundation. It used its prestige chiefly to the circumstance that the most weighty authorities in the biology and philosophy of the day decided in favour of it, especially Haller, Bonnet, and Lestibaud.

Albrecht Haller, professor at Göttingen, who is often called the father of physiology, was a man of wide and varied learning, but he does not occupy a very high position in regard to insight into natural phenomena. He made a vigorous defence of the "evolution theory" in his famous work, *Elementa physiologicæ*, affirming: "There is no such thing as formation (*nulla est epigenesis*). No part of the animal frame is made before another, all were made together." He thus denied that there was any evolution in the proper sense of the word, and even went so far as to say that the heart existed in the new-born child and the antlers in the hornless fawn, all the parts were there in advance, and were merely hidden from the eye of man for the time being. Haller even calculated the number of human beings that God must have created on the sixth day and stored away in Eve's ovary. He put the number at 300,000 millions, assuming the age of the world to be 5,000 years, the average age of a human being to be thirty years, and the population of the world at

* "Fading theory" would be the literal translation. *Scatula* in the Latin for a case or box.—TRANS.

that time to be 1,000 millions. And the famous Haller maintained all this nonsense, in spite of its ridiculous consequences, even after Wolff had discovered the real course of embryonic development and established it by direct observation!

Among the philosophers of the time the distinguished Leibnitz was the chief defender of the "preformation theory," and by his authority and literary prestige won many adherents to it. Supported by his system of monads, according to which body and soul are united in inseparable association and by their union form the individual, or the "monad," Leibnitz consistently extended the "scalutulum theory" to the soul, and held that this was no more evolved than the body. He says, for instance, in his *Philosophy*: "I mean that these souls, which one day are to be the souls of men, are present in the seed, like those of other species; in such wise that they existed in our ancestors as far back as Adam, or from the beginning of the world, in the form of organized bodies."

The theory seemed to receive considerable support from the observations of one of its most zealous supporters, Bonnet. In 1745 he discovered, in the plant-house, a case of parthenogenesis, or virgin-birth, an interesting form of reproduction that has lately been found by Siebold and others among various classes of the articulate, especially crustacea and insects. Among these and other animals of certain lower species the female may reproduce for several generations without having been fertilized by the male. These ova that do not need fertilization are called "false ova," pseudova or spores. Bonnet saw that a female plant-house, which he had kept in clostral isolation, and rigidly removed from contact with males, had on the eleventh day (after forming a new skin for the fourth time) a living daughter, and during the next twenty days ninety-four other daughters, and that all of them went on to reproduce in the same way without any contact with males. It seemed as if this furnished an irrefutable proof of the truth of the scalutulum theory, as it was held by the Ovidians; it is not surprising to find that the theory then secured general acceptance.

This was the condition of things when suddenly, in 1759, Caspar Friedrich Wolff adduced, and dealt a fatal blow at the whole preformation theory with his new theory of epigenesis. Wolff the son of a

Berlin tailor, was born in 1733, and went through his scientific and medical studies, first at Berlin under the famous anatomist Meckel, and afterwards at Halle. Here he secured his doctorate in his twenty-sixth year, and in his academic dissertation (November 28th, 1759), the *Theoria generionis*, expounded the new theory of a real development on a basis of epigenesis. This treatise is, in spite of its smallness and its obscure phraseology, one of the most valuable in the whole range of biological literature. It is equally distinguished for the mass of new and careful observations it contains, and the far-reaching and pregnant ideas which the author everywhere extracts from his observations and builds into a luminous and accurate theory of generation. Nevertheless, it met with no success at the time. Although scientific studies were then assiduously cultivated owing to the impulse given by Linné—although botanists and zoologists were no longer counted by dozens, but by hundreds, hardly any notice was taken of Wolff's theory. Even when he established the truth of epigenesis by the most rigorous observations, and demolished the airy structure of the preformation theory, the "exact" scientist Haller proved one of the most strenuous supporters of the old theory, and rejected Wolff's correct view with a dictatorial "There is no such thing as evolution." He even went on to say that religion was menaced by the new theory! It is not surprising that the whole of the physiologists of the second half of the eighteenth century submitted to the ruling of this physiological pontiff, and attacked the theory of epigenesis as a dangerous innovation. It was not until more than fifty years afterwards that Wolff's work was appreciated. Only when Meckel translated into German in 1812 another valuable work of Wolff's on *The Formation of the Illimentary Canal* (written in 1768), and called attention to its great importance, did people begin to think of him once more; yet this obscure writer had evinced a profounder insight into the nature of the living organism than any other scientist of the eighteenth century.

Wolff's idea led to an appreciable advance over the whole field of biology. There is such a vast number of new and important observations and pregnant thoughts in his writings that we have only gradually learned to appreciate them fully in the course of the nineteenth

century. He opened up the true path for research in many directions. In the first place, his theory of epigenesis gave us the first real insight into the nature of embryonic development. He showed convincingly that the development of every organism consists of a series of *new formations*, and that there is no trace whatever of the complete form either in the ovum or the spermatozoon. On the contrary, these are quite simple bodies, with a very different purport. The embryo which is developed from them is also quite different, in its internal arrangement and outer configuration, from the complete organism. There is no trace whatever of preformation or in-folding of organs. To-day we can scarcely call epigenesis a *theory*, because we are convinced it is a fact, and can demonstrate it at any moment with the aid of the microscope.

Wolff furnished the conclusive empirical proof of his theory in his classic dissertation on *The Formation of the Alimentary Canal* (1766). In its complete state the alimentary canal of the hen is a long and complex tube, with which the lungs, liver, salivary glands, and many other small glands, are connected. Wolff showed that in the early stages of the embryonic chick there is no trace whatever of this complicated tube with all its dependencies, but instead of it only a flat, leaf-shaped body; that, in fact, the whole embryo has at first the appearance of a flat, oval-shaped leaf. When we remember how difficult the exact observation of so fine and delicate a structure as the early leaf-shaped body of the chick must have been with the poor microscopes then in use, we must admire the

Wolff to make the most important discoveries in this most difficult part of embryology. By his laborious research he correct opinion that the embryonic body of all the higher animals, such as the birds, is for some time merely

a flat, thin, leaf-shaped disk—consisting at first of one layer, but afterwards of several. The lowest of these layers is the embryonic canal, and Wolff followed its development from its commencement to its completion. He showed how this leaf-shaped structure first turns into a groove, then the margins of this groove fold together and form a closed canal, and at length the two external openings of the tube (the mouth and anus) appear.

Moreover, the important fact that the other systems of organs are developed in the same way, from tubes formed out of simple layers, did not escape Wolff. The nervous system, muscular system, and vascular (blood-vessel) system, with all the organs appertaining (heart, etc., like the alimentary system, developed out of simple leaf-shaped structures. Hence, Wolff came to the view by 1768 which Pander developed in the *Theory of Germinal Layers* fifty years afterwards. His principles are not literally correct; but he comes as near to the truth in them as was possible at that time, and could be expected of him.

Our admiration of this gifted genius increases when we find that he was also the precursor of Goethe in regard to the metamorphosis of plants and of the famous cellular theory. Wolff had, as Huxley showed, a clear presentiment of this cardinal theory, since he recognised small microscopic globules as the elementary parts out of which the germinal layers arose.

Finally, I must invite special attention to the *metaphysical* character of the profound philosophic reflections which Wolff always added to his remarkable observations. He was a great monistic philosopher, in the best meaning of the word. It is unfortunate that his philosophic discoveries were ignored as completely as his observations for more than half a century. We must be careful to emphasise the fact of their clear monistic tendency.

CHAPTER III.

MODERN EMBRYOLOGY

We may distinguish three chief periods in the growth of our science of human embryology. The first has been considered in the preceding chapter; it embraces the whole of the preparatory period of research, and extends from Aristotle to Caspar Friedrich Wolff, or to the year 1759, in which the epoch-making *Theoria generationis* was published. The second period, with which we have now to deal, lasts about a century—that is to say, until the appearance of Darwin's *Origin of Species*, which brought about a change in the very foundations of biology, and, in particular, of embryology. The third period begins with Darwin. When we say that the second period lasted a full century, we must remember that Wolff's work had remained almost unnoticed during half the time—namely, until the year 1812. During the whole of these fifty-three years not a single book that appeared followed up the path that Wolff had opened, or extended his theory of embryonic development. We merely find his views—perfectly correct views, based on extensive observations of fact—mentioned here and there as erroneous; their opponents, who adhered to the dominant theory of preformation, did not even deign to reply to them. This unjust treatment was chiefly due to the extraordinary authority of Albrecht von Haller; it is one of the most unbecoming instances of a great authority, as such, preventing for a long time the recognition of established facts.

The general ignorance of Wolff's work was so great that at the beginning of the nineteenth century two scientists of Jena, Oken (1806) and Kleeer (1810), began independent research into the development of the alimentary canal of the chick, and hit upon the right clue to the embryonic puzzle, without knowing a word about Wolff's important treatise on the same subject. They were treading in his very footsteps without suspecting it. This can be easily proved from the fact that they did not travel as far as Wolff. It was not

until Meckel translated into German Wolff's book on the alimentary system, and pointed out its great importance, that the eyes of anatomists and physiologists were suddenly opened. At once a number of biologists instituted fresh embryological inquiries, and began to confirm Wolff's theory of epigenesis.

This resuscitation of embryology and development of the epigenesis-theory was chiefly connected with the university of Wurtzburg. One of the professors there at that time was Dollinger, an eminent biologist, and father of the famous Catholic historian who later distinguished himself by his opposition to the new dogma of papal infallibility. Dollinger was both a profound thinker and an accurate observer. He took the keenest interest in embryology, and worked at it a good deal. However, he is not himself responsible for any important result in this field. In 1816 a young medical doctor, whom we may at once designate as Wolff's chief successor, Karl Ernst von Baer, came to Wurtzburg. Baer's conversations with Dollinger on embryology led to a fresh series of most extensive investigations. Dollinger had expressed a wish that some young scientist should begin again under his guidance an independent inquiry into the development of the chick during the hatching of the egg. As neither he nor Baer had money enough to pay for an incubator and the proper control of the experiments, and for a competent artist to illustrate the various stages observed, the lead of the enterprise was given to Christian Pander, a wealthy friend of Baer's, who had been induced by Baer to come to Wurtzburg. An able engraver, Dalkow, was engaged to do the copper-plates. In a short time the embryology of the chick, in which Baer was taking the greatest indirect interest, was so far advanced that Pander was able to sketch the main features of it on the ground of Wolff's theory in the dissertation he published in 1817. He clearly enunciated the theory of germinal layers which Wolff

had anticipated, and established the truth of Wolff's idea of a development of the complicated systems of organs out of simple leaf-shaped primitive structures. According to Pander, the leaf-shaped object in the hen's egg divides, before the incubation has proceeded twelve hours, into two different layers, an external *ectoderm* layer and an internal *mesoderm* layer, between the two there develops later a third layer, the *vascular* (blood-vessel) layer.¹

Karl Ernst von Baer, who had set aside Pander's investigation, and had shown the liveliest interest in it after Pander's departure from Würzburg, began his own much more comprehensive research in 1810. He published the mature result nine years afterwards in his famous work, *Animal Embryology: Observations and Reflections* (not translated). This classic work still remains a model of careful observation united to profound philosophic speculation. The first part appeared in 1828, the second in 1837. The book proved to be the foundation on which the whole science of embryology has built down to our own day. It so far surpassed its predecessors, and Pander in particular, that it has become, after Wolff's work, the chief basis of modern embryology.

Baer was one of the greatest scientists of the nineteenth century, and exercised considerable influence on other branches of biology as well. He built up the theory of germinal layers, as a whole and in detail, so clearly and solidly that it has been the starting-point of embryological research ever since. He taught that in all the vertebrates first two and then four of those germinal layers are formed, and that the earliest rudimentary organs of the body arise by the conversion of these layers into tubes. He described the first appearance of the vertebrate embryo, as it may be seen in the globular yolk of the fertilized egg, as an oval disk which first divides into two layers. From the upper or *animal* layer are developed all the organs which accomplish the phenomena of animal life—the functions of sensation and motion, and the covering of the body. From the lower or *vegetative* layer come the organs which effect the vegetative life of the organism—nutrition, digestion, blood-formation, respiration, secretion, reproduction, etc.

Each of these original layers divides, according to Baer, into two thinner and superimposed layers or plates. He calls the two plates of the animal layer, the *skin-stratum* and *muscle-stratum*. From the upper of these plates, the *skin-stratum*, the external skin, or outer covering of the body, the central nervous system, and the sense-organs, are formed. From the lower, or *muscle-stratum*, the muscles, or fleshy parts and the bony skeleton—in a word, the motor organs—are evolved. In the same way, Baer said, the lower or vegetative layer splits into two plates, which he calls the *vascular-stratum* and the *musculo-stratum*. From the outer of the two (the *vascular*) the heart, blood-vessels, spleen, and the other vascular glands, the kidneys, and sexual glands, are formed. From the fourth or *innermost* layer, in fine, we get the internal and digestive lining of the alimentary canal and all its dependencies, the liver, lungs, salivary glands, etc. Baer had, in the main, correctly judged the significance of these four secondary embryonic layers, and he followed the conversion of them into the tube-shaped primitive organs with great perspicuity. He first solved the difficult problem of the transformation of this four-fold, flat, leaf-shaped, embryonic disk into the complete vertebrate body, through the conversion of the layers or plates into tubes. The flat leaves bend themselves in obedience to certain laws of growth, the borders of the curling plates approach nearer and nearer, until at last they come into actual contact. Thus out of the flat gut-plate is formed a hollow gut-tube, out of the flat spinal plate a hollow nerve-tube, from the skin-plate a skin-tube, and so on.

Among the many great services which Baer rendered to embryology, especially vertebrate embryology, we must not forget his discovery of the human ovum. Earlier scientists had, as a rule, of course, assumed that man developed out of an egg, like the other animals. In fact, the preformation theory held that the germs of the whole of humanity were stored already in Eve's ovum. But the real ovum escaped detection until the year 1827. This ovum is extremely small, being a tiny round vesicle about the size of a needle's eye as a tiny particle, but is otherwise quite invisible. This particle is formed in the ovary inside a much larger

¹ The technical terms which are bound to creep into this chapter will be fully understood later on.—TRANS.

globule, which takes the name of the Graafian follicle, from its discoverer, Graaf, and had previously been regarded as the true ovum. However, in 1827 Baer proved that it was not the real ovum, which is much smaller, and is contained within the follicle. (Compare the end of the twenty-ninth chapter.)

Baer was also the first to observe what is known as the *segmentation sphere* of the vertebrate; that is to say, the round vesicle which first develops out of the impregnated ovum, and the thin wall of which is made up of a single layer of regular, polygonal (many-cornered) cells (see the illustration in the twelfth chapter). Another discovery of his that was of great importance in constructing the vertebrate stem and the characteristic organization of this extensive group (to which man belongs) was the detection of the axial rod, or the *chorda dorsalis*. This is a long, round, cylindrical rod of cartilage which runs down the longer axis of the vertebrate embryo; it appears at an early stage, and is the first sketch of the spinal column, the solid skeletal axis of the vertebrate. In the lowest of the vertebrates, the amphioxus, the internal skeleton consists only of this rod throughout life. But even in the case of man and all the higher vertebrates it is round this rod that the spinal column and the brain are afterwards formed.

However, important as these and many other discoveries of Baer's were in vertebrate embryology, his researches were even more influential, from the circumstance that he was the first to employ the *comparative* method in studying the development of the animal frame. Baer occupied himself chiefly with the embryology of vertebrates (especially the birds and fishes). But he by no means confined his attention to these, gradually taking the various groups of the invertebrates into his sphere of study. As the general result of his comparative embryological research, Baer distinguished four different modes of development and four corresponding groups in the animal world. These chief groups or types are: 1, the vertebrates; 2, the articulates; 3, the molluscs; and 4, all the lower groups which were then wrongly comprehended under the general name of the radiata. Georges Cuvier had been the first to formulate this 'tinction', in 1812. He showed that these groups present specific differences in their whole internal structure, and the connection and

disposal of their systems of organs; and that, on the other hand, all the animals of the same type—say, the vertebrates—essentially agreed in their inner structure, in spite of the greatest superficial differences. But Baer proved that these four groups are also quite differently developed from the ovum; and that the series of embryonic forms is the same throughout for animals of the same type, but different in the case of other animals. Up to that time the chief aim in the classification of the animal kingdom was to arrange all the animals from lowest to highest, from the infusorium to man, in one long and continuous series. The erroneous idea prevailed nearly everywhere that there was one uninterrupted chain of evolution from the lowest animal to the highest. Cuvier and Baer proved that this view was false, and that we must distinguish four totally different types of animals, on the ground of anatomic structure and embryonic development.

Baer's epoch-making works aroused an extraordinary and widespread interest in embryological research. Immediately afterwards we find a great number of observers at work in the newly opened field, enlarging it in a very short time with great energy by their various discoveries in detail. Next to Baer's comes the admirable work of Heinrich Rathke, of Königsberg (died 1860); he made an extensive study of the embryology, not only of the invertebrates (crustaceans, insects, molluscs), but also, and particularly, of the vertebrates (fishes, tortoises, serpents, crocodiles, etc.). We owe the first comprehensive studies of mammal embryology to the careful research of Wilhelm Bischoff, of Munich; his embryology of the rabbit (1830), the dog (1832), the guinea-pig (1832), and the doe (1834), still form classical studies. About the same time a great impetus was given to the embryology of the invertebrates. The way was opened through this obscure province by the studies of the famous Berlin zoologist, Johannes Müller, on the echinodermata. He was followed by Albert Kölliker, of Würzburg, writing on the cuttle-fish (or the cephalopods), Siebold and Huxley on worms and zoophytes, Fritz Müller (Dietrich) on the crustaceans, Weismann on insects, and so on. The number of workers in this field has greatly increased of late, and a quantity of new and astonishing discoveries have been made. One notices, in several of these recent works on

embryology, that their authors are too little acquainted with comparative anatomy and classification. Paleontology is, unfortunately, altogether neglected by many of these new writers, although this interesting science furnishes most important facts for phylogeny, and thus often proves of very great service in ontogeny.

A very important advance was made in our science in 1839, when the cellular theory was established, and a new field of inquiry bearing on embryology was suddenly opened. When the famous botanist, M. Schleiden, of Jena, showed in 1838, with the aid of the microscope, that every plant was made up of innumerable elementary parts, which we call *cells*, a pupil of Johannes Muller at Berlin, Theodor Schwann, applied the discovery at once to the animal organism. He showed that in the animal body as well, when we examine its tissues in the microscope, we find these cells everywhere to be the elementary units. All the different tissues of the organism, especially the very dissimilar tissues of the nerves, muscles, bones, external skin, mucous lining, etc., are originally formed out of cells; and this is also true of all the tissues of the plant. These cells are separate living beings; they are the citizens of the State which the entire multicellular organism seems to be. This important discovery was bound to be of service to embryology, as it raised a number of new questions: What is the relation of the cells to the germinal layers? Are the germinal layers composed of cells, and what is their relation to the cells of the tissues that form later? How does the ovum stand in the cellular theory? Is the ovum itself a cell, or is it composed of cells? These important questions were now imposed on the embryologist by the cellular theory.

The most notable effort to answer these questions—which were attacked on all sides by different students—is contained in the famous work, *Inquiries into the Development of the Vertebrates* (not translated) of Robert Remak, of Berlin (1851). This gifted scientist succeeded in mastering, by a complete reform of the science, the great difficulties which the cellular theory had at first put in the way of embryology. A Berlin anatomist, Carl Boguslaus Reichert, had already attempted to explain the origin of the tissues. But this attempt was bound to miscarry, since its not very clear-headed author lacked a sound acquaintance with embryology and

the cell theory, and even with the structure and development of the tissue in particular. Remak at length brought order into the dreadful confusion that Reichert had caused; he gave a perfectly simple explanation of the origin of the tissues. In his opinion the animal ovum is always a *simple cell*: the germinal layers which develop out of it are always composed of cells; and these cells that constitute the germinal layers arise simply from the continuous and repeated cleaving (segmentation) of the original solitary cell. It first divides into two and then into four cells; out of these four cells are born eight, then sixteen, thirty-two, and so on. Thus, in the embryonic development of every animal and plant there is formed first of all out of the simple egg cell, by a repeated sub-division, a cluster of cells, as Kolliker had already stated in connection with the cephalopods in 1844. The cells of this group spread themselves out flat and form leaves or plates, each of these leaves is formed exclusively out of cells. The cells of different layers assume different shapes, increase, and differentiate; and in the end there is a further cleavage (differentiation) and division of work of the cells within the layers, and from these all the different tissues of the body proceed.

These are the simple foundations of *Autogeny*, or the science that treats of the development of the tissues (*histogenesis*), as it was established by Remak and Kolliker. Remak, in determining more closely the part which the different germinal layers play in the formation of the various tissues and organs, and in applying the theory of evolution to the cells and the tissues they compose, raised the theory of germinal layers, at least as far as it regards the vertebrates, to a high degree of perfection.

Remak showed that three layers are formed out of the two germinal layers which compose the first simple leaf-shaped structure of the vertebrate body (or the "germinal disk"), as the lower layer splits into two plates. These three layers have a very definite relation to the various tissues. First of all, the cells which form the outer skin of the body (the epidermis), with its various dependencies (hairs, nails, etc.)—that is to say, the entire outer envelope of the body—are developed out of the outer or upper layer; but these are also developed in a curious way out of the same layer the cells which form the central nervous system, the

brain and the spinal cord. In the second place, the inner or lower germinal layer gives rise only to the cells which form the epithelium (the whole inner lining) of the alimentary canal and all that depends on it (the lungs, liver, pancreas, etc.), and that

of the body.

The middle layer gives rise to all the rest of the body, the muscles, blood, bones, cartilage, etc. Remak further proved that this middle layer, which he calls "the motor-germinative layer," proceeds to subdivide into two secondary layers. Thus we find once more the four layers which Baer had indicated. Remak calls the outer secondary leaf of the middle layer (Baer's "concular layer") "the skin-fibre layer" (it would be better to say, skin-fibre layer), the inner secondary leaf (Baer's "vascular layer") he gave the name of the "alimentary-fibre layer," the outer envelope of the alimentary heart, the blood-vessels.

Remak for *Autogeny*, or the self-formation of the tissues, our has been gradually built up a in detail. There have been attempts to restrict and even destroy Remak's principles. The two anatomists, Reichert (of Berlin) and Wilhelm His (of Lauscha), especially, have endeavored in their works to introduce a new opinion of the embryonic development of the

primary germinal layers would not be the sole sources of formation. But these efforts were so seriously marred by ignorance of comparative anatomy, an imperfect acquaintance with ontogenesis, and a complete neglect of phylogenesis, that they could not have more than a passing success. We can only explain how these curious attacks of Reichert and His came to be regarded for a time as advances by the general lack of discrimination and of grasp of the true object of embryology.

Wilhelm His published, in 1868, his massive *Researches into the Earliest Form of the Vertebrate Body*; one of the curiosities of embryological literature. The author imagines that he has built

a "mechanical theory of embryonic development" by merely giving an exact description of the embryology of the chick, without any regard to comparative anatomy and phylogeny, and thus falls

parallel in the history of biological literature. As the final result of his laborious investigations, His tells us "that a comparatively simple law of growth is the initial thing in the first development. Every formation, whether it consist in cleavage of layers, or folding, or complete division, is a consequence of this fundamental law." Unfortunately, he

growth" is, just as other opponents of the theory of selection, who would put in its place a great "law of evolution."

Anything about the nature of this, nevertheless, it is quite clear from His's works that he imagines constructive nature to be a sort of skillful tailor. The ingenious operator succeeds in bringing

all the living things by cutting, tearing, and folding, tugging and splitting, and so on.

Embryological theories excited a good deal of interest at the time of publication, and have evoked a fair amount of literature in the last few decades. He professed to explain the most complicated parts of organic construction (such as the development of the brain) in the simplest way on mechanical principles, and to derive them immediately from simple physical processes (such as unequal distension of elastic plates).

It is quite true that a mechanical or monistic explanation (or a reduction of natural phenomena to physical and chemical processes) is the ideal of modern science, and this ideal would be realized if we could succeed in expressing these formative processes in mathematical formulae. His has, therefore, inserted plenty of numbers and measurements in his embryological works, and gives of "exact" quantity of

fortunately, they are of no value, and do not help us in the least in forming an "exact" acquaintance with the embryonic phenomena. Indeed, they wander from the true path altogether by neglecting the phylogenetic method; this, he thinks, is "a more by-path," and is "not necessary at all for the explanation of the facts of

embryology," which are the direct consequences of physiological principles. What this takes to be a simple physical process—for instance, the folding of the germinal layers (in the formation of the medullary tube, alimentary tube, etc.)—is, as a matter of fact, the direct result of the growth of the various cells which form these organic structures; but these growth-motions have themselves been transmitted by heredity from parents and ancestors, and are only the hereditary repetition of countless phylogenetic changes which have taken place for thousands of years in the race-history of the said ancestors. Each of these historical changes was, of course, originally due to adaptation; it was, in other words, physiological, and reducible to mechanical causes. But we have, naturally, no means of observing them now. It is only by the hypotheses of the science of evolution that we can form an approximate idea of the organic links in this historic chain.

All the best recent research in animal embryology has led to the confirmation and development of Baer and Remak's theory of the germinal layers. One of the most important advances in this direction of late was the discovery that the two primary layers out of which is built the body of all vertebrates (including man) are also present in all the invertebrates, with the sole exception of the lowest group, the unicellular protozoa. Huxley had detected them in the medusa in 1846. He showed that the two layers of cells from which the body of this zoophyte is developed correspond, both morphologically and physiologically, to the two original germinal layers of the vertebrate. The outer layer, from which come the external skin and the muscles, was then called by Allman (1853) the "ectoderm" (=outer layer, or skin); the inner layer, which forms the alimentary and reproductive organs, was called the "entoderm" (=inner layer). In 1867 and the following years the discovery of the germinal layers was extended to other groups of the invertebrates. In particular, the indefatigable Russian zoologist, Kowalevsky, found them in all the most diverse sections of the invertebrates—the worms, tunicates, echinoderms, molluscs, articulates, etc.

In my monograph on the sponges (1872) I proved that these two primary germinal layers are also found in that group, and

that they may be traced from it right up to man, through all the various classes, in identical form. This "homology of the two primary germinal layers" extends through the whole of the metazoa, or tissue-forming animals; that is to say, through the whole animal kingdom, with the one exception of its lowest section, the unicellular beings, or protozoa. These lowly organized animals do not form germinal layers, and therefore do not succeed in forming true tissue. Their whole body consists of a single cell (as is the case with the amoeba and infusoria), or of a loose aggregation of only slightly differentiated cells, though it may not even reach the full structure of a single cell (as with the murex). But in all other animals the organism first grows into two primary layers, the outer or *animal layer* (the ectoderm, epiblast, or ectoblast), and the inner or *vegetal layer* (the entoderm, hypoblast, or endoblast), and from these the tissues and organs are formed. The first and oldest organ of all these metazoa is the primitive gut (or pro-gaster) and its opening, the primitive mouth (prostoma). The typical embryonic form of the metazoa, as it is presented for a time by this simple structure of the two-layered body, is called the *gastrula*; it is to be conceived as the hereditary reproduction of some primitive common ancestor of the metazoa, which we call the *gastraea*. This applies to the sponges and other zoophytes, and to the worms, the molluscs, echinoderms, articulates, and vertebrates. All these animals may be comprised under the general heading of "gut animals," or metazoa, in contradistinction to the gutless protozoa.

I have pointed out in my *Study of the Gastraea Theory* [not translated] (1873) the important consequences of this conception in the morphology and classification of the animal world. I also divided the realm of metazoa into two great groups, the lower and higher metazoa. In the first are comprised the *ciliatederata* (also called zoophytes, or "plant-animals"). In the lower forms of this group the body consists throughout life merely of the primary germinal layers, with the cells sometimes more and sometimes less differentiated. But with the higher forms of the ciliatederata (the corals, higher medusae, ctenophore, and platodes) a middle layer, or mesoderm, often of considerable size, is developed between the

other two layers; but blood and an internal cavity are still lacking.

To the second great group of the metazoa I gave the name of the *coelomaria*, or *bilateralis* (or the bilateral higher forms). They all have a cavity within the body (coeloma), and most of them have blood and blood-vessels. In this are comprised the six higher stems of the animal kingdom, the annulata and their descendants, the mollusca, echinodermata, articulatæ, tunicata, and vertebrata. In all these bilateral organisms the two-sided body is formed out of four secondary germinal layers, of which the inner two construct the wall of the alimentary canal, and the outer two the wall of the body. Between the two pairs of layers lies the cavity (coeloma).

Although I laid special stress on the great morphological importance of this cavity in my *Study of the Gastræa Theory*, and endeavoured to prove the significance of the four secondary germinal layers in the organisation of the coelomaria, I was unable to deal satisfactorily with the difficult question of the mode of their origin. This was done eight years afterwards by the brothers Oscar and Richard Hertwig in their careful and extensive comparative studies. In their masterly *Coelom Theory. An Attempt to Explain the Middle Germinal Layer* [not translated] (1881) they showed that in most of the metazoa, especially in all the vertebrates, the body-cavity arises in the same way, by the outgrowth of two sacs from the inner layer. These two coelom-pouches proceed from the rudimentary mouth of the gastrula, between the two primary layers. The inner plate of the two-layered coelom-pouch (the visceral layer) joins itself to the endoderm; the outer plate (parietal layer) unites with the ectoderm. Thus are formed the double-layered gut-wall within and the double-layered body-wall without; and between the two is formed the cavity of the coelom, by the blending of the right and left coelom-sacs. We shall see this more fully in Chap. X.

The many new points of view and fresh ideas suggested by my gastræa theory and Hertwig's coelom theory led to the publication of a number of writings on the theory of germinal layers. Most of them set out to oppose it at first, but in the end the majority supported it. Of late years both theories are accepted in their essential features by nearly every competent man of science, and light and order

have been introduced into this once dark and contradictory field of research. A further cause of congratulation for this solution of the great embryological controversy is that it brought with it a recognition of the need for phylogenetic study and explanation.

Interest and practice in embryological research have been remarkably stimulated during the past thirty years by this appreciation of phylogenetic methods. Hundreds of assiduous and able observers are now engaged in the development of comparative embryology and its establishment on a basis of evolution, whereas they numbered only a few dozen not many decades ago. It would take too long to enumerate even the most important of the countless valuable works which have enriched embryological literature since that time. References to them will be found in the latest manuals of embryology of Kolliker, Balfour, Hertwig, Kollman, Korschelt, and Heider.

Kolliker's *Entwickelungsgeschichte des Menschen und der höheren Thiere*, the first edition of which appeared forty-two years ago, had the rare merit at that time of gathering into presentable form the scattered attainments of the science, and expounding them in some sort of unity on the basis of the cellular theory and the theory of germinal layers. Unfortunately, the distinguished Würzburg anatomist, to whom comparative anatomy, histology, and ontogeny owe so much, is opposed to the theory of descent generally and to Darwinism in particular. All the other manuals I have mentioned take a decided stand on evolution. Francis Balfour has carefully collected and presented with discrimination, in his *Manual of Comparative Embryology* (1880), the very scattered and extensive literature of the subject; he has also widened the basis of the gastræa theory by a comparative description of the rise of the organs from the germinal layers in all the chief groups of the animal kingdom, and has given a most thorough empirical support to the principles I have formulated. A comparison of his work with the excellent *Text-Book of the Embryology of the Vertebrates* (1890) [translation 1895] of Korschelt and Heider shows what astonishing progress has been made in the science in the course of ten years. I would especially recommend the manuals of Julius Kollmann and Oscar Hertwig to those readers who are stimulated to further study by these chapters on human

embryology. Kollmar's book is commendable for its clear statement of the subject and very fine original illustrations; its author adheres firmly to the biogenetic law, and uses it throughout with considerable profit. That is not the case in Oscar Hertwig's recent *Text-book of the Embryology of Man and the Mammals* (translative 1892 and 1899) (seventh edition, 1903). This able anatomist has of late often been quoted as an opponent of the biogenetic law, although he himself had demon-

strated that the oscillation is partly due to the middle of the series. We must be careful with regard to hypotheses; though it is quite impossible to make any headway in the explanation of facts without them. However, the purely descriptive part of embryology in Hertwig's *Text-book* is very thorough and reliable.

A new branch of embryological research has been studied very assiduously in the last decade of the nineteenth century—namely, "experimental embryology." The great importance which has been attached to the application of physical experiments to the living organism for the last hundred years, and the valuable results that it has given to physiology in the study of the vital phenomena, have led to its extension to embryology. I was the first to make experiments of this kind during a stay of four months on the Canary Island, Lanzarote, in 1866. I there made a thorough investigation of the almost unknown embryology of the siphonophore. I cut a number of the embryos of these animals (which develop freely in the water, and pass through a very curious transformation), at an early stage, into several pieces, and found that a fresh organism (more or less complete,

according to the size of the piece) was developed from each particle. More recently some of my pupils have made similar experiments with the embryos of vertebrates (especially the frog) and some of the invertebrates. Wilhelm Roux, in particular, has made extensive experiments, and based on them a special "mechanical embryology," which has given rise to a good deal of discussion and controversy. Roux has published a special journal for these subjects since the *Archiv für Entwicklungs-*

the contribution of many of them are valuable papers on the physiology and pathology of the embryo. Pathological experiments—the placing of the embryo in abnormal conditions—have yielded many interesting results; just as the physiology of the normal body has for a long time derived assistance from the pathology of the diseased organism. Older of these mechanical-embryological articles return to the erroneous methods of Hux, and are only misleading. There must be said of the many contributions of mechanical embryology which take up a position of hostility to the theory of descent and its chief embryological foundation—the biogenetic law. This law, however, when rightly understood, is not opposed to, but is the best and most solid support of, a sound mechanical embryology. Impartial reflection and a due attention to paleontology and comparative anatomy should convince these so-called mechanicians that the facts they have discovered—and, indeed, the whole embryological process—cannot be fully understood without the theory of descent and the biogenetic law.

CHAPTER IV.

THE OLDER PHYLOGENY

THE embryology of man and the animals, the history of which we have reviewed in the last two chapters, was mainly a descriptive science forty years ago. The earlier investigations in this province were

chiefly directed to the discovery, by careful observation, of the wonderful facts of the embryonic development of the animal body from the ovum. Forty years ago no one dared attack the question of the

causes of these phenomena. For fully a century, from the year 1755, when Wolff's solid *Theoria generationis* appeared, until 1859, when Darwin published his famous *Origin of Species*, the real causes of the embryonic processes were quite unknown. No one thought of seeking the agencies that effected this marvellous succession of structures. The task was thought to be so difficult as almost to pass beyond the limits of human thought. It was reserved for Charles Darwin to initiate us into the knowledge of these causes. This compels us to recognise in this great genius, who wrought a complete revolution in the whole field of biology, a founder at the same time of a new period in embryology. It is true that Darwin occupied himself very little with direct embryological research, and even in his chief work he only touches incidentally on the embryonic phenomena, but by his reform of the theory of descent and the founding of the theory of selection he has given us the means of attaining to a real knowledge of the causes of embryonic formation. That is, in my opinion, the chief feature in Darwin's incalculable influence on the whole science of evolution.

When we turn our attention to this latest period of embryological research, we pass into the second division of organic evolution—stem-evolution, or phylogeny. I have already indicated in the first chapter the important and intimate causal connection between these two sections of the science of evolution—between the evolution of the individual and that of his ancestors. We have formulated this connection in the biogenetic law, the shorter evolution, that of the individual, or *ontogenesis*, is a rapid and summary repetition, a condensed recapitulation, of the larger evolution, or that of the species. In this principle we express all the essential points relating to the causes of evolution; and we shall seek throughout this work to confirm this principle and lend it the support of facts. When we look to its causal significance, perhaps it would be better to formulate the biogenetic law thus: "The evolution of the species and the stem (*phylon*) shows us, in the physiological functions of heredity and adaptation, the conditioning causes on which the evolution of the individual depends"; or, more briefly: "Phylogenesis is the mechanical cause of ontogenesis."

But before we reach the most

achievement by which Darwin revealed the causes of evolution to us, we must glance at the efforts of earlier scientists on this object. Our historical inquiry into these will be even shorter than that into the work done in the field of ontogeny. We have very few names to consider here. At the head of them we find the great French naturalist, Jean Lamarck, who first established evolution as a scientific theory in 1809. Even before his time, however, the chief philosopher, Kant, and the chief poet, Goethe, of Germany had occupied themselves with the subject. But their efforts passed almost without recognition in the eighteenth century. A "philosophy of nature" did not arise until the beginning of the nineteenth century. In the whole of the time before this no one had ventured to raise seriously the question of the origin of species, which is the culminating point of phylogeny. On all sides it was regarded as an insoluble enigma.

The whole science of the evolution of man and the other animals is intimately connected with the question of the nature of species, or with the problem of the origin of the various animals which we group together under the name of species. Thus the definition of the species becomes important. It is well known that this definition was given by Linné, who, in his famous *Systema Naturæ* (1735), the first to classify and name the various groups of animals and plants, and drew up an orderly scheme of the species then known. Since that time "species" has been the most important and indispensable idea in descriptive natural history, in zoological and botanical classification, although there have been endless controversies as to its real meaning.

What, then, is this "organic species"? Linné himself appealed directly to the Mosaic narrative; he believed that, as it is stated in *Genesis*, one pair of each species of animals and plants was created in the beginning, and that all the individuals of each species are the descendants of these created couples. As for the hermaphrodites (organisms that have male and female organs in one being), he thought it sufficed to assume the creation of one sole individual, since this would be fully competent to propagate its species. Further developing these mystic ideas, Linné went on to borrow from *Genesis* the account of the deluge and of Noah's ark as a record for a catalogue of the mammalian

and topographical distribution of organisms. He accepted the story that all the plants, animals, and men on the earth were swept away in a universal deluge, except the couples preserved with Noah in the ark, and ultimately landed on Mount Ararat. This mountain seemed to Linné particularly suitable for the landing, as it reaches a height of more than 16,000 feet, and thus provides in its higher zones the several climates demanded by the various species of animals and plants. The animals that were accustomed to a cold climate could remain at the summit; those used to a warm climate could descend to the foot; and those requiring a temperate climate could remain half-way down. From this point the re-population of the earth with animals and plants could proceed.

It was impossible to have any scientific notion of the method of evolution in Linné's time, as one of the chief sources of information, paleontology, was still wholly unknown. This science of the fossil remains of extinct animals and plants is very closely bound up with the whole question of evolution. It is impossible to explain the origin of living organisms without appealing to it. But this science did not rise until a much later date. The real founder of scientific paleontology was Georges Cuvier, the most distinguished zoologist who, after Linné, worked at the classification of the animal world, and effected a complete revolution in systematic zoology at the beginning of the nineteenth century. In regard to the nature of the species he associated himself with Linné and the Mosaic story of creation, though this was more difficult for him with his acquaintance with fossil remains. He clearly showed that a number of quite different animal populations have lived on the earth; and he claimed that we must distinguish a number of stages in the history of our planet, each of which was characterized by a special population of animals and plants. These successive populations were, he said, quite independent of each other, and therefore the supernatural creative act, which was demanded as the origin of the animals and plants by the dominant creed, must have been repeated several times. In this way a whole series of different creative periods must have succeeded each other; and in connection with these he had to assume that stupendous revolu-

tions or cataclysms—something like the legendary deluge—must have taken place repeatedly. Cuvier was all the more interested in these catastrophes or cataclysms as geology was just beginning to assert itself, and great progress was being made in our knowledge of the structure and formation of the earth's crust. The various strata of the crust were being carefully examined, especially by the famous geologist Werner and his school, and the fossils found in them were being classified, and these researches also seemed to point to a variety of creative periods. In each period the earth's crust, composed of the various strata, seemed to be differently constituted, just like the population of animals and plants that then lived on it. Cuvier combined this notion with the results of his own paleontological and zoological research, and in his effort to get a consistent view of the whole process of the earth's history he came to form the theory which is known as "the catastrophic theory," or the theory of terrestrial revolutions. According to this theory, there have been a series of mighty cataclysms on the earth, and these have suddenly destroyed the whole animal and plant population then living on it, after each cataclysm there was a fresh creation of living things throughout the earth. As this creation could not be explained by natural laws, it was necessary to appeal to an intervention on the part of the Creator. This catastrophic theory, which Cuvier described in a special work, was soon generally accepted, and retained its position in biology for half a century.

However, Cuvier's theory was completely overthrown sixty years ago by the geologists, led by Charles Lyell, the most distinguished worker in this field of science. Lyell proved in his famous *Principles of Geology* (1830) that the theory was false, in so far as it concerned the crust of the earth; that it was totally unnecessary to bring in supernatural agencies or general catastrophes in order to explain the structure and formation of the mountains; and that we can explain them by the familiar agencies which are at work to-day in altering and reconstructing the surface of the earth. These causes are—the action of the atmosphere and water in its various forms (snow, ice, fog, rain, the wear of the river, and the stormy ocean), and the volcanic action which is exerted by the molten central

mass. Lyell convincingly proved that these natural causes are quite adequate to explain every feature in the build and formation of the crust. Hence Cuvier's theory of cataclysms was very soon driven out of the province of geology, though it remained for another thirty years in undisputed authority in biology. All the zoologists and botanists who gave any thought to the question of the origin of organisms adhered to Cuvier's erroneous idea of revolutions and new creations.

In order to illustrate the complete stagnancy of biology from 1830 to 1859 on the question of the origin of the various species of animals and plants, I may say, from my own experience, that during the whole of my university studies I never heard a single word said about this most important problem of the science. I was fortunate enough at that time (1852-1857) to have the most distinguished masters for every branch of biological science. Not one of them ever mentioned this question of the origin of species. Not a word was ever said about the earlier efforts to understand the formation of living things, nor about Lamarck's *Philosophie Zoologique* which had made a fresh attack on the problem in 1809. Hence it is easy to understand the enormous opposition that Darwin encountered when he took up the question for the first time. His views seemed to float in the air, without a single previous effort to support them. The whole question of the formation of living things was considered by biologists, until 1859, as pertaining to the province of religion and transcendentalism, even in speculative philosophy, in which the question had been approached from various sides, no one had ventured to give it serious treatment. This was due to the dualistic system of Immanuel Kant, who taught a natural system of evolution as far as the inorganic world was concerned, but, on the whole, adopted a supernaturalist system as regards the origin of living things. He even went so far as to say: "It is quite certain that we cannot even satisfactorily understand, much less explain, the nature of an organism and its internal forces on purely mechanical principles; it is so certain, indeed, that we may confidently say: 'It is absurd for a man to imagine even that some day a Newton will arise who will explain the origin of a single blade of grass by natural laws not controlled by design'—such a hope is entirely forbidden us." In these

words Kant definitely adopts the dualistic and teleological point of view for biological science.

Nevertheless, Kant deserted this point of view at times, particularly in several remarkable passages which I have dealt with at length in my *Natural History of Creation* (chap. v.), where he expresses himself in the opposite, or monistic, sense. In fact, these passages would justify one, as I showed, in claiming his support for the theory of evolution. However, these monistic passages are only stray gleams of light; as a rule, Kant adheres in biology to the obscure dualistic ideas, according to which the forces at work in inorganic nature are quite different from those of the organic world. This dualistic system prevails in academic philosophy to-day—most of our philosophers still regarding these two provinces as totally distinct. They put, on the one side, the inorganic or "lifeless" world, in which there are at work only mechanical laws, acting necessarily and without design; and, on the other, the province of organic nature, in which none of the phenomena can be properly understood, either as regards their inner nature or their origin, except in the light of preconceived design, carried out by final or purposive causes.

The prevalence of this unfortunate dualistic prejudice prevented the problem of the origin of species, and the connected question of the origin of man, from being regarded by the bulk of people as a scientific question at all until 1859. Nevertheless, a few distinguished students, free from the current prejudice, began, at the commencement of the nineteenth century, to make a serious attack on the problem. The merit of this attaches particularly to what is known as "the older school of natural philosophy," which has been so much misrepresented, and which included Jean Lamarck, Buffon, Geoffroy St. Hilaire, and Blainville in France, Wolfgang Goethe, Reinhold Treviranus, Schelling, and Lorenz Oken in Germany (and Erasmus Darwin in England).

The gifted natural philosopher who treated this difficult question with the greatest sagacity and comprehensiveness was Jean Lamarck. He was born at Bazentin, in Picardy, on August 1st, 1744; he was the son of a clergyman, and was destined for the Church. But he turned to seek glory in the army, and eventually devoted himself to science.

His *Philosophie Zoologique* was the

first scientific attempt to sketch the real course of the origin of species, the first "natural history of creation" of plants, animals, and men. But, as in the case of Wolff's book, this remarkably able work had no influence whatever; neither one nor the other could obtain any recognition from their prejudiced contemporaries. No man of science was stimulated to take an interest in the work, and to develop the germs it contained of the most important biological truths. The most distinguished botanists and zoologists entirely rejected it, and did not even deign to reply to it. Cuvier, who lived and worked in the same city, has not thought fit to devote a single syllable to this great achievement in his memoir on progress in the sciences, in which the pettiest observations found a place. In short, Lamarck's *Philosophie Zoologique* shared the fate of Wolff's theory of development, and was for half a century ignored and neglected. The German scientists, especially Oken and Goethe, who were occupied with similar speculations at the same time, seem to have known nothing about Lamarck's work. If they had known it, they would have been greatly helped by it, and might have carried the theory of evolution much farther than they found it possible to do.

To give an idea of the great importance of the *Philosophie Zoologique*, I will briefly explain Lamarck's leading thought. He held that there was no essential difference between living and lifeless beings. Nature is one united and connected system of phenomena; and the forces which fashion the lifeless bodies are the only ones at work in the kingdom of living things. We have, therefore, to use the same method of investigation and explanation in both provinces. Life is only a physical phenomenon. All the plants and animals, with man at their head, are to be explained, in structure and life, by mechanical or efficient causes, without any appeal to final causes, just as in the case of minerals and other inorganic bodies. This applies equally to the origin of the various species. We must not assume any original creation, or repeated creations (as in Cuvier's theory), to explain this, but a natural, continuous, and necessary evolution. The whole evolutionary process has been uninterrupted. All the different kinds of animals and plants which we see to-day, or that have ever lived, have descended in a natural way from earlier and different

species; all come from one common stock, or from a few common ancestors. These remote ancestors must have been quite simple organisms of the lowest type, arising by spontaneous generation from inorganic matter. The succeeding species have been constantly modified by adaptation to their varying environment (especially by use and habit), and have transmitted their modifications to their successors by heredity.

Lamarck was the first to formulate as a scientific theory the natural origin of living things, including man, and to push the theory to its extreme conclusions—the rise of the earliest organisms by spontaneous generation (or abiogenesis) and the descent of man from the nearest related mammal, the ape. He sought to explain this last point, which is of especial interest to us here, by the same agencies which he found at work in the natural origin of the plant and animal species. He considered use and habit (adaptation) on the one hand, and heredity on the other, to be the chief of these agencies. The most important modifications of the organs of plants and animals are due, in his opinion, to the function of these very organs, or to the use or disuse of them. To give a few examples, the woodpecker and the humming-bird have got their peculiarly long tongues from the habit of extracting their food with their tongues from deep and narrow folds or canals; the frog has developed the web between his toes by his own swimming; the giraffe has lengthened his neck by stretching up to the higher branches of trees, and so on. It is quite certain that this use or disuse of organs is a most important factor in organic development, but it is not sufficient to explain the origin of species.

To adaptation we must add heredity as the second and not less important agency, as Lamarck perfectly recognised. He said that the modification of the organs in any one individual by use or disuse was slight, but that it was increased by accumulation in passing by heredity from generation to generation. But he missed altogether the principle which Darwin afterwards found to be the chief factor in the theory of transformation—namely, the principle of natural selection in the struggle for existence. It was partly owing to his failure to detect this supremely important element, and partly to the poor condition of all biological science at the time, that Lamarck did not

succeed in establishing more firmly his theory of the common descent of man and the other animals.

Independently of Lamarck, the older German school of natural philosophy, especially Reinhold Treviranus, in his *Biologie* (1802), and Lorenz Oken, in his *Naturphilosophie* (1804), turned its attention to the problem of evolution about the end of the eighteenth and beginning of the nineteenth century. I have described its work in my *History of Creation* (chap. iv.) Here I can only deal with the brilliant genius whose evolutionary ideas are of special interest—the greatest of German poets, Wolfgang Goethe. With his keen eye for the beauties of nature, and his profound insight into its life, Goethe was early attracted to the study of various natural sciences. It was the Livornite occupation of his leisure hours throughout life. He gave particular and protracted attention to the theory of colours. But the most valuable of his scientific studies are those which relate to that "living, glorious, precious thing," the organism. He made profound research into the science of structures or morphology (morphos = form). Here, with the aid of comparative anatomy, he obtained the most brilliant results, and went far in advance of his time. I may mention, in particular, his verified theory of the skull, his discovery of the pineal gland in man, his system of the metamorphosis of plants, etc. These morphological studies led Goethe on to research into the formation and modification of organic structures, which we must count as the first germ of the science of evolution. He approaches so near to the theory of descent that we must regard him, after Lamarck, as one of its earliest founders. It is true that he never formulated a complete scientific theory of evolution, but we find a number of remarkable suggestions of it in his splendid miscellaneous essays on morphology. Some of them are really among the very basic ideas of the science of evolution. He says, for instance (1807): "When we compare plants and animals in their most rudimentary forms, it is almost impossible to distinguish between

them. But we may say that the plants and animals, beginning with an almost inseparable closeness, gradually advance along two divergent lines, until the plant at last grows in the solid, enduring tree and the animal attains in man to the highest degree of mobility and freedom." That Goethe was not merely speaking in a poetical, but in a literal genealogical, sense of this close affinity of organic forms is clear from other remarkable passages in which he treats of their variety in outward form and unity in internal structure. He believes that every living thing has arisen by the interaction of two opposing formative forces or impulses. The internal or "centripetal" force, the type or "impulse to specification," seeks to maintain the constancy of the specific forms in the succession of generations: this is *Archedy*. The external or "centrifugal" force, the element of variation or "impulse to metamorphosis," is continually modifying the species by changing their environment; this is *adaptation*. In these significant conceptions Goethe approaches very close to a recognition of the two great mechanical factors which we now assign as the chief causes of the formation of species.

Now, over, in order to appreciate Goethe's views on morphology, one must associate his decidedly pantheistic conception of nature with his pantheistic philosophy. The warm and keen interest with which he followed, in his last years, the controversies of contemporary French scientists, and especially the struggle between Cuvier and Geoffroy St. Hilaire (see chap. iv of *The History of Creation*), is very characteristic. It is also necessary to be familiar with his style and general tenour of thought in order to appreciate rightly the many allusions to evolution found in his writings. Otherwise, one is apt to make serious errors.

He approached so close, at the end of the eighteenth century, to the principles of the science of evolution that he may well be described as the first forerunner of Darwin, although he did not go so far as to formulate evolution as a scientific system, as Lamarck did.

CHAPTER V.

THE MODERN SCIENCE OF EVOLUTION

We owe so much of the progress of scientific knowledge to Darwin's *Origin of Species* that its influence is almost without parallel in the history of science. The literature of Darwinism grows from day to day, not only on the side of academic zoology and botany, the sciences which were chiefly affected by Darwin's theory, but in a far wider circle, so that we find Darwinism discussed in popular literature with a vigour and zest that are girant to no other scientific conception. This remarkable success is due chiefly to two circumstances. In the first place, all the sciences, and especially biology, have made astounding progress in the last half-century, and have furnished a very vast quantity of proofs of the theory of evolution. In striking contrast to the failure of Lamarck and the older scientists to attract attention to their effort to explain the origin of living things and of man, we have this second and successful effort of Darwin, which was able to gather to its support a large number of established facts. Availing himself of the progress already made, he had very different scientific proofs to allege than Lamarck, or St. Hilaire, or Goethe, or Treviranus had had. But, in the second place, we must acknowledge that Darwin had the special distinction of approaching the subject from an entirely new side, and of basing the theory of descent on a consistent system, which now goes by the name of Darwinism.

Lamarck had unsuccessfully attempted to explain the modification of organisms that descend from a common form chiefly by the action of habit and the use of organs, though with the aid of heredity. But Darwin's success was complete when he independently sought to give a mechanical explanation, on a quite new ground, of this modification of plant and animal structures by adaptation and heredity. He was impelled to his theory of selection on the following grounds. He compared the origin of the various kinds of animals and plants which we modify artificially — by the action of

artificial selection in horticulture and among domestic animals — with the origin of the species of animals and plants in their natural state. He then found that the agencies which we employ in the modification of forms by artificial selection are also at work in Nature. The chief of these agencies he held to be "the struggle for life." The gist of this peculiarly Darwinian idea is given in this formula: "The struggle for existence produces new species without premeditated design in the life of Nature, in the same way that the will of man consciously selects new races in artificial conditions. The gardener or the farmer selects new forms as he wills for his own profit, by ingeniously using the agency of heredity and adaptation for the modification of structures; so, in the natural state, the struggle for life is always unconsciously modifying the various species of living things. This struggle for life, or competition of organisms in securing the means of subsistence, acts without any conscious design, but it is none the less effective in modifying structures. As heredity and adaptation enter into the closest reciprocal action under its influence, new structures, or alterations of structure, are produced, and these are purposive in the sense that they serve the organism when formed, but they were produced without any pre-conceived aim."

This simple idea is the central thought of Darwinism, or the theory of selection. Darwin conceived this idea at an early date, and then, for more than twenty years, worked at the collection of empirical evidence in support of it before he published his theory. His grandfather, Erasmus Darwin, was an able scientist of the older school of natural philosophy, who published a number of natural-philosophic works about the end of the eighteenth century. The most important of them is his *Zoonomia*, published in 1794, in which he expounds views similar to those of Goethe and Lamarck, without really knowing anything of the work of these

contemporaries. However, in the writings of the grandfather the plastic imagination rather outran the judgment, while in Charles Darwin the two were better balanced.

Darwin did not publish any account of his theory until 1859, when Alfred Russel Wallace, who had independently reached the same theory of selection, published his own work. In the following year appeared the *Origin of Species*, in which he develops it at length and supports it with a mass of proof. Wallace had reached the same conclusion, but he had not so clear a perception as Darwin of the effectiveness of natural selection in forming species, and did not develop the theory so fully. Nevertheless, Wallace's writings, especially those on mimicry, etc., and an admirable work on *The Geographical Distribution of Animals*, contain many fine original contributions to the theory of selection. Unfortunately, this gifted scientist has since devoted himself to spiritualism.

Darwin's *Origin of Species* had an extraordinary influence, though not at first on the experts of the science. It took zoologists and botanists several years to recover from the astonishment into which they had been thrown through the revolutionary idea of the work. But its influence on the special sciences with which we zoologists and botanists are concerned has increased from year to year; it has introduced a most healthy fermentation in every branch of biology, especially in comparative anatomy and ontogeny, and in zoological and botanical classification. In this way it has brought about almost a revolution in the prevailing views.

However, the point which chiefly concerns us here—the extension of the theory to man—was not touched at all in Darwin's first work in 1859. It was believed for several years that he had no thought of applying his principles to man, but that he shared the current idea of man holding a special position in the universe. Not only ignorant laymen (especially several theologians), but also a number of men of science, said very naively that Darwinism in itself was not to be opposed; that it was quite right to use it to explain the origin of the various

species of plants and animals, but that it was totally inapplicable to man.

In the meantime, however, it seemed to a good many thoughtful people, laymen as well as scientists, that this was wrong; that the descent of man from some other animal species, and immediately from some ape-like mammal, followed logically and necessarily from Darwin's reformed theory of evolution. Many of the acuter opponents of the theory saw at once the justice of this position, and, as this consequence was intolerable, they wanted to get rid of the whole theory.

The first scientific application of the Darwinian theory to man was made by Huxley, the greatest zoologist in England. This able and learned scientist, to whom zoology owes much of its progress, published in 1863 a small work entitled *Evidence as to Man's Place in Nature*. In the extremely important and interesting lectures which made up this work he proved clearly that the descent of man from the ape followed necessarily from the theory of descent. If that theory is true, we are bound to conceive the animals which most closely resemble man as those from which humanity has been gradually evolved. About the same time Carl Vogt published a larger work on the same subject. We must also mention Gustav Jaeger and Friedrich Rolle among the zoologists who accepted and taught the theory of evolution immediately after the publication of Darwin's book, and maintained that the descent of man from the lower animals logically followed from it. The latter published, in 1866, a work on the origin and position of man.

About the same time I attempted, in the second volume of my *General Animal Biology* (1866), to apply the theory of evolution to the whole organic kingdom, including man.¹ I endeavoured to sketch the probable ancestral trees of the various classes of the animal world, the protists, and the plants, as it seemed necessary to do on Darwinian principles, and as we can actually do now with a high degree of confidence. If the theory of descent, which Lamarck first clearly formulated and Darwin thoroughly established, is true, we should be able to draw up a natural classification of plants and animals in the light of their genealogy, and to conceive the large and small divisions of

¹ Darwin and Wallace arrived at the theory quite independently. *See Wallace's Contributions to the Theory of Natural Selection (1859) and Darwinism (1881).*

² Roughly spoken of this "as one of the greatest ideas the world ever published."—TENNANT.

the system as the branches and twigs of an ancestral tree. The eight genealogical tables which I inserted in the second volume of the *General Morphology* are the first sketches of their kind. In the twenty-seventh chapter, particularly, I trace the chief stages in man's ancestry, as far as it is possible to follow it through the vertebrate stem. I tried especially to determine, as well as one could at that time, the position of man in the classification of the mammals and its genealogical significance. I have greatly improved this attempt, and treated it in a more popular form, in chaps. xxvi-xxviii of my *History of Creation* (1886).¹

It was not until 1871, twelve years after the appearance of *The Origin of Species*, that Darwin published the famous work which made the much-contested application of his theory to man, and crowned the splendid structure of his system. This important work was *The Descent of Man, and Selection in Relation to Sex*. In this Darwin expressly drew the conclusion, with rigorous logic, that man also must have been developed out of lower species, and described the important part played by sexual selection in the elevation of man and the other higher animals. He showed that the careful selection which the sexes exercise on each other in regard to sexual relations and procreation, and the aesthetic feeling which the higher animals develop through this, are of the utmost importance in the progressive development of forms and the differentiation of the sexes. The males choosing the handsomest females in one class of animals, and the females choosing only the finest-looking males in another, the special features and the sexual characteristics are increasingly accentuated. In fact, some of the higher animals develop in this connection a finer taste and judgment than man himself. But, even as regards man, it is in this sexual selection that we owe the family-life, which is the chief foundation of civilisation. The rise of the human race is due for the most part to the advanced sexual selection which our ancestors exercised in choosing their mates.

Darwin accepted in the main the general outlines of man's ancestral tree, as I gave it in the *General Morphology* and the *History of Creation*, and admitted that his

studies led him to the same conclusion. That he did not at once apply the theory to man in his first work was a commendable piece of discretion; such a sequel was bound to excite the strongest opposition to the whole theory. The first thing to do was to establish it as regards the animal and plant worlds. The subsequent extension to man was bound to be made sooner or later.

It is important to understand this very clearly. If all living things come from a common root, man must be included in the general scheme of evolution. On the other hand, if the various species were separately created, man, too, must have been created, and not evolved. We have to choose between these two alternatives. This cannot be too frequently or too strongly emphasised. *Either* all the species of animals and plants are of supernatural origin—created, not evolved—and in that case man also is the outcome of a creative act, as religion teaches, or the different species have been evolved from a few common, simple ancestral forms, and in that case man is the highest fruit of the tree of evolution.

We may state this briefly in the following principle—*The descent of man from the lower animals is a special deviation which necessarily follows from the general inductive law of the whole theory of evolution.* In this principle we have a clear and plain statement of the matter. Evolution is in reality nothing but a great induction, which we are compelled to make by the comparative study of the most important facts of morphology and physiology. But we must draw our conclusion according to the laws of induction, and not attempt to determine scientific truths by direct measurement and mathematical calculation. In the study of living things we can scarcely ever directly and fully, and with mathematical accuracy, determine the nature of phenomena, as is done in the simpler study of the inorganic world—in chemistry, physics, mineralogy, and astronomy. In the latter, especially, we can always use the simplest and absolutely safest method—that of mathematical determination. But in biology this is quite impossible for various reasons; one very obvious reason being that most of the facts of the sciences are very complicated and much too intricate to allow a direct mathematical analysis. The greater part of the phenomena that biology deals with are

¹ Of which Darwin said that the *Descent of Man* would probably never have been written if he had seen it earlier.—L. AGASSI.

complicated *historical processes*, which are related to a far-reaching past, and as a rule can only be approximately estimated. Hence we have to proceed by *induction*—that is to say, to draw general conclusions, stage by stage, and with proportionate confidence, from the accumulation of detailed observations. These inductive conclusions cannot command absolute confidence, like mathematical axioms, but they approach the truth, and gain increasing probability, in proportion as we extend the basis of observed facts on which we build. The importance of these inductive laws is not diminished from the circumstance that they are looked upon merely as temporary acquisitions of science, and may be improved to any extent in the progress of scientific knowledge. The same may be said of the attainments of many other sciences, such as geology or archaeology. However much they may be altered and improved in detail in the course of time, these inductive truths may retain their substance unchanged.

Now, when we say that the theory of evolution in the sense of Lamarck and Darwin is an inductive law—in fact, the greatest of all biological inductions—we rely, in the first place, on the facts of paleontology. This science gives us some direct acquaintance with the historical phenomena of the changes of species. From the strata in which we find the fossils in the various strata of the earth we gather confidently, in the first place, that the living population of the earth has been gradually developed, as clearly as the earth's crust itself, and that, in the second place, several different populations have succeeded each other in the various geological periods. Modern geology teaches that the formation of the earth has been gradual, and unbroken by any violent revolutions. And when we compare together the various kinds of animals and plants which succeed each other in the history of our planet, we find, in the first place, a constant and gradual increase in the number of species from the earliest times until the present day; and, in the second place, we notice that the forms in each great group of animals and plants also constantly improve as the ages advance. Thus, of the vertebrates there are at first only the lower fishes; then come the higher fishes, and later the amphibians. Still later appear the three higher classes of vertebrates—the reptiles,

birds, and mammals, for the first time; only the lowest and least perfect forms of the mammals are found at first; and it is only at a very late period that placental mammals appear, and man belongs to the latest and youngest branch of these. Thus perfection of form increases as well as variety from the earliest to the latest stage. That is a fact of the greatest importance. It can only be explained by the theory of evolution, with which it is in perfect harmony. If the different groups of plants and animals do really descend from each other, we must expect to find this increase in their number and perfection under the influence of natural selection, just as the succession of fossils actually discloses it to us.

Comparative anatomy furnishes a second series of facts which are of great importance for the forming of our inductive law. This branch of morphology compares the adult structures of living things, and seeks in the great variety of organic forms the stable and simple law of organisation, or the common type or structure. Since Cuvier founded this science at the beginning of the nineteenth century it has been a favourite study of the most distinguished scientists. Even before Cuvier's time Goethe had been greatly stimulated by it, and induced to take up the study of morphology. Comparative osteology, or the philosophic study and comparison of the bony skeleton of the vertebrates—one of its most interesting sections—especially fascinated him, and led him to form the theory of the skull which I mentioned before. Comparative anatomy shows that the internal structure of the animals of each stem and the plants of each class is the same in its essential features, however much they differ in external appearance. Thus man has so great a resemblance in the chief features of his internal organisation to the other mammals that no comparative anatomist has ever doubted that he belongs to this class. The whole internal structure of the human body, the arrangement of its various systems of organs, the distribution of the bones, muscles, blood-vessels, etc., and the whole structure of these organs in the larger and the finer scale, agree so closely with those of the other mammals (such as the apes, rodents, ungulates, cetaceans, marsupials, etc.) that their external differences are of no account whatever. We learn further from comparative anatomy that the chief features of animal structure

are so similar in the various classes (fifty to sixty in number altogether) that they may all be comprised in from eight to twelve great groups. But even in these groups, the stem-forms or animal types, certain organs (especially the alimentary canal) can be proved to have been originally the same for all. We can only explain by the theory of evolution this essential unity in internal structure of all these animal forms that differ so much in outward appearance. This wonderful fact can only be really understood and explained when we regard the internal resemblance as an inheritance from common-stem forms, and the external differences as the effect of adaptation to different environments.

In recognising this, comparative anatomy has itself advanced to a higher stage. Gegenbaur, the most distinguished of recent students of this science, says that with the theory of evolution a new period began in comparative anatomy, and that the theory in turn found a touchstone in the science. "Up to now there is no fact in comparative anatomy that is inconsistent with the theory of evolution; indeed, they all lead to it. In this way the theory receives back from the science all the service it rendered to its method." Until then students had marvelled at the wonderful resemblance of living things in their inner structure without being able to explain it. We are now in a position to explain the causes of this, by showing that this remarkable agreement is the necessary consequence of the inheriting of common stem-forms; while the striking difference in outward appearance is a result of adaptation to changes of environment. Heredity and adaptation alone furnish the true explanation.

But one special part of comparative anatomy is of supreme interest and of the utmost philosophic importance in this connection. This is the science of rudimentary or useless organs; I have given it the name of "dysteleology" in view of its philosophic consequences. Nearly every organism (apart from the very lowest), and especially every highly-developed animal or plant, including man, has one or more organs which are of no use to the body itself, and have no share in its function or vital aims. That all have, in various parts of our frame, muscles which we never use, as, for instance, in the shell of the ear and adjoining parts. In most of the mammals,

especially those with pointed ears, these internal and external ear-muscles are of great service in altering the shell of the ear, so as to catch the waves of sound as much as possible. But in the case of man and other short-eared mammals these muscles are useless, though they are still present. Our ancestors having long abandoned the use of them, we cannot work them at all to-day. In the inner corner of the eye we have a small crecent-shaped fold of skin, this is the last relic of a third inner eyelid, called the nictitating (winking) membrane. This membrane is highly developed and of great service in some of our distant relations, such as fishes of the shark type and several other vertebrates, in us it is shrunken and useless. In the intestines we have a process that is not only quite useless, but may be very harmful—the vermiform appendage. This small intestinal appendage is often the cause of a fatal illness. If a cherry-stone or other hard body is unfortunately squeezed through its narrow aperture during digestion, a violent inflammation is set up, and often proves fatal. This appendix has no use whatever now in our frame, it is a dangerous relic of an organ that was much larger and was of great service in our vegetarian ancestors. It is still large and important in many vegetarian animals, such as apes and rodents.

There are similar rudimentary organs in all parts of our body, and in all the higher animals. They are among the most interesting phenomena to which comparative anatomy introduces us, partly because they furnish one of the clearest proofs of evolution, and partly because they most strikingly refute the teleology of certain philosophers. The theory of evolution enables us to give a very simple explanation of these phenomena.

We have to look on them as organs which have fallen into disuse in many generations. With the decrease in the use of its function, the organ itself shrivels up gradually, and finally disappears. There is no other way of explaining rudimentary organs. Hence they are also of great interest in philosophy, they show clearly that the material or mechanical view of the organism is the only correct one, and that the *disastrous* or teleological conception is wrong. The ancient legend of the direct creation of man according to a pre-conceived plan and the empty phrases about

"design" in the organism are completely shattered by them. It would be difficult to conceive a more thorough refutation of teleology than is furnished by the fact that all the higher animals have these rudimentary organs.

The theory of evolution finds its broadest inductive foundation in the natural classification of living things, which arranges all the various forms in larger and smaller groups, according to their degree of affinity. These groupings or categories of classification—the varieties, species, genera, families, orders, classes, etc.—show such constant features of co-ordination and subordination that we are bound to look on them as *genealogical*, and represent the whole system in the form of a branching tree. This is the genealogical tree of the variously related groups; their likeness in form is the expression of a real affinity. As it is impossible to explain in any other way the natural tree-like form of the system of organisms, we must regard it at once as a weighty proof of the truth of evolution. The careful construction of these genealogical trees is, therefore, not an amusement, but the chief task of modern classification.

Among the chief phenomena that bear witness to the inductive law of evolution we have the geographical distribution of the various species of animals and plants over the surface of the earth, and their topographical distribution on the summits of mountains and in the depths of the ocean. The scientific study of these features—the "science of distribution," or chorology (*chora* = a place)—has been pursued with lively interest since the discoveries made by Alexander von Humboldt. Until Darwin's time the work was confined to the determination of the facts of the science, and chiefly aimed at settling the spheres of distribution of the existing large and small groups of living things. It was impossible at that time to explain the causes of this remarkable distribution, or the reasons why one group is found only in one locality and another in a different place, and why there is this manifold distribution at all. Here, again, the theory of evolution has given us the solution of the problem. It furnishes the only possible explanation when it teaches that the various species and groups of species descend from common stem-forms, whose ever-branching offspring have gradually spread

themselves by migration over the earth. For each group of species we must admit a "centre of production," or common home; this is the original habitat in which the ancestral form was developed, and from which its descendants spread out in every direction. Several of these descendants become in their turn the stem-forms for new groups of species, and these also scattered themselves by active and passive migration, and so on. As each migrating organism found a different environment in its new home, and adapted itself to it, it was modified, and gave rise to new forms.

This very important branch of science that deals with active and passive migration was founded by Darwin, with the aid of the theory of evolution; and at the same time he advanced the true explanation of the remarkable relation or similarity of the living population in any locality to the fossil forms found in it. Moritz Wagner very ably developed his idea under the title of "the theory of migration." In my opinion, this famous traveller has rather over-estimated the value of his theory of migration when he takes it to be an indispensable condition of the formation of new species and opposes the theory of selection. The two theories are not opposed in their main features. Migration (by which the stem-form of a new species is isolated) is really only a special case of selection. The striking and interesting facts of chorology can be explained only by the theory of evolution, and therefore we must count them among the most important of its inductive bases.

The same must be said of all the remarkable phenomena which we perceive in the economy of the living organism. The many and various relations of plants and animals to each other and to their environment, which are treated in *bionomy* (from *nomos*, law or norm, and *bios*, life), the interesting facts of parasitism, domesticity, care of the young, social habits, etc., can only be explained by the action of heredity and adaptation. Formerly people saw only the guidance of a beneficent Providence in these phenomena; to-day we discover in them admirable proofs of the theory of evolution. It is impossible to understand them except in the light of this theory and the struggle for life.

Finally, we must, in my opinion, count among the chief inductive bases of the

theory of evolution the fatal development of the individual organism, the whole science of embryology or ontogeny. But as the later chapters will deal with this in detail, I need say nothing further here. I shall endeavour in the following pages to show, step by step, how the whole of the embryonic phenomena form a massive chain of proof for the theory of evolution; for they can be explained in no other way. In thus appealing to the close causal connection between ontogenesis and phylogenesis, and taking our stand throughout on the biogenetic law, we shall be able to prove, stage by stage, from the facts of embryology, the evolution of man from the lower animals.

The general adoption of the theory of evolution has definitely closed the controversy as to the nature or definition of the species. The word has no *absolute* meaning whatever, but is only a group-name, or category of classification, with a purely relative value. In 1857, it is true, a famous and gifted, but inaccurate and dogmatic, scientist, Louis Agassiz, attempted to give an absolute value to these "categories of classification." He did this in his *Essay on Classification*, in which he turns upside down the phenomena of organic nature, and, instead of tracing them to their natural causes, examines them through a theological prism. The true species (*bona species*) was, he said, an "incarnate idea of the Creator." Unfortunately, this pretty phrase has no more scientific value than all the other attempts to save the absolute or intrinsic value of the species.

The dogma of the fixity and creation of species lost its last great champion when Agassiz died in 1873. The opposite theory, that all the different species descend from common stem-forms, encounters no serious difficulty to-day. All the endless research into the nature of the species, and the possibility of several species descending from a common ancestor, has been closed to-day by the removal of the sharp limits that had been set up between species and varieties on the one hand, and species and genera on the other. I gave an analytic proof of this in my monograph on the sponges (1872), having made a very close study of variability in this small but highly instructive group, and shown the impossibility of making any dogmatic distinction of species. According as the classifier takes his ideas of genus, species, and variety in a broader

or in a narrower sense, he will find in the small group of the sponges either one genus with three species, or three genera with 238 species, or 113 genera with 591 species. Moreover, all these forms are so connected by intermediate forms that we can convincingly prove the descent of all the sponges from a common stem-form, the *dyothus*.

Here, I think, I have given an analytic solution of the problem of the origin of species, and so met the demand of certain opponents of evolution for an actual instance of descent from a stem-form. Those who are not satisfied with the synthetic proofs of the theory of evolution which are provided by comparative anatomy, embryology, paleontology, dyseleology, chorology, and classification, may try to refute the analytic proof given in my treatise on the sponge, the outcome of five years of assiduous study. I repeat. It is now impossible to oppose evolution on the ground that we have no convincing example of the descent of all the species of a group from a common ancestor. The monograph on the sponges furnishes such a proof, and, in my opinion, indisputable proof. Any man of science who will follow the protracted steps of my inquiry and test my assertions will find that in the case of the sponges we can follow the actual evolution of species in a concrete case. And if this is so, if we can show the origin of all the species from a common form in one single class, we have the solution of the problem of man's origin, because we are in a position to prove clearly his descent from the lower animals.

At the same time, we can now reply to the often-repeated assertion, even heard from scientists of our own day, that the descent of man from the lower animals, and proximately from the apes, still needs to be "proved with certainty." These "certain proofs" have been available for a long time, one has only to open one's eyes to see them. It is a mistake to seek them in the discovery of intermediate forms between man and the ape, or the conversion of an ape into a human being by skilful education. The proofs lie in the great mass of empirical material we have already collected. They are furnished in the strongest form by the data of comparative anatomy and embryology, completed by paleontology. It is not a question now of detecting new proofs of the evolution of man, but of examining

and understanding the proofs we already have.

I was almost alone thirty-six years ago when I made the first attempt, in my *General Morphology*, to put organic science on a mechanical foundation through Darwin's theory of descent. The association of ontogeny and phylogeny and the proof of the intimate causal connection between these two sections of the science of evolution, which I expounded in my work, met with the most spirited opposition on nearly all sides. The next ten years were a terrible "struggle for life" for the new theory. But for the last twenty-five years the tables have been turned. The phylogenetic method has met with so general a reception, and found so prolific a use in every branch of biology, that it seems superfluous to treat any further here of its validity and results. The proof of it lies in the whole morphological literature of the last three decades. But no other science has been so profoundly modified in its leading thoughts by this adoption, and been forced to yield such far-reaching consequences, as that science which I am now seeking to establish—monistic anthropogeny.

This statement may seem to be rather audacious, since the very next branch of biology, anthropology in the stricter sense, makes very little use of these results of anthropogeny, and sometimes expressly opposes them.¹ This applies especially to the attitude which has characterized the German Anthropological Society (the *Deutsche Gesellschaft für Anthropologie*) for some thirty years. Its powerful president, the famous pathologist, Rudolph Virchow, is chiefly

responsible for this. Until his death (September 5th, 1902) he never ceased to reject the theory of descent as unproven, and to ridicule its chief consequence—the descent of man from a series of mammal ancestors—as a fantastic dream. I need only recall his well-known expression at the Anthropological Congress at Vienna in 1894, that "it would be just as well to say man came from the sheep or the elephant as from the ape."

Virchow's assistant, the secretary of the German Anthropological Society, Professor Johannes Ranke of Munich, has also indefatigably opposed transformism; he has succeeded in writing a work in two volumes (*Der Mensch*), in which all the facts relating to his organization are explained in a sense hostile to evolution. This work has had a wide circulation, owing to its admirable illustrations and its able treatment of the more interesting facts of anatomy and physiology—exclusive of the sexual organs! But, as it has done a great deal to spread erroneous views among the general public, I have included a criticism of it in my *History of Craniace*, as well as met Virchow's attacks on anthropogeny.

Neither Virchow, nor Ranke, nor any other "exact" anthropologist, has attempted to give any other natural explanation of the origin of man. They have either set completely aside this "question of questions" as a transcendental problem, or they have appealed to religion for its solution. We have to show that this rejection of the rational explanation is totally without justification. The fund of knowledge—*Wissen*—accumulated in the progress of biology in the nineteenth century is quite adequate to furnish a rational explanation, and to establish the theory of the evolution of man on the solid facts of his embryology.

¹ This does not apply to English anthropologists, who are almost all evolutionists.

CHAPTER VI.

THE OVUM AND THE AMCEBA

In order to understand clearly the course of human embryology, we must select the more important of its wonderful and manifold processes for fuller explanation, and then proceed from these to the innumerable features of less importance. The most important feature in this sense, and the best starting-point for ontogenetic study, is the fact that man is developed from an ovum, and that this ovum is a simple cell. The human ovum does not materially differ in form and composition from that of the other mammals, whereas there is a distinct difference between the fertilized ovum of the mammal and that of any other animal.

This fact is so important that few should be unaware of its extreme significance, yet it was quite unknown in the first

embryonic development. Hence we must say a few words about the actual condition of the theory and the significance of the views it has suggested.

In order properly to appreciate the cellular theory, the most important element in our science, it is necessary to understand in the first place that the cell is a *unified organism*, a self-contained living being. When we anatomically dissect the fully-formed animal or plant into its various organs, and then examine the finer structure of these organs with the microscope, we are surprised to find that all these different parts are ultimately made up of the same structural element or unit. This common unit of structure is the cell. It does not matter whether we thus dissect a leaf, flower, or fruit, or a bone, muscle, gland, or bit of skin, etc.; we find in every case the same ultimate constituent, which has been called the cell since Schleiden's discovery. There are many opinions as to its real nature, but the essential point in our view of the cell is to look upon it as a self-contained or independent living unit. It is, in the words of Brücke, "an elementary organism." We may define it most precisely as the ultimate organic unit, and, as the cells are the sole active principles in every vital function, we may call them the "plastids," or "formative elements." This unity is found in both the anatomic structure and the physiological function. In the case of the protists, the entire organism usually consists of a single independent cell throughout life. But in the tissue-forming animals and plants, which are the great majority, the organism begins its career as a simple cell, and then grows into a cell-community, or, more correctly, an organized cell-state. Our own body is not really the simple unity that it is generally supposed to be. On the contrary, it is a very elaborate social system of countless microscopic organisms, a colony or commonwealth, made up of innumerable independent units, or very different tissue-cells.



FIG. 1.—The human ovum, magnified 300 times. The globular mass of yolk (a) is enclosed by a transparent membrane (the cytoplasm or zona pellucida) (b) and contains a non-central nucleus (the germinal vesicle, c). Cf. Fig. 4.


quarter of the nineteenth century. As we have seen, the human and mammal ovum was not discovered until 1857, when Carl Ernst von Beer detected it. Up to that time the larger vesicles, in which the real and much smaller ovum is contained, had been wrongly regarded as ova. The important circumstance that this mammal ovum is a simple cell, like the ovum of other animals, could not, of course, be recognized until the cell theory was established. This was not done, by Schleiden for the plant and Schwann for the animal, until 1858. As we have seen, this cell theory is of the greatest service in explaining the human frame and its

In reality, the term "cell," which existed long before the cell theory was formulated, is not happily chosen. Schleiden, who first brought it into scientific use in the sense of the cell theory, gave this name to the elementary organisms because, when you find them in the dissected plant, they generally have the appearance of chambers, like the cells in a bee-hive, with firm walls and a fluid or pulpy content. But some cells, especially young ones, are entirely without the enveloping membrane, or stiff wall. Hence we now generally describe the cell as a living, vacuous particle of protoplasm, enclosing a firmer nucleus in its albuminoid body. There may be an enclosing membrane, as there actually is in the case of most of the plants; but it may be wholly lacking, as in the case with most of the animals. There is no membrane at all in the first stage. The young cells are usually round, but they vary much in shape later on. Illustrations of this will be found in the cells of the various parts of the body shown in Figs. 3-7.

Hence the essential point in the modern idea of the cell is that it is made up of two different active constituents—an inner and an outer part. The smaller and inner part is the nucleus (or *caryon* or *embolaster*, Fig. 1c and Fig. 2b). The outer and larger part, which encloses the other, is the body of the cell (*cellous*, *plax*, or *cytosome*). The soft living substance of which the two are composed has a peculiar chemical composition, and belongs to the group of the albuminoid plasma-substances ("formative matter," or protoplasm). The essential and indispensable element of the nucleus is called nucleolus (or *caryoplasm*), that of the cell body is called plastid (or *cytoplasm*). In the most rudimentary cases both substances seem to be quite simple and homogeneous, without any visible structure. But, as a rule, when we examine them under a high power of the microscope, we find a certain structure in the protoplasm. The chief and most common form of this is the fibrous or net-like "strenid structure" (Frommann) and the feathery "honey-comb structure" (Butschli).

The shape or outer form of the cell is infinitely varied, in accordance with its endless power of adapting itself to the most diverse activities or environments. In its simplest form the cell is globular (Fig. 1a). This normal round form is especially found in cells of the simplest

structure, and those that are developed in a free fluid without any external pressure. In such cases the nucleus also is not infrequently round, and located in the centre of the cell-body (Fig. 2b). In other cases, the cells have no definite shape; they are constantly changing their form owing to their automatic movements. This is the case with the amoebae (Figs. 1g and 16) and the amoeboid travelling cells (Fig. 11), and also with very young ova (Fig. 13). However, as a rule, the cell assumes a definite form in the course of its career. In the tissues of the multicellular organism, in which a number of similar cells are bound together in virtue of certain laws of heredity, the shape is determined partly by the form of their connection and partly by their special



functions. Thus, for instance, we find in the mucous lining of our tongue very thin and delicate flat cells of roundish shape (Fig. 3). In the outer skin we find similar, but harder, covering cells, joined together by saw-like edges (Fig. 4). In the liver and other glands there are thicker and softer cells, linked together in rows (Fig. 5).

The best-named tissues (Figs. 3-5) belong to the simplest and most primitive type, the group of the "covering-tissues," or *epithelia*. In these "primary tissues" (to which the germinal layers belong) simple cells of the same kind are arranged in layers. The arrangement and shape are more complicated in the "secondary tissues," which are gradually developed out of the primary.

the muscles, nerves, bones, etc. In the bones, for instance, which belong to the group of supporting or connecting organs,

the cells (Fig. 6) are star-shaped, and are joined together by numbers of net-like interlacing processes, so, also, in the tissues of the teeth (Fig. 7), and in other forms of supporting-tissue, in which a soft or hard substance (intercellular matter, or base) is inserted between the cells.

The cells also differ very much in size. The great majority of them are invisible to the naked eye, and can be seen only through the microscope (being as a rule between $\frac{1}{100}$ and $\frac{1}{200}$ inch in diameter). There are many of the smaller plants—such as the famous bacteria—which only come into view with a very high magnifying power. On the other hand, many cells attain a considerable size, and run occasionally to several inches in diameter, as do certain kinds of rhizopods among

The *passive* portions consist third, those are subsequently formed from the others, and I have given them the name of "plasma-products." They are partly external (cell-membranes and intercellular matter) and partly internal (cell-sap and cell-contents).

The nucleus (or caryon), which is usually of a simple roundish form, is quite structureless at first (especially in very young cells), and composed of homogeneous nuclear matter or karyoplasm (Fig. 28). But, as a rule, it forms a sort of vesicle later on, in which we can distinguish a more solid *nuclear base* (*carotidic*) and a softer or fluid *nuclear sap* (*carotidoplasm*). In a mesh of the nuclear network (or it may be on the inner side of the nuclear envelope) there is, as a rule, a dark, very



FIG. 2.



FIG. 3.



FIG. 5.

FIG. 2.—Three epithelial cells from the mucous lining of the tongue.

FIG. 3.—Five spiral or grooved cells, with edges joined, from the outer skin (epidermis) one of them (B) is isolated.

FIG. 5.—Ten liver cells, one of them (B) has two nuclei.

the unicellular protists (such as the radiaria and thalamophora). Among the tissue-cells of the animal body many of the muscular fibres and nerve fibres are more than four inches, and sometimes more than a yard, in length. Among the largest cells are the yolk-filled ova, as, for instance, the yellow "yolk" in the hen's egg, which we shall describe later (Fig. 15).

Cells also vary considerably in structure. In this connection we must first distinguish between the active and passive components of the cell. It is only the former, or *active* parts of the cell, that really live, and effect that marvellous world of phenomena to which we give the name of "organic life." The first of these is the inner nucleus (*carotidic*), and the second the body of the cell (*cytoplasm*).

opaque, solid body, called the *nucleolus*. Many of the nuclei contain several of these nucleoli (as, for instance, the germinal vesicle of the ova of fishes and amphibia). Recently a very small, but particularly important, part of the nucleus has been distinguished as the *central body* (*centrosome*)—a tiny particle that is originally found in the nucleus itself, but is usually outside it, in the cytoplasm; as a rule, fine threads stream out from it in the cytoplasm. From the position of the central body with regard to the other parts it seems probable that it has a high physiological importance as a centre of movement, but it is lacking in many cells.

The cell-body also consists originally, and in its simplest form, of a homogeneous viscid plasmonic matter. But, as a rule,

only the smaller part of it is formed of the living cell-subst

.. (cytoplasm), the greater part consists of dead, passive plasma-products (metaplasm). It is useful to distinguish between the inner and outer of these. External plasma-products (which are thrust out from the protoplasm as solid "structural matter") are the cell-membranes and the intercellular matter. The internal plasma-products are either the fluid cell-sap or hard structures. As a rule, in mature and differentiated cells these various parts are so arranged that the protoplasm (like the karyoplasm in the round nucleus) forms a sort of skeleton or frame-work.

The spaces of this network are filled partly with the fluid cell-sap and partly by hard structural products.

The simple round ovum, which we take as the starting-point of our study (Figs. 1 and 2), has in many cases the vague, sensitive & an




FIG. 6.—Nine star-shaped bone-cells with interlaced branches.

itself the whole multicellular body. It is the common parent of all the countless generations of cells which form the different tissues.

their powers fully or in g to this, the neurite cell in the brain (Fig. 9) develops along one rigid line. It cannot, like the ovum, beget endless generations of cells, of which some will

the brain. The ovum stands potentially for the entire organism—in other words, it has the faculty of building up out of



all of a bunch, joined together by di

wires at a large telegraphic centre, cross, and recross in the delicate protoplasm of the nerve cell, and pass out in the branching processes which proceed from it and put it in communication with other nerve-cells or nerve-fibres (a, b). We can only partly follow their intricate paths in the fine matter of the body of the cell.

Here we have a most elaborate apparatus, the delicate structure of which we are just beginning to appreciate through

conjecture than knowledge. Its in case structure corresponds to the very complicated functions of the mind. Nevertheless, this elementary organ of psychic activity—of which there are thousands in our brain—is nothing but a single cell. Our whole mental life is only the joint result of the combined activity of all these nerve-cells, or soul-cells. In the centre of each cell there is a large transparent nucleus, containing a small and dark nuclear body. Here, as elsewhere, it is the nucleus that determines the individuality of the cell; it proves that the whole structure, in spite of its intricate composition, amounts to only a single cell.

In contrast with this very elaborate and very strictly differentiated psychic cell (Fig. 9), we have our ovum (Figs. 1 and 2), which has hardly any structure at all



FIG. 2.—Unfertilized ovum of an echinoderm (from Hertwig). The vacuolar surface (or "granular vesicle") is globular, half the size of the animal ovum and contains a nuclear framework in the central spot of which there is a dark nucleolar film. (Hertwig, 1892, p. 1.)

But even in the case of the ovum we must infer from its properties that its protoplasmic body has a very complicated chemical composition and a fine molecular structure which escapes our observation. This presumed molecular structure of the plasma is now generally admitted, but it has never been seen, and, indeed, lies far beyond the range of microscopic vision. It must not be confused—as it is often done—with the structure of the plasma (the fibrous network, granules, honey-comb, etc.) which does come within the range of the microscope.

But when we speak of the cells as the elementary organisms, or structural units, or "ultimate individualities," we must bear in mind a certain restriction of the phrase. I mean, that the cells are not,

is often supposed, the very lowest stage organic individuality. The very lowest stage of organicity to which I allude is the "cytode." There is no such thing as a "cell" in the sense in which we call the "cytode" (cyto = cell),

certain living, independent beings, consisting only of a particle of plasma—an albuminoid substance, which is not yet differentiated into cytoplasm and cytoplasm, but combines the properties of both. These remarkable beings called the *cytodes*—especially the *chironomus* and *bacterium*—are specimens of these simple cytodes. (Compare the nineteenth Chapter.) To be quite accurate, then, we must say the elementary organism, or the ultimate individual, is found in two different stages. The first and lower stage is the cytode, which consists merely of a particle of plasma, or quite simple plasma. The second and higher stage is the cell, which is already divided or differentiated into nuclear matter and cellular matter. We comprise both kinds—the cytodes and the cells—under the name of *plastids* ("formative particles"), because they are the real builders of the organism. However, these cytodes are not found, as a rule, in the higher animals and plants; here we have only real cells with a nucleus. Hence, in these tissue-forming organisms (both plant and animal) the organic unit always consists of a chemically and anatomically different parts—the outer cell-body and the nucleus.

order to conceive oneself that this cell is really an independent organism, we are only to observe the development and vital phenomena of one of them. We see then that it performs all the essential functions of life—both vegetal and animal—which we find in the entire organism. Each of these tiny beings grows and nourishes itself independently. It takes its food from the surrounding fluid, sometimes, even, the naked cells take in solid particles at certain points of their surface—in other words, "eat" them—without needing any special mouth and stomach for the purpose (cf. Fig. 19).

Further, each cell is able to reproduce itself. This multiplication, in most cases, takes the form of a simple cleavage, sometimes direct, sometimes indirect; the simple direct (or "amitotic") division is less common, and is found, for instance, in the blood cells (Fig. 10). In these the nucleus first divides into two equal parts by constriction. The indirect (or "mitotic")

THE OVUM AND THE ANCEBA




Fig. 9.—A large branching nerve-cell, or "star-cell," from the brain of an electric fish (*Torpedo*), magnified 600 times. In the middle of the cell is the large, prominent round nucleus, one nucleolus, and within the latter again, a nucleolus. The protoplasm of the cell is split into radiating fine threads (or fibrils, which are embedded in the branching processes of the cell (b). One

cleavage is much more frequent; in this the cytoplasm of the nucleus and the cytoplasm of the cell-body act upon each other in a peculiar way, with a partial dissolution (*carolysis*), the formation of knots and loops (*mitosis*), and a movement of the halved plasma-particles towards two mutually repulsive poles of attraction (*carotekinesis*, Fig. 11).

The intricate physiological processes which accompany this "mitosis" have been very closely studied of late years. The inquiry has led to the detection of certain laws of evolution which are of extreme importance in connection with heredity. As a rule, two very different parts of the nucleus play an important part in these changes. They are the *chromatin*, or coloured nuclear substance,



FIG. 11.—Blood-cells, multiplying by direct division, from the blood of the embryo of a frog. Originally each blood-cell has a nucleus and a small vac. When it is going to multiply, the nucleus divides into two (*a, b*). Then the protoplasmic bands are contracted between the two angles and these move apart from each other (*c*). Finally the constriction is completed, and the cell splits into two daughter-cells (*d, e*). (From *Proc.*)

which has a peculiar property of tinging itself deeply with certain colouring-matters (carotene, haematoxylin, etc.), and the *achromatin* (or *linin*, or *achromatic*), a colourless nuclear substance that lacks this property. The latter generally forms in the dividing cell a sort of spindle, at the poles of which there is a very small particle, also colourless, called the "central body" (*centromere*). This acts as the centre or focus in a "sphere of attraction" for the granules of protoplasm in the surrounding cell-body, and assumes a star-like appearance (the cell-star, or *monaster*). The two central bodies, standing opposed to each other at the poles of the nuclear spindle, form "the double-star" (or *amphaster*, Fig. 11, B, C). The

chromatin often forms a long, irregularly wound thread—"the coil" (*spireme*, Fig. A). At the commencement of the cleavage it gathers at the equator of the cell, between the stellar poles, and forms a crown of U-shaped loops (generally four or eight, or some other definite number). The loops split lengthwise into two halves (B), and these back away from each other towards the poles of the spindle (C). Here each group forms a crown once more, and thus, with the corresponding half of the divided spindle, forms a fresh nucleus (D). Then the protoplasm of the cell-body begins to contract in the middle, and gather about the new daughter-nuclei, and at last the two daughter-cells become independent beings.


Between this common mitosis, or *indirect* cell-division, which is the normal cleavage-process in most cells of the higher animals and plants, and the simple *direct* division (Fig. 10) we find every grade of segmentation; in some circumstances even one kind of division may be converted into another.

The plastid is also endowed with the functions of movement and sensation. The single cell can move and creep about, when it has space for free movement and is not prevented by a hard envelope; it then thrusts out at its surface processes like fingers, and quickly withdraws them again, and thus changes its shape (Fig. 12). Finally, the young cell is sensitive, or more or less responsive to stimuli; it makes certain movements on the application of chemical and mechanical irritations. Hence we can ascribe to the individual cell all the chief functions which we comprehend under the general heading of "life"—sensation, movement, nutrition, and reproduction. All these properties of the multicellular and highly developed animal are also found in the single animal-cell, at least in its younger stages. There is no longer any doubt about this, and so we may regard it as a solid and important basis of our physiological conception of the elementary organism.

Without going any further here into these very interesting phenomena of the life of the cell, we will pass on to consider the application of the cell theory to the ovum. Here comparative research yields the important result that *every ovum is at first a simple cell*. I say this is very important, because our whole science of embryology now resolves itself into the problem: "How does the multicellular

organism arise from the unicellular?" Every organic individual is at first a simple cell, and as such an elementary organism, or a unit of individuality. This cell produces a cluster of cells by segmentation, and from these develop the multicellular organisms, or individual of higher rank.

internal constitution. Later, though the ova remain unicellular, they differ in size and shape, enclose various kinds of yolk-particles, have different envelopes, and so on. But when we examine them at their birth, in the ovary of the female animal, we find them to be always of the same form in the first stages of their life. In



When we examine a little closer the original features of the ovum, we notice that, in its first stage the ovum is just the same simple and indefinite structure in the case of man and all the animals (Fig. 13). We are unable to detect any material difference between them, either in outer shape or


the beginning each ovum is a very simple, roundish, naked, mobile cell, without a nucleus, is composed merely of a particle of cytoplasm enclosing a nucleus (Fig. 13). Special names have been given to these parts of the ovum; the cell-body is called the *yolk (vitellus)*, and the cell-nucleus the *germinal vesicle*. As a rule, the

nucleus of the ovum is soft, and looks like a small purple or vesicle.

... many other cells.

nuclear skeleton or frame and a third, hard nuclear body (the *nucleolus*). In the ovum this is called the *germinal spot*. Finally, we find in many ova (but not in all) a still further point within the germinal spot, a "nucleolin," which goes by the name of the *germinal point*. The latter parts (germinal spot and germinal point) have, apparently, a minor importance, in comparison with the other two (the yolk and germinal vesicle) in the

so must distinguish the active *first yolk* (or *protoplasm* = *first plasma*) the passive *nurturing yolk* (or *deutoplasm*).



14.—Mobile cells from the inflamed eye of

1. From the watery fluid of the eye. (See Hamster 2.)

The naked cells creep freely about, by (like the amoeba, or rheopod) protruding fine processes from the uncontracted protoplasmic body. These bodies vary continually in number, shape, and size. The nucleus of these amoeboid "protoplasts" ("traveling cells," or *plasmocytes*) is usually brown surrounded by the nucleus of fine granules which are scattered in the protoplasm. (From Frey)

ity of the lower animals (such as sponges, polyps, and medusae) the naked ova retain their original simple appearance until impregnation. But in most animals they at once begin to change; the change consists partly in the formation of connections with the yolk, which serve to nourish the ovum, and partly of external membranes for their protection (the *ovolemma*, or *prochorion*). A membrane of this sort is formed in all the mammals in the course of the embryonic process. The little globule is surrounded by a thick capsule of glass-like transparency, the *zona pellucida*, or *ovolemma*

pellucidum (Fig. 14). When we examine it closely under the microscope, we

ally we

whether fertile

not, cannot be distinguished from that of most of the other mammals. It is nearly the same everywhere in form, size, and composition. When it is fully formed, it has a diameter of (on an average) about $\frac{1}{16}$ of an inch. When the mammal ovum has been carefully isolated, and held against the light on a glass-plate, it

naked eye. The ova of the higher mammals are about the same size. The diameter of the ovum is almost always

It has always the same globular shape, the same characteristic membrane, the same transparent germinal vesicle with its dark germinal spot. Even when we use the most powerful microscope with its highest power, we can detect no material difference between the ova of man, the ape, the dog, and so on. I do not mean to say that there are no differences between the ova of these different mammals. On the contrary, we are bound to assume that there are such, at least as regards chemical composition. Even the ova of different men must differ from each other; otherwise we should not have a different individual from each ovum. It is true that our crude and imperfect apparatus cannot detect these subtle individual differences, which are probably in the molecular structure. However, such a striking resemblance of their ova in form, so great as to

as and the other mammals. From non germ-form we infer a common form. On the other hand, they

easily distinguish the mammal from the fertilised ovum of the birds, amphibians, fishes, and other vertebrates (see the close of the twenty-ninth chapter).

The fertilised bird-ovum (Fig. 15) is notably different. It is true that in its earliest stage (Fig. 15 E) this ovum also is very like that of the mammal (Fig. 15 A). But afterwards, while still within the oviduct, it takes up a quantity of nourishment and works this into the familiar large yellow yolk. When we examine a very young ovum in the hen's oviduct, we

find it to be a simple, small, naked, amoeboid cell, just like the young ova of other animals (Fig. 13). But it then grows to the size we are familiar with in the round yolk of the egg. The nucleus of the ovum, or the germinal vesicle, is thus pressed right to the surface of the globular ovum, and is embedded there in a small quantity of transparent matter, the so-called white yolk. This forms a round white spot, which is known as the "tread" (*eccentricale*) (Fig. 15 *b*). From

the tread a thin column of the white yolk penetrates through the yellow yolk to the centre of the globular cell, where it swells into a small, central globule (wrongly called the yolk-cavity, or *latebra*, Fig. 15 *d*). The yellow yolk-matter which surrounds this white yolk has the appearance in the egg (when boiled hard) of concentric layers (*c*). The yellow yolk is also enclosed in a delicate structureless membrane (the *membrana vitellina*, *d*).

As the large yellow ovum of the bird




FIG. 15.—Ova of various animals, exhibiting amoeboid movements, highly magnified. All the ova are noted cells of varying shape. In the dark background protoplasm (yolk) is a large vesicular nucleus (the germinal vesicle), and in this is seen a nuclear body (the germinal spot), in which again we often see a germinal point. Figs. A1-A4 represent the ovum of a sponge (*Leucosphaera echinaria*) at four successive moments. B1-B4 are the ovum of a paramete ova (*Chondrocythoe serratus*), at eight successive moments. (From Edmund van Beneden.) C1-C5 show the ovum of the cat at various stages of movement (from Pfeffer); Fig. F the ovum of a trout; G the ovum of a shad; H a human ovum.

attains a diameter of several inches in the bigger birds, and encloses round yolk-particles, there was formerly a reluctance to consider it as a simple cell. This was a mistake. Every animal that has only one cell-nucleus, every amoeba, every graminia, every infusorian, is unicellular, and remain unicellular whatever variety of matter it feeds on. So the ovum remains a simple cell here.



FIG. 14.—The human ovum, taken from the female ovary, magnified 300 times. The whole ovum is a single round cell. The chief part of the granular mass is formed by the nuclear yolk (nucleoplasm), which is evenly distributed in the active protoplasm and consists of numbers of fine yellow granules. In the upper part of the yolk is the transparent round germinal vesicle, which corresponds to the nucleus. The nucleus is darker granular, the germinal vesicle which encloses a nucleolus. The granular yolk is surrounded by the thick transparent germinal membrane (zona pellucida, or zona follicularis). This is traversed by numbers of lines as fine as hairs, which are directed radially towards the centre of the ovum. These are called the yolk-canals, it is through these that the streaming cytoplasm penetrates into the yolk at impregnation.

ever much yellow yolk it afterward accumulates within its protoplasm. It is, of course, different, with the bird's ovum — *as has been* — has been fertilised. The ovum then consists of as many cells as there are nuclei in the tread. Hence, in the fertilised egg which we eat daily, the yellow yolk is already a multicellular body. Its tread is composed of several cells, and is now commonly called the

germinal disc. We shall return to this *discogerminal* in the sixth chapter.

When the mature bird-ovum has left the ovary and been fertilised in the oviduct, it covers itself with various membranes which are secreted from the wall of the oviduct. First, the large clear albuminous layer is deposited around the yellow yolk; afterwards, the hard external shell with a fine inner skin. All these

gradually forming envelopes and processes are of no importance in the formation of the embryo; they serve merely for the protection of the original simple ovum. We sometimes

very large

of other animals, such as fishes of the shark type. Here, also, the ovum is originally of the same character as it is in the mammal, it is a perfectly simple and naked cell. But, as in the case of the bird, a considerable quantity of nutritive yolk is accumulated inside the original yolk as food for the developing embryo; and various coverings are formed round the egg. The ovum of many other animals has the same internal and external features. They have, however, only a physiological, not a morphological, importance; they have no direct influence on the formation of the foetus. They are partly consumed as food by the embryo, and partly serve as protective envelopes. Hence

we may leave them out of consideration altogether here, and restrict ourselves to material points—to the substantial identity of the original ovum in man and the rest of the animals (Fig. 13).

Now, let us for the first time make use of our biogenetic law, and directly apply this fundamental law of evolution to the human ovum. We reach a very simple, but very important, conclusion. From

the fact that the human ovum and that of all other animals consists of a single cell, it follows immediately, according to the biogenetic law, that all the animals, including man, descend from a unicellular organism.



FIG. 13.—A fertilized ovum from the oviduct of a hen. The yellow yolk (*c*) consists of several concentric layers (*a*), and is enclosed in a thin yellow membrane (*b*). The nucleus or germinal vesicle is seen above in the center or "beard" (*d*). From that point the whole yolk penetrates to the central yolk cavity (*e*). The two kinds of yolk do not differ very much.

If our biogenetic law is true, if the embryonic development is a summary or condensed recapitulation of the stem-history—and there can be no doubt about it—we are bound to conclude, from the fact that all the ova are at first simple cells, that all the multicellular organisms originally sprang from a unicellular being. And as the original ovum in man and all the other animals has the same simple and indefinite appearance, we may assume with some probability that this unicellular stem-form was the common ancestor of the whole animal world, including man. However, this last hypothesis does not seem to me as inevitable and as absolutely certain as our first conclusion.

This inference from the unicellular embryonic form to the unicellular ancestor is so simple, but so important, that we cannot sufficiently emphasize it. We must, therefore, turn next to the question whether there are to-day any unicellular organisms, from the features of which we may draw some approximate conclusion as to the unicellular ancestors of the multicellular organisms. The answer is: Most

certainly there are. There are still unicellular organisms which are, in their whole nature, really nothing more than permanent ova. There are independent unicellular organisms of the simplest character which develop no further, but reproduce themselves as such, without any further growth. We know

to-day of a great number of these little beings, such as the *grogarium*, flagellata, actineta, infusoria, etc. However, there is one of them that has an especial interest for us, because it at once suggests itself when we raise our question, and it must be regarded as the unicellular being that approaches nearest to the real ancestral form. This organism is the *Amoeba*.

For a long time now we have comprised under the general name of amoebae a number of microscopic unicellular organisms, which are very widely distributed, especially in fresh water, but also in the ocean; in fact, they have lately been discovered in damp soil. There are also parasitic amoebae which live inside other animals. When we place one of these amoebae in a drop of water under the microscope and examine it with a high power, it generally appears as a roundish particle of a very irregular and varying shape (Figs. 16 and 17) slimy, semi-fluid substance, which consists of protoplasm, we see only the solid globular particle it contains, the nucleus. This unicellular body moves about continually, creeping in every direction on the glass on which we are examining it. The movement is effected by the shapeless body thrusting out finger-like processes at various parts of its surface, and these are slowly but continually changing, drawing the rest of the body after them. After a time, perhaps, the action changes—



FIG. 16.—A creeping amoeba (highly magnified). The whole organism is a simple animal cell, and moves about by means of the changing an "arms" it thrusts out of and withdraws into the protoplasmic body. Inside it is the central nucleus with its nucleolus.

The amoeba suddenly stands still, withdraws its projections, and assumes a globular shape. In a little while, however, the round body begins to expand again, thrusts out arms in another

direction, and moves on once more. These changeable processes are called "false feet," or pseudopodia, because they act physiologically as feet, yet are not special organs in the anatomic sense. They disappear as quickly as they come, and are nothing more than temporary projections of the semi-fluid and structureless body.

If you touch one of these creeping amoebae with a needle, or put a drop of acid in the water, the whole body at once contracts in consequence of this mechanical

which it comes in contact. The latter process may be observed at any moment by forcing it to eat. If finely ground colouring matter, such as carmine or indigo, is put into the water, you can see the body of the amoeba prodding these coloured particles into itself, the substance of the cell closing round them. The amoeba can take in food in this way at any point on its surface, without having any special organs for intussusception and digestion, or a real mouth or gut.

The amoeba grows by thus taking in food and dissolving the particles eaten in its protoplasm. When it reaches




FIG. 17.—Division of a unicellular amoeba (*Amoeba proteus*) in its vegetative stage. (From P. G. Schuster.) The dark spot is the nucleus, the lighter spot is contractile vacuole in the protoplasm. The latter redivides in one of the daughter-cells.

reproduce. This is done by the simple process of cleavage (Fig. 17). First, the nucleus divides into two parts. Then the protoplasm is separated between the two new nuclei, and the whole cell splits into two daughter-cells, the protoplasm gathering about each of the nuclei. The thin bridge of protoplasm which at first connects the daughter-cells soon breaks. Here we have the simple form of direct cleavage of the nuclei. Without mitosis, or formation of threads, the homogeneous nucleus divides into two halves. These move away from each other, and become centres of attraction for the enveloping matter, the protoplasm. The same direct cleavage of the nuclei is also witnessed in the reproduction

other protists, while other

rule, the body
shape. In
case, if the

impurity of the water lasts some time, the amoeba begins to develop a covering. It exudes a membrane or capsule, which immediately hardens, and assumes the appearance of a round cell with a protective membrane. The amoeba either takes its food directly by imbibition of matter floating in the water, or by prodding into its protoplasmic body solid particles with

unicellular organisms show the indirect division of the cell.

Hence, although the amoeba is nothing but a simple cell, it accomplishes all the functions of the multicellular organism. It moves, feels, nourishes itself, and reproduces. Some kinds of these amoebae can be seen with the naked eye, but most of them are microscopically small. It is for the following reasons that we regard the amoebae as the unicellular organisms which have

special phylogenetic (or evolutionary) relations to the ovum. In many of the lower animals the ovum reverts its original naked form until fertilisation, develops no membranes, and is then often indistinguishable from the ordinary amoeba. Like the amoeba, these naked ova may thrust out processes, and move about as travelling cells. In the sponges, these mobile ova move about freely in the maternal body like independent amoebae (Fig. 17). They had been observed by earlier scientists, but described as foreign bodies—namely, parasitic amoebae, living parasitically on the body of the sponge. Later, however, it was discovered that they were not parasites, but the ova of the sponge. We also find this remarkable phenomenon among other animals, such as the graceful, bell-shaped zoophytes, which we call polyps and medusae. Their ova remain naked cells, which thrust out amoeboid projections, nourish themselves, and move about. When they have been fertilised, the multicellular organism is formed from them by repeated segmentation.

It is, therefore, no audacious hypothesis, but a perfectly sound conclusion, to regard the amoeba as the particular unicellular organism which offers us an approximate illustration of the ancient common unicellular ancestor of all the metazoa, or multicellular animals. The simple naked amoeba has a less definite and more original character than any other cell. Moreover, there is the fact that recent research has discovered such amoeba-like cells everywhere in the mature body of the multicellular animals. They are found, for instance, in the human blood, side by side with the red corpuscles, as colourless blood-cells, and it is the same with all the vertebrates. They are also found in many of the invertebrates—for instance, in the blood of the snail. I showed, in 1899, that these colourless blood-cells can, like the independent amoebae, take up solid particles, or "eat" (whence they are called *phagocytes* = "eating-cells," Fig. 19). Lately, it has been discovered that many different cells may, if they have room enough, execute the same movements, creeping about and eating. They behave just like amoebae (Fig. 12). It has also been shown that these "travelling-cells," or *plasmocytes*, play an important part in man's physiology and pathology (as means of transport for food, infectious matter, bacteria, etc.).

The power of the naked cell to execute these characteristic amoeba-like movements comes from the contractility (or automatic mobility) of its protoplasm. This seems to be a universal property of young cells. When they are not enclosed by a firm membrane, or confined in a "cellular prison," they can always accomplish these amoeboid movements. This is true of the naked ova as well as of any other naked cells, of the "travelling-cells," of various kinds in connective tissue, lymph-cells, mucus-cells, etc.

We have now, by our study of the ovum and the comparison of it with the amoeba, provided a perfectly sound and most valuable foundation for both the embryology and the evolution of man. We have learned that the human ovum is a simple cell, that this ovum is not materially different from that of other



FIG. 15.—OVUM of a sponge (*Obolus*). The ovum creeps about in the body of the sponge by thrusting out ever-changing processes. It is indistinguishable from the common amoeba.

mammals, and that we may infer from it the existence of a primitive unicellular ancestral form, with a substantial resemblance to the amoeba.

The statement that the earliest progenitors of the human race were simple cells of this kind, and led an independent unicellular life like the amoeba, has not only been ridiculed as the dream of a natural philosopher, but also been violently censured in theological journals as "shameful and immoral." But, as I observed in my essay *On the Origin and Ancestral Tree of the Human Race* in 1870, this offended piety must equally protest against the "shameful and immoral" fact that each human individual is developed from a simple ovum, and that this human ovum is indistinguishable from those of the other mammals, and in its earliest stage is like a naked amoeba.

We can show this to be a fact any day with the microscope, and it is little use to close one's eyes to "immoral" facts of this kind. It is as indisputable as the momentous conclusions we draw from it and as the vertebrate character of man (see Chapter XI).

We now see very clearly how extremely important the cell theory has been for our whole conception of organic nature. "Man's place in nature" is settled beyond

can understand how the elaborate mind of the higher vertebrates, and especially of man, was gradually evolved from them. The academic psychologists who lack this zoological equipment are unable to do so.

This naturalistic and realistic conception is a stumbling-block to our modern idealistic metaphysicians and their theological colleagues. Fenced about with their transcendental and dualistic prejudices, they attack not only the material system as established on our scientific knowledge, but even the plainest facts which go to form its foundation. An instructive instance of this was seen a few years ago, in the academic discourse delivered by a distinguished theologian, Wilhelm Reyschlag, at Halle, January 12th, 1900, on the occasion of the centenary festival.



FIG. 19.—Blood-cells that eat, or phagocytes, from a naked sea-slug (*Aplysia*, greatly magnified). It was the first to observe in the blood-cells of this animal the important fact that "the blood-cells of the invertebrates are unspecialized pieces of plasma, and take as food, by means of their peculiar movements, like the amoebae. I had (in Naples, on May 26th, 1894) injected into the blood-vessels of one of these animals an infusion of water and ground indigo, and was greatly astonished to find the blood-cells themselves more or less filled with the particles of indigo after a few hours. After repeated injections I succeeded in "observing the very entrance of the coloured particles in the blood-cells, which took place just in the same way as with the amoebae. I have given further particulars about this in my *Monograph on the Radiolarians*."

question by it. Apart from the cell theory, man is an insoluble enigma to us. Hence philosophers, and especially physiologists, should be thoroughly conversant with it. The soul of man can only be really understood in the light of the cell-soul, and we have the simplest form of this in the amceba. Only those who are acquainted with the simple psychic functions of the unicellular organisms and their gradual evolution in the series of lower animals

can understand how the elaborate mind of the higher vertebrates, and especially of man, was gradually evolved from them. The academic psychologists who lack this zoological equipment are unable to do so. This naturalistic and realistic conception is a stumbling-block to our modern idealistic metaphysicians and their theological colleagues. Fenced about with their transcendental and dualistic prejudices, they attack not only the material system as established on our scientific knowledge, but even the plainest facts which go to form its foundation. An instructive instance of this was seen a few years ago, in the academic discourse delivered by a distinguished theologian, Wilhelm Reyschlag, at Halle, January 12th, 1900, on the occasion of the centenary festival. "Man's place in nature" is settled beyond question by it. Apart from the cell theory, man is an insoluble enigma to us. Hence philosophers, and especially physiologists, should be thoroughly conversant with it. The soul of man can only be really understood in the light of the cell-soul, and we have the simplest form of this in the amceba. Only those who are acquainted with the simple psychic functions of the unicellular organisms and their gradual evolution in the series of lower animals

idea that the greatest of men, Luther and Christ, were descended from a mere globule of protoplasm." Nevertheless, not a single informed and impartial scientist doubts the fact that these great men were, like all other men—and all other vertebrates—developed from an impregnated ovum, and that this simple nucleated globule of protoplasm has the same chemical constitution in all the mammals.

CHAPTER VII.

CONCEPTION

THE recognition of the fact that every man begins his individual existence as a simple cell is the solid foundation of all research into the genesis of man. From this fact we are forced, in virtue of our biogenetic law, to draw the weighty phylogenetic conclusion that the earliest ancestors of the human race were also unicellular organisms; and among these protozoa we may single out the vague form of the amoeba as particularly important (cf. Chapter VI). That these unicellular amoeboid forms did once exist follows directly from the phenomena which we perceive every day in the fertilised ovum. The development of the multicellular organism from the ovum, and the formation of the germinal layers and the tissues, follow the same laws in man and all the higher animals. It will, therefore, be our next task to consider more closely the integrated ovum and the process of conception which produces it.

The process of impregnation or sexual conception is one of those phenomena that people love to conceal behind the mystic veil of supernatural power. We shall soon see, however, that it is a purely mechanical process, and can be reduced to familiar physiological functions. Moreover, this process of conception is of the same type, and is effected by the same organs, in man as in all the other mammals. The pairing of the male and female has in both cases for its main purpose the introduction of the ripe matter of the male seed or sperm into the female body, in the sexual canals of which it encounters the ovum. Conception then uses by the blending of

We must observe, first, that this important process is by no means so widely distributed in the animal and plant world as is commonly supposed. There is a very large number of lower organisms which propagate asexually, or by monogony; these are especially the sexless amoebae, (chromonades, bacteria, etc.), but also many other protists, such as the amoebae, foraminifera, radiolaria,

myxomycetes, etc. In these the multiplication of individuals takes place by unsexual reproduction, which takes the form of cleavage, budding, or sporulation. The copulation of two coalescing cells, which in these cases often precedes the reproduction, cannot be regarded as a sexual act unless the two copulating plastids differ in size or structure. On the other hand, sexual reproduction is the general rule with all the higher organisms, both animal and plant; very rarely do we find asexual reproduction among them. There are, in particular, no cases of parthenogenesis (virginal conception) among the vertebrates.

Sexual reproduction offers an infinite variety of interesting forms in the different classes of animals and plants, especially as regards the mode of conception, and the conveyance of the spermatozoon to the ovum. These features are of great importance not only as regards conception itself, but for the development of the organic form, and especially for the differentiation of the sexes. There is a particularly curious correlation of plants and animals in this respect. The splendid studies of Charles Darwin and Hermann Muller on the fertilisation of flowers by insects have given us very interesting particulars of this.* This reciprocal service has given rise to a most intricate sexual apparatus. Equally elaborate structures have been developed in man and the higher animals, serving partly for the isolation of the sexual products on each side, partly for bringing them together in conception. But, notwithstanding these phenomena are in themselves, we cannot go into them here, as they have only a minor importance—if any at all—in the real process of conception. We must, however, try to get a very clear idea of this process and the meaning of sexual reproduction.

* See Darwin's work, *On the Forces by which Grains are Fertilised* (1862).

In every act of conception we have, as I said, to consider two different kinds of cells—a female and a male cell. The female cell of the animal organism is always called the ovum (or *ovule*, egg, or egg-cell); the male cells are known as the sperm or seed-cells, or the spermatozoa (also spermium and zoospermium). The ripe ovum is, on the whole, one of the largest cells we know. It attains colossal dimensions when it absorbs great quantities of nutritive yolk, as is the case with birds and reptiles and many of the fishes. In the great majority of the animals the ripe ovum is rich in yolk and much larger than the other cells. On the other hand, the next cell which we

Firstly, they are extraordinarily small, being usually the smallest cells in the body; and, secondly, they have, as a rule, a peculiarly lively motion, which is known as spermatozoic motion. The shape of the cell has a good deal to do with this motion. In most of the animals, and also in many of the lower plants (but not the higher), each of these spermatozoa has a very small, naked cell-body, enclosing an elongated nucleus, and a long thread hanging from it (Fig. 20). It was long before we could recognise that these structures are simple cells. They were formerly held to be special organisms, and were called "seed animals" (spermato-zoa, or spermato-




FIG. 20.—Spermia or spermatozoa of various animals. The pear-shaped flattened nucleus is seen from the front as *f*, and sideways as *f'*. *S* is the nucleus, *m* = *f'*, part (peritoplasm), *x* the middle, tapering end (or whip), *N* from human spermatozoa, *A* four spermatozoa from the ape, *K* from the cat, *H* from the mouse, *C* from the dog. *S* " "

have to consider in the process of conception, the male sperm-cell or spermatozoon, is one of the smallest cells in the animal body. Conception usually consists in the bringing into contact with the ovum of a shiny fluid secreted by the male, and this may take place either inside or out of the female body. This fluid is called sperm, or the male seed. Sperm, like saliva or blood, is not a simple fluid, but a thick agglomeration of innumerable cells swimming about in a comparatively small quantity of fluid. It is not the fluid, but the independent male cells that swim in it, that cause conception.

The spermatozoa of the great majority of animals have two characteristic features.

Firstly, they are now scientifically known as *sperms* or *spermata*, or as *spermatozoids* (seed-bodies) or *spermatozoa* (seed threads). It took a good deal of comparative research to convince us that each of these spermatozoa is really a simple cell. They have the same shape as in many other vertebrates and most of the invertebrates. However, in many of the lower animals they have quite a different shape. Thus, for instance, in the craw fish they are large round cells, without any movement, equipped with stiff outgrowths like bristles (Fig. 21 *f*). They have also a peculiar form in some of the worms, such as the thread-worms (*Nematodes*); in this case they are seen

amoeboid and like very small ova (Fig. 21 c-e). But in most of the lower animals (such as the sponges and polyps) they have the same pine-cone shape — and the other mammals (Fig. a, b).



FIG. 21.—Spermatozoa or spermella of various animals. (From Leag.) a of a sea slug, b of a sea slug, c of a sea slug, d of a sea slug. (Note the shape of the spermatozoa from a sea slug.)

1. Dutch naturalist Leag

hoek discovered these thread-like lively particles in 1677 in the male sperm. It was generally believed that they were special, independent, tiny animalcules, like the infusoria, and that the whole mature organism existed already, with all its parts, but very small and packed together, in each spermatozoon (see p. 12). We now know that the mobile spermatozoa are nothing but simple and real cells, of the kind that we call "ciliated" (equipped with lashes, or cilia). In the previous illustrations we have distinguished in the spermatozoon a head, trunk, and tail. The "head" (Fig. 20 b) is merely the oval nucleus of the cell; the body or middle-part (m) is an accumulation of cell-matter; and the tail (x) is a thread-like prolongation of the same.

Moreover, we now know that these spermatozoa are not at all a peculiar form of cell; precisely similar cells are found in various other parts of the body. If they have many short threads projecting, they are called *ciliated*; if only one long, whip-shaped process (or, more rarely, two or four), *caudate* (tailed) cells.

Very careful recent examination of the sperm, under a very high microscopic power (Fig. 22 a, b), has detected some further details in the finer structure of the

ciliated cell, and these are common to man and the anthropoid ape. The head (h) encloses the elliptic nucleus in a thin envelope of cytoplasm; it is a little flattened on one side, and thus looks rather pear-shaped from the front (b). In the central piece (m) we can distinguish a short neck and a longer connective piece (with central body). The tail consists of a long main section (l) and a short, very fine tail (x).

The process of fertilization by sexual conception consists, therefore, essentially in the coalescence and fusing together of two different cells. The lively spermatozoon travels towards the ovum by its serpentine movements, and bores its way into the female cell (Fig. 23). The nuclei of both sexual cells, attracted by a certain "affinity," approach each other and melt into one.

Is quite a

unfertilized cell. For if we must regard the sperm as real cells no less than the ova, and the process of conception of the two

under the microscope, it bears in and independent organism.

It bears in the cell and nuclear matter of the penetrating spermatozoa a part of the father's body, and in the protoplasm and karyoplasm of the ovum a part of the mother's body. This is clear from the fact that the child inherits many features from both parents. It inherits from the father by means of the spermatozoon, and from the mother by means of the ovum. The




FIG. 22.—A single human spermatozoon magnified 1000 times. a shows it from the broader and b from the narrower side. h head (with nucleus), m middle-piece, l long-section, and x tail. (From Leag.)

actual blending of the two cells produce a third cell, which is the germ of a child, or the new organism conceived. One

is that the stem-cell is a *single* *prokaryote*; it unites both sexual and asexual in itself.

I think the fundamental importance of this, but often unappreciated, feature in order to have a correct and clear idea of conception. With that end, I have given a special name to the new cell from which the child develops, and which is generally loosely called "the fertilized ovum," or "the first segmentation sphere." I call it "the stem-cell" (*cytula*). The name "stem-cell" seems to me the simplest and most suitable, because all the other cells of the body are derived



FIG. 25.—The Fertilized

it, and because

mother and stem-mother of all the countless generations of cells of which the multicellular organism is to be composed. That complicated molecular movement of the protoplasmic life

thing quite different

what we find in the two parent-cells, from the coalescence of which it has issued. The life of the stem-cell or cytula is the product or resultant of the paternal life-movement that is conveyed in the spermatozoon and the maternal life-movement that is contributed by the ovum.

The admirable work done by

animals, coincidences with the formation

of a simple "stem-cell" of this cell and that this then passes, by repeated segmentation (or cleavage),

known as "the segmentation sphere" or "segmentation cell." The process is most clearly observed in the ova of the echinoderms (star-fishes, sea-urchins, etc.). The investigators Oscar and Richard Hertwig were chiefly directed to these. The main results may be summed up as follows:—

Conception is preceded by certain preliminary changes, which are very necessary—in fact, usually indispensable—for its occurrence. They are comprised under the general heading of "Changes prior to impregnation." In these the original nucleus of the ovum, the germinal vesicle, is lost. Part of it is extruded, and part dissolved in the cell contents; only a very small part of it is left to form the basis of a fresh nucleus, the *pro-nucleus feminae*. It is the latter alone that combines in conception with the invading nucleus, the fertilizing spermatozoon (the *pro-nucleus masculinus*).

The impregnation of the ovum commences with a decay of the germinal vesicle, or the original nucleus of the ovum (Fig. 8). We have seen that this is in most unripe ova a large, transparent, round vesicle. This germinal vesicle contains a viscous fluid (the *caryolymph*). The first nuclear fringe (*caryoblast*) is formed of the enveloping membrane and a mesh-work of nuclear threads running across the interior, which is filled with the nuclear sap. In a knot of the network is contained the dark, stiff, opaque nuclear corpuscle or nucleolus. When the impregnation of the ovum sets in, the greater part of the germinal vesicle is dissolved in the cell, the nuclear membrane and mesh-work disappear, the nuclear sap is distributed in the protoplasm, a small portion of the nuclear base is extruded, another small

secondary

female pro-nucleus (Fig. 24 c & d).

The small portion of the nuclear base which is extruded from the impregnated ovum is known as the "directive bodies" or "polar cells"; there are many disputes as to their origin and significance, but

they

round granules, of the appearance as the remaining pro-nucleus. They are detached cell-buds; their separation from the large mother-cell takes

place in the same way as in ordinary "indirect cell-division." Hence, the polar cells are probably to be conceived as "abortive ova," or "rudimentary ova," which proceed from a single ovum by cleavage in the same way that several sperm-cells arise from one "sperm-mother-cell," in reproduction from sperm. The male sperm-cells in the testicles must undergo similar changes in view of the

the ova in the maturing of the each of the original seed-cells by double segmentation into four daughter-cells, each furnished with fourth of the original nuclear matter (the hereditary chromatin), and each of those

pendant cells become a *spermatozoon*, ready for impregnation. Thus is prevented the doubling of the chromatin in the coalescence of the two nuclei at conception. As the two polar cells are extruded and lost, and have no further part in the fertilisation of the ovum, we need not discuss them any further. But we must give more attention to the female pro-nucleus which alone remains after the polar cells and the divisional vesicle (Fig. 23 c & d) corpuscle of chromatin

male pro-nucleus. The product of this blending, which is the most important part of the act of impregnation, is the

sequent embryonic processes.

Hertwig has shown that the tiny transparent ova of the echinoderms are the

of this important
We can, in this
fully accomplish
and follow the

in by step within the space of ten minutes. If we put ripe ova of the starfish

fluid, we find each ovum within five minutes. Though fine, described as "sperm-threads" (Fig. 20), make their way to the ova, owing to a sort of chemical sensitive action which may be called "smell." But only one of

these innumerable spermatozoa is chosen—namely, the one that first reaches the ovum by the serpentine motions of its tail, and touches the ovum with its head.

the spot where the touches the surface of the ovum the protoplasm of the latter is raised in the form of a small wart, the "impregnation rise" (Fig. 25 A). The spermatozoon then bores its way into this with its head, a tail outside wriggling about all the time (Fig. 25 B, C). Presently the tail is disappears within the ovum. At the same time the ovum secretes a thin external yolk-membrane (Fig. 25 C), starting from the point of impregnation, and this prevents any more spermatozoa

inside the impregnated cell a rapid series of

The pear-shaped head of the




FIG. 24.—An impregnated echinoderm ovum with male hemispherical nucleus (x4). (From

son," grows larger and round is converted into the male pro-nucleus (Fig. 26 a & b). This has an influence on the fine granules

They move towards each other inside the yolk with increasing speed, the male nucleus moves more outside

male nucleus takes with it the radiating protoplasm which spreads like a star about the male nucleus. The male nucleus (usually in the centre of the globular ovum), lies close together, are flattened at the points of contact, and coalesce into a common mass. The small central particles of

nucleus which is formed from this combination of the nuclei is the stem-nucleus or the first segmentation nucleus; the new-formed cell, the product of the impregnation, is our stem-cell, or "first segmentation sphere" (Fig. 2).

Hence the one essential point in the process of sexual reproduction is the impregnation of the egg-cell, the stem-cell, by the combination of two originally different cells, the female ovum and the male spermatozoon. This process is of the highest importance, and merits our closest attention; all that happens in the later development of this first cell and in the life of the organism that comes of it is determined from the first by the chemical and morphological composition of the stem-cell, its nucleus and its body. We must, therefore, make a very careful

nucleus the function of generalist... broadly, and to the nutritive protoplasm, the duties of nutrition and adaptation. As, moreover, there is a complete coalescence of the mutually attracted nuclear substances in conception, and the new nucleus formed (the stem-nucleus) is the real starting-point for the development of the fresh organism, the further may be drawn that the male nucleus conveys to the child the qualities of the father, and the female nucleus the features of the mother. We must not forget, however, that the protoplasmic bodies of the copulating cells also fuse together in the act of impregnation; the cell-body of the invading spermatozoon (the trunk and all of the male ciliated cell) is dissolved in the yolk of the female ovum.

— yolk of the female ovum



FIG. 21.—Impregnation of the ovum of a star-fish. (From Hertwig.) Only a small part of one of the sperm is shown. One of the numerous spermatozoa approaches the "impregnation cone" (A), and then penetrates into the protoplasm of the ovum (C).

study of the rise and structure of the stem-cell.

The first question that arises is as to the behaviour of the two different active elements, the nucleus and the protoplasm. It is obvious that the nucleus plays the more important part in this. Hence Hertwig puts his theory of conception in the principle: "Conception consists in the copulation of two cell-nuclei, which come from a male and a female cell." And as the phenomenon of heredity is inseparably connected with the reproductive process, we may further conclude that these two copulating nuclei "convey the characteristics which are transmitted from parents to offspring." In this sense I had in 1866 (in the ninth chapter of the *General Morphology*) ascribed to the reproductive

the nuclei, but it must not be overlooked; though it is known to us, we see clearly at least the formation of the star-like figure (the radial arrangement of the particles in the plasma) in it (Figs. 26-27).

The older theories of impregnation generally went astray in regarding the large ovum as the sole base of the new organism, and only ascribed to the spermatozoon the work of stimulating and originating its development. The stimulus which it gave to the ovum was sometimes thought to be purely chemical, at other times rather physical (on the principle of transferred movement), or again a myotic and transcendental process. This error was partly due to the imperfect knowledge at that time of the facts of impregnation, and partly to the striking

difference in the sizes of the two sexual cells. Most of the earlier observers thought that the spermatozoon did not penetrate into the ovum. And even when this had been demonstrated, the spermatozoon was believed to disappear in the ovum without leaving a trace. However, the splendid research made in the last three decades with the finer technical methods of our time has completely exposed the error of this. It has been shown that the tiny sperm-cell is *not subordinated to, but co-ordinated with*, the large ovum. The nuclei of the two cells, as the vehicles of the hereditary features of the parents, are of equal physiological importance. In some cases we have succeeded in proving that the mass of the active nuclear substance which combines in the copulation of the two

The striking differences of the respective sexual cells in size and shape, which occasioned the erroneous views of earlier scientists, are easily explained on the principle of division of labour. The inert, motionless ovum grows in size according to the quantity of provision it stores up in the form of nutritive yolk for the development of the germ. The active swimming sperm-cell is reduced in size in proportion to its need to seek the ovum and bore its way into its yolk. These differences are very conspicuous in the higher animals, but they are much less in the lower animals. In those protists (unicellular plants and animals) which have the first rudiments of sexual reproduction the two copulating cells are at first quite equal. In these cases the act of impregnation is nothing more than a




FIG. 46.

Impregnation of the ovum of the *a* (ab) moves towards the larger nucleus of the surrounding waste of protoplasm.



FIG. 47.

(From Hertwig.) In Fig. 46 the little sperm-cell (ab) in Fig. 47 they nearly touch and are surrounded by

sexual nuclei is originally the same for both.

These morphological facts are in perfect harmony with the familiar physiological truth that the child inherits from both parents, and that on the average they are equally distributed. I say "on the average," because it is well known that a child may have a greater likeness to the father or to the mother; that goes without saying, as far as the primary sexual characters (the sexual glands) are concerned. But it is also possible that the determination of the latter—the weighty determination whether the child is to be a boy or a girl—depends on a slight qualitative or quantitative difference in the nucleus or the coloured nuclear matter which which comes from both parents in the act of conception.

sudden growth, in which the originally simple cell doubles its volume, and is thus prepared for reproduction (cell-division). Afterwards slight differences are seen in the size of the copulating cells; though the smaller ones will have the same shape as the larger ones. It is only when the difference in size is very pronounced that a notable difference in shape is found: the sprightly sperm-cell changes more in shape and the ovum

square in harmony with this new conception of the equivalence of the two gonads, or the equal physiological importance of the male and female sex-cells and their equal share in the process of heredity, is the important fact established by Hertwig (1875), that in normal impregnation only one single spermatozoon

copulation with one ovum; the membrane which is raised on the surface of the yolk immediately after one sperm-cell has penetrated (Fig. 25 C) prevents any others from entering. All the rivals of the fortunate penetrator are excluded, and die without. But if the ovum passes into a morbid state, if it is made stiff by a lowering of its temperature or stupefied with narcotics (chloroform, morphia, nicotine, etc.), two or more spermatozoa may penetrate into its yolk-body. We then witness *poly spermism*. The more Hertwig chloroformed the ovum, the spermatozoa were able to bore it

conscious body.




FIG. 25.—Stem-cell of a rabbit, magnified 100 times. In the center of the granular protoplasm of the fertilized ovum (a) is seen the little, bright, comma-shaped nucleus. b is the cytoplasm with a nucleus. c are small spermatozoa.

These remarkable facts of impregnation

in psychology, especially as regards the theory of the cell-soul, which I consider to be its chief foundation. The phenomena we

and explained by ascribing a certain degree of psychic activity to the principles. They feel each other's unity, and are drawn to true impulse (probably related to smell); they move towards each other, and do not rest until they fuse together. Physiologists may say that it is only a question of a peculiar physico-chemical phenomenon, and not a psychic action, but in two cannot be separated. Even the psychic functions, in the strict sense of the word, are only complex physical

processes, or "psycho-physical" phenomena, which are determined in all cases exclusively by the chemical composition of their material substratum.

The mosaic view of the matter becomes clear enough when we remember the radical importance of impregnation as regards heredity. It is well known that not only the most delicate bodily structures, but also the subtlest traits of mind, are transmitted from the parents to the children. In this the chromatic matter of the male nucleus is just as important a vehicle as the large karyoplasmic sub-

and the other those of the mother. blending of the two parental nuclei determines the individual psychic character of the child.

But there is another important psychological question—the most important of all—that has been definitely answered by the recent discoveries in connection with conception. This is the question of the immortality of the soul. No fact throws more light on it and refutes it more convincingly than the elementary process of conception that we have described. For this copulation of the two sexual nuclei (Figs. 26-27) indicates the precise moment at which the individual begins to exist. All the bodily and mental features of the newborn child are the sum-total of the hereditary qualities which it has received in reproduction from parents and ancestors. All that man acquires afterwards in life by the exercise of his organs, the influence of his environment, and education—in a word, by adaptation—cannot obliterate that general outline of his being which he inherited from his parents. But this hereditary disposition, the essence of every human soul, is not "eternal," but "temporal"; it comes into being only at the moment when the sperm-nucleus of the father and the nucleus of the maternal ovum meet and fuse together. It is clearly irrational to assume an "eternal life without end" for an individual phenomenon, the commencement of which we can indicate to a moment by direct visual observation.

The great importance of the process of impregnation in answering such questions is quite clear. It is true that studied a opically in all its details in the human life—but withholding its occurrence at every moment—for reasons that are

various enough. However, the two cells which need consideration, the female ovum and the male spermatozoon, proceed in

in all the other mammals, the human foetus or embryo which results from copulation has the same form as with the other animals. Hence, no scientist who is acquainted with the facts doubts that the processes of impregnation are just the same in man as in the other animals.

The stem-cell which is produced, and with which every man begins his career, cannot be distinguished in appearance from those of other mammals, such as the rabbit (Fig. 28). In the case of man, also, this stem-cell differs materially from the original ovum, both in regard to form (morphologically), in regard to material composition (chemically), and in regard

to vital properties (physiologically). It is partly from the father and partly from the mother. Hence it is not surprising that the child who is developed from it inherits from both parents. The vital movements of each of these cells form a sum of mechanical processes which in the last analysis are due to movements of the smallest vital parts, or the molecules, of the living substance. If we agree to call this active substance *plasma*, and its molecules *plastules*, we may say that the individual physiological character of each of these cells is due to its molecular plastidule-movement. Hence, the *plastidule-movement of the ovula is the resultant of the combined plastidule-movements of the female ovum and the male sperm-cell.*¹

CHAPTER VIII.

THE GASTRÆA THEORY

There is a substantial agreement throughout the animal world in the first changes which follow the impregnation of the ovum and the formation of the stem-cell; they begin in all cases with the segmentation of the ovum and the formation of the germinal layers. The only exception is found in the protozoa, the very lowest and simplest forms of animal life; these remain unicellular throughout life. To this group belong the amoebæ, gregarinas, rhizopoda, infusoria, etc. As their whole organism consists of a single cell, they can never form germinal layers, or definite strata of cells. But all the other animals—all the tissue-forming animals, or metazoa, as we call them, in contradistinction to the protozoa—construct real germinal layers by the repeated cleavage of the impregnated ovum. This we find in the lower cœlaria and worms, as well

as in the more highly-developed molluscs, echinoderms, articulates, and vertebrates.

In all these metazoa, or multicellular animals, the chief embryonic processes are substantially alike, although they often seem to a superficial observer to differ considerably. The stem-cell that proceeds from the impregnated ovum always passes by repeated cleavage into a number of simple cells. These cells are all direct descendants of the stem-cell, and are, for reasons we shall see presently, called segmentation-cells. The repeated cleavage of the stem-cell, which gives rise to these segmentation-spheres, has long been known as "segmentation." Sooner or later the segmentation-cells join together to form a round (at first, globular) embryonic sphere (*blastula*); they then form into two very different groups, and arrange themselves

¹ The plasma of the sperm-cell or ovula may, from the anatomical point of view, be regarded as homogeneous and structureless, like that of the amoeba. This is not inconsistent with our hypothetical ascription to the plastidules (or molecules of the plasma) of a complex molecular structure. The complexity of this is the greater in proportion to the complexity of the organism that is developed from it and the length of the chain of its ancestry, or to the subtlety of automatic processes of heredity and adaptation.

in two separate strata—the two primary germinal layers. These enclose a digestive cavity, the primitive gut, with an opening, the primitive mouth. We give the name of the *gastræa* to the important embryonic form that has these primitive organs, and the name of *gastrulation* to the formation of it. This ontogenetic process has a very great significance, and is the real starting-point of the construction of the multicellular animal body.

The fundamental embryonic processes of the cleavage of the ovum and the formation of the germinal layers have been very thoroughly studied in the last thirty years, and their real significance has been appreciated. They present a striking variety in the different groups, and it was no light task to prove their essential identity in the whole animal world. But since I formulated the gastræa theory in 1872, and afterwards (1875) reduced all the various forms of segmentation and gastrulation to one fundamental type, their identity may be said to have been established. We have thus mastered the law of unity which governs the first embryonic processes in all the animals.

Man is like all the other higher animals, especially the apes, in regard to these earliest and most important processes. As the human embryo does not essentially differ, even at a much later stage of development—when we already perceive the cerebral vesicles, the eyes, ears, gill-arches, etc.—from the similar forms of the other higher mammals, we may confidently assume that they agree in the earliest embryonic processes, segmentation and the formation of germinal layers. This has not yet, it is true, been established by observation. We have never yet had occasion to dissect a woman immediately after impregnation and examine the stem-cell or the segmentation-cells in her oviduct. However, as the earliest human embryos we have examined, and the later and more developed forms, agree with those of the rabbit, dog, and other higher mammals, no reasonable man will doubt but that the segmentation and formation of layers are the same in both cases.

But the special form of segmentation and layer formation which we find in the mammal is by no means the original, simple, palingenetic form. It has been much modified and cenogenetically

altered by a very complex adaptation to embryonic conditions. We cannot, therefore, understand it altogether in itself. In order to do this, we have to make a comparative study of segmentation and layer-formation in the animal world; and we have especially to seek the original, palingenetic form from which the modified cenogenetic (see p. 4) form has gradually been developed.

This original unaltered form of segmentation and layer-formation is found to-day in only one case in the vertebrates, to which man belongs—the lowest and oldest member of the stem, the wonderful lancelet or amphioxus (cf. Chapters XVI and XVII). But we find a precisely similar palingenetic form of embryonic development in the case of many of the invertebrate animals, as, for instance, the remarkable ascidia, the pond-animal (*Limnoria*), the arrow-worm (*Sagitta*), and many of the echinoderms and ctenidaria, such as the common star-fish and sea-urchin, many of the medusæ and corals, and the simpler sponges (*Obolus*). We may take as an illustration the palingenetic segmentation and germinal layer-formation in an eight-fold irregular coral, which I discovered in the Red Sea, and described as *Monoceros Dactylus*.

The impregnated ovum of this coral (Fig. 29 A, B) first splits into two equal cells (C). First, the nucleus of the stem-cell and its central body divide into two halves. These recede from and repel each other, and act as centres of attraction on the surrounding protoplasm, in consequence of this, the protoplasm is constricted by a circular furrow, and, in turn, divides into two halves. Each of the two segmentation-cells thus produced splits in the same way into two equal cells. The four segmentation-cells (grand-daughters of the stem-cell) lie in one plane. Now, however, each of them subdivides into two equal halves, the cleavage of the nucleus again preceding that of the surrounding protoplasm. The eight cells which thus arise break into sixteen, these into thirty-two, and then (each being constantly halved) into sixty-four, 128, and so on.* The final result of this

* The number of segmentation-cells thus produced increases geometrically in the original gastrulation, or the proper palingenetic form of cleavage. However, in different animals the number reaches a different height, so that the medusa, and also the Mollusca, may consist sometimes of thirty-two, sometimes of sixty-four, and sometimes of still more cells.




Fig. 9.—Gastrulation of a coel (*Alouatta palliata*). A, B, stem-cell (tertiary) or unimpregnated stem-cell. C (immediately after impregnation) the nucleus is irregular. In Fig. H (a little later) it is more clear. G, symmetrical cells. D, four segmentation cells. E, amphioxys (segmentation). F, blastodermis (blastodermis). H, gastrula, or mesopharynx, external appearance. I, gastrula (longitudinal section).

repeated cleavage is the formation of a globular cluster of similar segmentation-cells, which we call the mulberry-formation or morula. The cells are thickly pressed together like the parts of a mulberry or blackberry, and this gives a lumpy appearance to the surface of the sphere (Fig. E).¹

When the cleavage is thus ended, the mulberry-like mass changes into a hollow globular sphere. Watery fluid or jelly gathers inside the globe; the segmentation-cells are loosened, and all rise to the surface. There they are flattened by mutual pressure, and assume the shape of truncated pyramids, and arrange themselves side by side in one regular layer (Figs. F, G).² This layer of cells is called the germinal membrane (or blastoderm), the homogeneous cells which compose its simple structure are called blastodermic cells; and the whole hollow sphere, the walls of which are made of the preceding, is called the *Blastula* or *Blastosphere*.³

In the case of our coral, and of many other lower forms of animal life, the young embryo begins at once to move independently and swim about in the water. A fine, long, thread-like process, a sort of whip or lash, grows out of each blastodermic cell, and this independently executes vibratory movements, slow at first, but quicker after a time (Fig. F). In this way each blastodermic cell becomes a ciliated cell. The combined force of all these vibrating lashes causes the whole blastula to move about in a rotatory fashion. In many other animals, especially those in which the embryo develops within enclosed membranes, the ciliated cells are only formed at a later stage, or even not formed at all. The blastosphere may grow and expand by the blastodermic cells (at the surface of the sphere) dividing and increasing, and more fluid is secreted in the internal cavity. There are still to-day some organisms that remain throughout life at the structural stage of the blastula—hollow vesicles that swim about by a ciliary movement in the water,

the wall of which is composed of a single layer of cells, such as the volvox, the magosphaera, splanura, etc. We shall speak further of the great phylogenetic significance of this fact in the nineteenth Chapter.

A very important and remarkable process now follows—namely, the curving or invagination of the blastula (Fig. H). The vesicle with a single layer of cells for wall is converted into a cup with a wall of two layers of cells (cf. Figs. G, H, I). A certain spot at the surface of the sphere is flattened, and then bent inward. This depression sinks deeper and deeper, growing at the cost of the internal cavity. The latter decreases as the hollow deepens. At last the internal cavity disappears altogether, the inner side of the blastoderm (that which lines the depression) coming to lie close on the outer side. At the same time, the cells of the two sections assume different sizes and shapes, the inner cells are more round and the outer more oval (Fig. I). In this way the embryo takes the form of a cup or jar-shaped body, with a wall made up of two layers of cells, the inner cavity of which opens to the outside at one end (the spot where the depression was originally formed). We call this very important and interesting embryonic form the "cup-embryo" or "cup-larva" (*Gastrula*, Fig. 29, I longitudinal section, K external view). I have in my *Natural History of Crotalus* given the name of *deplula* to the remarkable intermediate form which appears at the passage of the blastula into the gastrula. In this intermediate stage there are two cavities in the embryo—the original cavity (*blastocoel*) which is disappearing, and the primitive gut-cavity (*progut*) which is forming.

I regard the gastrula as the most important and significant embryonic form in the animal world. In all real animals (that is, excluding the unicellular protists) the segmentation of the ovum produces either a pure, primitive, paleogenetic gastrula (Fig. 29 I, K) or an equally instructive cenogenetic form, which has been developed in time from the first, and can be directly reduced to it. It is certainly a fact of the greatest interest and instructiveness that animals of the most different stamens—vertebrates and tunicates, molluscs and articulate, echinoderms and annelids, ciliata and sponges—proceed from one and the same embryonic form. In illustration I give a few

¹ The segmentation-cells which make up the morula after the close of the paleogenetic cleavage usually tend to be quite regular, and to present no differences as to size, form, and composition. That, however, does not prevent them from differentiating into animal and vegetative cells, even during the cleavage.

² The blastula of the lower animals must not be confused with the very different blastula of the mammal, which is properly called the *gastrula* or *blastogastrula*. This cenogenetic gastrula and the paleogenetic blastula are sometimes very strongly compared under the common name of *Blastula* or *vesicular blastula*.

pure gastrula forms from various groups of animals (Figs. 30-35, explanation given below each).

In view of this extraordinary significance of the gastrula, we must make a very careful study of its original structure. As a rule, the typical gastrula is very small, being invisible to the naked eye, or

half round, or even almost round, and in others lengthened out, or almost cylindrical.

I give the name of primitive gut (*pro-gaster*) and primitive mouth (*prostoma*) to the internal cavity of the gastrula-body and its opening, because this cavity is the first rudiment of the digestive cavity of




FIG. 30 (A).—Gastrula of a very simple primitive-gut animal or protozoan (metaphysan) (*Monard*.)

FIG. 31 (B).—Gastrula of a worm (*Sagitta*). (From *Monard*.)

FIG. 32 (C).—Gastrula of an echinoderm (starfish, *Urechis*), not completely folded in (*Aquila*). (From *Alexander Agassiz*.)

FIG. 33 (D).—Gastrula of an arthropod (primitive crab, *Necoriscus*) (see 36).

FIG. 34 (E).—Gastrula of a mollusk (pond-snail, *Lymnaea*). (From *West Field*.)

FIG. 35 (F).—Gastrula of a vertebrate (lanollet *Amphioxus*). (From *Monard*.) (Front view.)

In each figure *a* is the primitive-gut cavity, *b* primitive mouth, *c* gastrula-body-cavity, *d* ectoderm (epithelium).

at the most only visible as a fine point under very favourable conditions, and measuring generally $\frac{1}{16}$ to $\frac{1}{8}$ of an inch (less frequently $\frac{1}{4}$ inch, or even more) in diameter. In shape it is usually like a roundish drinking-cup. Sometimes it is rather oval, at other times more ellipsoidal or spindle-shaped; in some cases it is

the organism, and the opening originally serving to take food into it. Naturally, the primitive gut and mouth change very considerably afterwards in the various classes of animals. In most of the cnidaria and many of the annelids (worm-like animals) they remain unchanged throughout life. But in most of the

higher animals, and so in the vertebrates, only the larger central part of the later alimentary canal develops from the primitive gut, the later mouth is a fresh development, the primitive mouth disappearing or changing into the anus. We must therefore distinguish carefully between the primitive gut and mouth of the gastrula and the later alimentary canal and mouth of the fully developed vertebrate.¹

The two layers of cells which line the gut-cavity and compose its wall are of extreme importance. These two layers, which are the sole builders of the whole organism, are no other than the two primary germinal layers, or the primitive

all the metazoa or multicellular animals. The skin-layer forms the external skin, the gut-layer forms the internal skin or lining of the body. Between these two germinal layers are afterwards developed the middle germinal layer (*mesoderm*) and the body-cavity (*coelome*) filled with blood or lymph.

The two primary germinal layers were first distinguished by Pander in 1817 in the incubated chick. Twenty years later (1840) Huxley pointed out that in many of the lower zoophytes, especially the medusæ, the whole body consists throughout life of these two primary germinal layers. Soon afterwards (1853) Allman introduced the names which have come




FIG. 31.—Gastrula of a lower sponge (*Sycon*). A reticular view. B longitudinal section through the axis & primitive gut cavity. a primitive mouth-aperture, b outer cell-layer (ectoderm, ectoblast, gut-layer), c ectoderm cell-layer (outer germinal layer, ectoderm, ectoblast, or skin-layer).

germ-layers. I have spoken in the introductory section (Chapter III.) of their radical importance. The outer stratum is the skin-layer, or *ectoderm* (Figs. 30-35c); the inner stratum is the gut-layer, or *endoderm* (c). The former is often also called the ectoblast, or epiblast, and the latter the endoblast, or hypoblast. *From these two primary germinal layers alone is developed the entire organism of*

into general use; he called the outer layer the *ectoderm* ("outer-skin"), and the inner the *endoderm* ("inner-skin"). But in 1867 it was shown, particularly by Kowalevsky, from comparative observation, that even in invertebrates, also, of the most different classes—annelids, molluscs, echinoderms, and articulates—the body is developed out of the same two primary layers. Finally, I discovered them (1872) in the lowest tissue-forming animals, the sponges, and proved in my gastræa theory that these two layers must be regarded as identical throughout the animal world, from the sponges and corals

¹ My distinction (1872) between the primitive gut and mouth and the later permanent stomach (*stomach*) and mouth (*mouth*) has been much criticised, but it is as much justified as the distinction between the outer embryo and the permanent kidney. Prof. or E. Ray-Lankester attributed these views after

"insects and vertebrates, including This fundamental "homology

[identity] of the primary germinal layers and the primitive gut¹ has been confirmed during the last thirty years by the careful research of many able observers, and is now pretty generally admitted for the whole of the metazoa.

As a rule, the cells which compose the two primary germinal layers show appreciable differences in their early development stage. Generally (if not always) the cells of the skin-layer or ectoderm (Figs. 36c, 37e) are the smaller, more numerous,

layer, or entoderm (e), are larger, less numerous, and darker. The protoplasm of the ectodermic (outer) cells is clear

matter of the entodermic (inner) cells, the latter are, as a rule, much richer in yellow granules (albumen and fatty particles) than the former. Also the cells of the gut-layer have, as a rule, a stronger affinity for colouring matter, and take on a tinge in a solution of carmine, aniline, etc., more quickly and appreciably than the cells of the skin-layer. The nuclei of the entoderm-cells are usually roundish, while those of the ectoderm-cells are oval.

When the doubling-process is complete, very striking histological differences between the cells of the two layers are found (Fig. 37). The tiny, light ectoderm-cells (e) are sharply distinguished from the larger and darker entoderm-cells (e). Frequently this differentiation of the cell-forms sets in at a very early stage, during

very appreciable in the blastula.

We have, up to the present, considered that form of segmental gastrulation which, for many and weighty reasons, we may regard as the original, primordial, or palingenetic form. We

segment

... resemblance to each other at first (and often until the formation of the blastoderm). We give the name of the "bell-gastrula," or *stolidogastrula*, to the gastrula that succeeds it. In just the


... (Fig. 29), we find it in lowest zoophytes (the gastrophytes, Fig. 70), and the simplest sponges (lynthus, Fig. 36); also in many of the medusæ and hydrapolytes, lower types of worms of various classes (brachiopod, arrow-worm, Fig. 31), tunicates (ascidii), many of the echinoderms (Fig. 32), lower articulate (Fig. 33), and molluscs (Fig. 34), and,

finally, in a slightly modified form, in the lowest vertebrate (the amphioxus, Fig. 35).

The gastrulation of the amphioxus is especially interesting because this lowest and oldest of all the vertebrates is of the highest significance in connection

... of the vertebrate stem, a ... of man (con... Chapters XVI. and XVII.). Just as the comparative anatomist traces the most elaborate features in the structures of the vertebrates to divergent development from this simple primitive ... comparative ... secondary forms of ve

formation of the germinal layers in the



amphioxus. Although this form is distinguished from the congeneric modifications

hole be regarded as palingenetic, it is nevertheless different in some feature from the quite primitive gastrulation such as we have, for instance, in the *Monostoma* (Fig. 39) and the *Segella*. Haeckel rightly observes that the segmentation of the ovum in the amphioxus is not strictly equal, but almost equal, and approaches the unequal. The difference in size between the two groups of cells continues to be very noticeable in the further course of the segmentation; the smaller animal cells of the upper hemisphere divide more quickly than the larger vegetal cells of the

(Fig. 38 A, B). Her blastoderm, which forms the single-layer wall of the globular Naetula at the end of the cleavage-process, does not consist of

homogeneous cells of equal size, as in the *Sagitta* and the *Monaxonus*; the cells of the upper half of the blastoderm (the mother-cells of the ectoderm) are more numerous and smaller, and the cells of the lower half (the mother-cells of the endoderm) less numerous and larger. Moreover, the segmentation-cavity of the blastula (Fig. 38 C, A) is not quite globular, but forms a flattened spheroid with unequal poles of its vertical axis. While the blastula is being folded into a cup at the vegetal pole of its axis, the difference in the size of the blastodermic cells increases (Fig. 38 D, E); it is most conspicuous when the invagination is complete and the segmentation cavity has

the blastoderm (or the increase of its cells) being brisker on one side than on the other; the side that grows more quickly, and so is more curved (Fig. 39 v), will be the anterior or belly-side, the opposite, flatter side will form the back (d). The primitive mouth, which at first, in the typical archigastrea, lay at the vegetal pole of the main axis, is forced away to the dorsal side; and whereas its two lips lay at first in a plane at right angles to the chief axis, they are now so far thrust aside that their plane cuts the axis at a sharp angle. The dorsal lip is therefore the upper and more forward, the ventral lip the lower and hinder. In the latter, at the ventral passage of the endoderm into

the ectoderm, there lie side by side a pair of very large cells, one to the right and one to the left (Fig. 39 p); these are the important polar cells of the primitive mouth, or "the primitive cells of the mesoderm." In consequence of these considerable variations arising in the course of the gastrulation, the primitive uniaxial form of the archigastrea in the amphioxus has already become triaxial, and thus the two-sidedness, or bilateral symmetry, of the vertebrate body has already been determined.




FIG. 38.—Gastrulation of the amphioxus, from *Helfferich* (vertical) section through the axis of the ovum. A, B, C three stages on the formation of the blastula; D, E, F closing of the blastula; F complete gastrula. A segmentation-cavity; g primitive gut-cavity.

disappeared (Fig. 38 F). The larger vegetal cells of the endoderm are richer in granules, and so darker than the smaller and lighter animal cells of the ectoderm.

But the unequal gastrulation of the amphioxus diverges from the typical equal cleavage of the *Sagitta*, the *Monaxonus* (Fig. 29), and the *Obolus* (Fig. 36), in another important particular. The pure archigastrea of the latter forms is uniaxial, and it is round in its whole length in transverse section. The vegetal pole of the vertical axis is just in the centre of the primitive mouth. This is not the case in the gastrula of the amphioxus. During the folding of the blastula the ideal axis is already bent on one side, the growth of

This has been transmitted from amphioxus to all the other modified gastrula-forms of the vertebrate stem.

Apart from this bilateral structure, the gastrula of the amphioxus resembles the typical archigastrea of the lower animals (Fig. 30-36) in developing the two primary germinal layers from a single layer of cells. This is clearly the oldest and original form of the metazoic embryo. Although the animals I have mentioned belong to the most diverse classes, they nevertheless agree with each other, and many more animal forms, in having returned to the present day, by a conservative heredity, this palaeogenetic form of gastrulation which they have from their

earliest common ancestors. But this is not the case with the great majority of the animals. With these the original embryonic process has been gradually less altered in the course of

segmentation of the ovum and the subsequent gastrulation have in this way been considerably changed. In fact, these various

the course of it was not rightly understood animals, and the gastrula was unrecognized. It was not until I had made an extensive comparative study, lasting a considerable time (in the years 1866-75), in animals of the most diverse classes, that I succeeded in showing the same common typical process in these apparently very different forms of gastrulation, and tracing them all to one original form. I regard all those that diverge from the primary palingeopetic gastrulation as secondary, modified, and cenogenetic. The more or less divergent forms of gastrula that is produced may be called a secondary, modified gastrula, or a metagastrula. The reader will find a schema of these different kinds of segmentation and gastrulation at the close of this chapter.

By far the most important process that determines the various cenogenetic forms of gastrulation is the change in the nutrition of the ovum and the accumulation in it of nutritive yolk. By this we understand various chemical substances (chiefly granules of albumin and fat-particles) which serve exclusively as reserve-matter or food for the embryo. As the metazoic embryo in its earlier stages of development

obtain its food and so build up the frame, the necessary material has to be stored up in the ovum. Hence we distinguish in the ova two chief elements—the active formative yolk (protoplasm) and the passive food-yolk (deutoplasm, wrongly spoken of as “the yolk”). In the little palingeopetic ova, the segmentation of which we have already considered, the yolk-granules are so small and so regularly distributed in the protoplasm of the ovum that the even and repeated cleavage is not affected by them. But in the great majority of the animal ova the food-yolk is more or less considerable, and is stored in a certain part of the ovum, so that even in the unfertilized ovum the “granary”

can clearly be distinguished from the formative plasma. As a rule, the formative-yolk (with the germinal vesicle) then usually gathers at one pole and the food-yolk at the other. The first is the animal, and the second the vegetal, pole of the

In these “telolecithal” ova, or ova with the yolk at one end (for instance, in the cyclostoma and amphibia), the gastrula-

way that in the cleavage of the impregnated ovum the animal (usually the upper) half splits up more quickly than the vegetal (lower). The contractions of the active protoplasm, which effect this continual cleavage of the cells, meet a greater resistance in the lower vegetal half from the passive deutoplasm than in the upper animal half. Hence we find in the latter




Fig. 20.—Gastrula of the amphibia, seen from left side (deutoplasmic mass active). (From Haeckel's "Gastrulae")

more but smaller, and in the former fewer but larger, cells. The animal cells pro-

the internal, germinal layer.

Although this unequal segmentation of the cyclostoma, graptolite, and amphibia seems at first sight to differ from the original equal segmentation (for instance, in the marisoma, Fig. 25), they both have this in common, that the cleavage process throughout affects the whole cell, hence Rernak called it *total* segmentation, and the ova in question *holoblastic*, or “whole-cleaving.” It is otherwise with the second chief group of ova, which he distinguished from these as *meroblastic*, or “partially-cleaving.” To this class belong the familiar large eggs of birds and reptiles, and of most fishes. The inert mass of the passive food-yolk is so

large in these cases that the protoplasmic contractions of the active yolk cannot effect any further cleavage. In consequence, there is only a partial segmentation. While the protoplasm in the animal section of the ovum continues briskly to divide, multiplying the nuclei, the deutoplasm in the vegetal section remains more or less undivided; it is merely consumed as food by the forming cells. The larger the accumulation of food, the more restricted is the process of segmentation. It may, however, continue for some time (even after the gastrulation is more or less complete) in the sense that the vegetal cell-nuclei distributed in the deutoplasm slowly increase by cleavage; as each of them is surrounded by a small quantity of protoplasm, it may afterwards appropriate a portion of the food-yolk, and thus form a real "yolk-cell" (*merocyte*). When this vegetal cell-formation continues for a long time, after the two primary germinal layers have been formed, it takes the name of the "after-segmentation."

The meroblastic ova are only found in the larger and more highly developed animals, and only in those whose embryo needs a longer time and richer nourishment within the fetal membranes. According as the food-yolk accumulates at the centre or at the side of the ovum, we distinguish two groups of dividing ova, periblastic and discoblastic. In the periblastic the food-yolk is in the centre, enclosed inside the ovum (hence they are also called "centrolecithal" ova); the formative yolk surrounds the food-yolk, and so suffers itself a superficial cleavage. This is found among the articulate (crabs, spiders, insects, etc.). In the discoblastic ova the food-yolk gathers at one side, at the vegetal or lower pole of the vertical axis, while the nucleus of the ovum and the great bulk of the formative yolk lie at the upper or animal pole (hence these ova are also called "teleolecithal"). In these cases the cleavage of the ovum begins at the upper pole, and leads to the formation of a dorsal divided embryo. This is the case with all meroblastic vertebrates, most fishes, the reptiles and birds, and the viviparous mammals (the monotremes).

The gastrulation of the discoblastic ova, which chiefly concerns us, offers serious difficulties to microscopic investigation and philosophic consideration. These, however, have been mastered by

the comparative embryological research which has been conducted by a number of distinguished observers during the last few decades—especially the brothers Hertwig, Rabi, Kupffer, Selenka, Ruckert, Goette, Rauber, etc. These thorough and careful studies, aided by the most perfect modern improvements in technical method (in tinting and dissection), have given a very welcome support to the views which I put forward in my work, *On the Gastrula and the Segmentation of the Animal Ovum* [not translated], in 1875. As it is very important to understand these views and their phylogenetic foundation clearly, not only as regards evolution in general, but particularly in connection with the genesis of man, I will give here a brief statement of them as far as they concern the vertebrate-stem.

1. All the vertebrates, including man, are phylogenetically (or genealogically) related—that is, are members of one single natural stem.

2. Consequently, the embryonic features in their individual development must also have a genetic connection.

3. As the gastrulation of the amphioxus shows the original palæogenetic form in its simplest features, that of the other vertebrates must have been derived from it.

4. The cenogenetic modifications of the latter are more approx. while the more food-yolk is stored up in the ovum.

5. Although the mass of the food-yolk may be very large in the ova of the discoblastic vertebrates, nevertheless in every case a blastula is developed from the morula, as in the holoblastic ova.

6. Also, in every case, the gastrula develops from the blastula by curving or invagination.

7. The cavity which is produced in the folds by this curving is, in each case, the primitive gut (pro-gut), and its opening the primitive mouth (pro-stoma).


8. The food-yolk, whether large or small, is always stored in the ventral wall of the primitive gut; the cells (called "merocytes") which may be formed in it subsequently (by "after-segmentation") also belong to the inner germinal layer, like the cells which immediately enclose the primitive gut-cavity.

9. The primitive mouth, which at first lies below at the lower pole of the vertical axis, is formed, by the growth of the yolk, backwards and then upwards,

towards the dorsal side of the embryo; the vertical axis of the primitive gut is thus gradually converted into horizontal.

20. The primitive mouth is closed sooner or later in all the vertebrates, and does not evolve into the permanent mouth-aperture; it rather corresponds to the "propræstoma," or region of the anus. From this important point the formation of the middle germinal layer proceeds, between the two primary layers.

The wide comparative studies of the scientists I have named have further shown that in the case of the discoblastic higher vertebrates (the three classes of animals) the primitive mouth of the embryonic disc, which was long looked for in vain, is found always, and is nothing else than the familiar "primitive groove." Of this we shall see more as we proceed. Meantime we realize that gastrulation may be reduced to one and the same process in all the vertebrates. Moreover, the various forms it takes in the invertebrates can always be reduced to one of the four types of segmentation described above. In relation to the distinction between total and partial segmentation, the grouping of the various forms is as follows:—



The lowest metazoans we know—namely, the lower zoophytes (sponges, simple polyps, etc.)—remain throughout life at a stage of development which differs little from the gastrula, their whole body consists of two layers of cells. This is a fact of extreme importance. We see that man, and also other vertebrates, pass quickly through a stage of development in which they consist of two layers, just as these low zoophytes do throughout life. If we apply our biogenetic law to this matter, we at once reach this important conclusion: "Man and all the other animals which pass through the two-layer stage,

or gastrula-form, in the course of their embryonic development, must descend from a primitive simple stem-form, the whole body of which consisted throughout life (as is the case with the lower zoophytes to-day) merely of two cell-strata or germinal layers." We will call this primitive stem-form, with which we shall deal more fully later on, the *gastræa*—that is to say, "primitive-gut animal."

According to this gastræa-theory there was originally in all the multicellular animals *one organ* with the same structure and function. This was the primitive gut; and the two primary germinal layers which form its wall must also be regarded as identical in all. This important homology or identity of the primary germinal layers is proved, on the one hand, from the fact that the gastrula was originally formed in the same way in all cases—namely, by the curving of the blastula; and, on the other hand, by the fact that in every case the same fundamental organs arise from the germinal layers. The outer or animal layer, or ectoderm, always forms the chief organs of animal life—the skin, nervous system, sense-organs, etc.; the inner or vegetal layer, or endoderm, gives rise to the chief organs of vegetative life—the organs of nourishment, digestion, blood-formation, etc.

In the lower zoophytes, whose body remains at the two-layer stage throughout life, the gastræads, the simplest sponges (*Olynthus*), and polyps (*Hydra*), these two groups of functions, animal and vegetative, are strictly divided between the two simple primary layers. Throughout life the outer or animal layer acts simply as a covering for the body, and accomplishes its movement and sensation. The inner or vegetative layer of cells acts throughout life as a gut-lining, or nutritive layer of entere cells, and often also yields the reproductive cells.

The best known of these "gastræads," or "gastrula-like animals," is the common fresh-water polyp (*Hydra*). This simplest of all the animals, as it is true, a crown of tentacles round its mouth. Also its outer germinal layer has certain special modifications. But these are secondary additions, and the inner germinal layer is a simple stratum of cells. On the whole, the hydra has preserved to our day by heredity the simple structure of our primitive ancestor, the *gastræa* (cf. Chapter XIX.)

In all other animals, particularly the vertebrates, the gastrula is merely a brief transitional stage. Here the two-layer stage of the embryonic development is quickly succeeded by a three-layer, then four-layer, stage. With

appearance of the four superimposed germinal layers we reach again a firm and steady standing-ground, from which we may follow the further, and much more difficult and complicated, course of embryonic development.

SUMMARY OF THE CHIEF DIFFERENCES IN THE OVUM-SEGMENTATION AND GASTRULATION OF ANIMALS.

The animal stems are indicated by the letters *a-g*: *a* Zoophyta, *z* Annelida, *c* Mollusca, *e* Echinodermata, *s* Arthropoda, *f* Terebrata, *g* Vertebrata.

Gastrula without	II. Unequal Segmentation. Acyrtularian or a.	
Partial Segmentation. Acyrtularian	III. Discoid Segmentation. Discoblastic or a. Discoid gastrula.	<div> <i>c</i> Cephalopods or cuttle-fish. <i>z</i> <i>Uca</i> and a few other crustaceans. </div> <div> <i>f</i> Primitive fishes, reptiles, birds, &c. </div>
Gastrula with separate food-yolk. Mecyrtularian	IV. Superficial Segmentation. Periblastic ova. Ectoblastic or a.	<div> </div>

THE GASTRULATION OF THE VERTEBRATE¹

THE remarkable processes of gastrulation, ovum-segmentation, and formation of germinal layers present a most conspicuous variety. There is to-day only the lowest of the vertebrates, the amphioxus, that exhibits the original form of these processes, or the palæogenetic gastrulation which we have considered in the preceding chapter, and which culminates in the formation of the archigastrula (Fig. 38). In all other extant vertebrates these fundamental processes have been more or less modified by adaptation to the conditions of embryonic development (especially by changes in the food-yolk), they exhibit various cenogenetic types of the formation of germinal layers. However, the different classes vary considerably from each other. In order to grasp the unity that underlies the many

phenomena and

the unity of the
"phylogenetic

unity," which I developed in my *General Morphology* in 1866, is now generally accepted. All impartial zoologists agree to-day that all the vertebrates, from the fish to the ape and

in, hence
"the primitive vertebrate." Hence the embryonic processes, by which each individual vertebrate is developed, must also be capable of being reduced to one common type of embryonic development; and this primitive type is most certainly exhibited to-day by the amphioxus.

It must, therefore, be our next task to make a comparative study of the various forms of vertebrate gastrulation, and trace them backwards to that of the lancelet. Broadly speaking, they fall first into two groups: the older cyclostoma, the earliest fishes, most of the amphibia, and the viviparous mammals, have *Aleo-*

lastic ova—that is to say, ova with total, unequal segmentation; while the younger cyclostoma, most of the fishes, the cephalopoda, reptiles, birds, and monotremes, have *meroblastic ova*, or ova with partial discoid segmentation. A closer study of them shows, however, that these two groups do not present a natural unity, and that the historical relations between their several divisions are very complicated. In order to understand them properly, we must first consider the various modifications of gastrulation in these classes. We may begin with that of the amphibia.

The most suitable and most available objects of study in this class are the eggs of our indigenous amphibia, the tailless frog and toad, and the tailed salamander. In spring they are to be found in clusters in every pond, and careful examination of

real features of the

In order to understand the whole process rightly and follow the formation of the germinal layers and the gastru-

if the frog and salamander must be carefully hardened; then the thinnest possible sections must be made of the hardened ova with the microscope, and the tinted sections must be very closely compared under a powerful microscope.

The ova of the frog or toad are globular in shape, about the twelfth of an inch in diameter, and are clustered in jelly-like masses, which are lumped together in the case of the frog, but form long strings in the case of the toad. When we examine the opaque, grey, brown, or blackish ova closely, we find that the upper half is darker than the lower. The middle of the upper half is in many species black, while the middle of the lower half is white. In this way we get a definite axis of the ovum with two poles. To give a clear

¹ Cf. Baillard's *Manual of Comparative Embryology*, vol. ii.; Theodore Morgan's *The Development of the Frog's Egg*.

² The colouring of the eggs of the amphibia is caused by the accumulation of dark-coloured matter at the animal pole of the ovum. In consequence of this the animal cells of the ectoderm are darker than the vegetal cells of the ectoderm. We find the reverse of this in the case of most animals, the protoplasm of the ectoderm cells being usually darker and more greenish.

idea of the segmentation of this ovum, it is best to compare it with a globe, on the surface of which are marked the various parallels of longitude and latitude. The superficial dividing lines between the different cells, which come from the repeated segmentation of the ovum, look like deep furrows on the surface, and hence the whole process has been given the name of furcation. In reality, however, this "furcation," which was formerly regarded as a very mysterious process, is

in this position throughout the course of the segmentation, and its cells multiply much more briskly. Hence the cells of the lower hemisphere are found to be larger and less numerous. The cleavage of the stem-cell (Fig. 40 *A*) begins with the formation of a complete furrow, which starts from the north pole and reaches to the south (*B*). An hour later a second furrow arises in the same way, and this cuts the first at a right angle (Fig. 40 *C*). The ovum is thus divided into four equal




FIG. 40.—The cleavage of the frog's ovum (magnified ten times). *A* stem-cell. *B* the first two eggs taken out. *C* four cells. *D* eight cells (4 animal and 4 vegetative). *E* sixteen cells (8 animal and 8 vegetative). *F* sixteen cells (8 animal and 8 vegetative). *G* twenty-four cells (16 animal and 8 vegetative). *H* thirty-two cells (16 animal and 16 vegetative). *I* forty-eight cells. *J* sixty-four cells. *K* eighty-four cells. *L* one hundred and twenty-eight cells. *M* one thousand and twenty-four cells (400 animal and 300 vegetative).

nothing but the familiar, repeated cell-segmentation. Hence also the segmentation-cells which result from it are real cells.

The unequal segmentation which we observe in the ovum of the amphibia has the special feature of beginning at the upper and darker pole (the north pole of the terrestrial globe in our illustration), and slowly advancing towards the lower and brighter pole (the south pole). Also the upper and darker hemisphere remains

parts. Each of these four "segmented cells" has an upper and darker and lower, brighter half. A few hours later a third furrow appears, vertically to first two (Fig. 40 *D*). The globular we now consists of eight cells, four small ones above (northern) and four large ones below (southern). Next, each of the four upper divides by a cleavage beginning from the pole, so that we now have eight above and four below (Fig. 40 *E*). Later, the

four new longitudinal divisions extend gradually to the lower cells, and the number rises from twelve to sixteen (*P*). Then a second circular furrow appears, parallel to the first, and nearer to the north pole, so that we may compare it to the north polar circle. In this way we get twenty-four segmentation-cells—sixteen upper, smaller, and darker ones, and eight smaller and lighter ones below.

In succession, forty, forty-eight, fifty-six, and at last sixty-four cells (*I, K*). In the meantime, the two hemispheres differ more and more from each other. Whereas the sluggish lower hemisphere long remains at thirty-two cells, the lively northern hemisphere briskly sub-divides twice, producing first sixty-four and then 128 cells (*L, M*). Thus we reach a stage in which we count on the surface




FIG. 41



FIG. 42




FIG. 44

ovum of the toad, in four success

(*D* dorsal half of the embryo, *V* ventral half, *F* yolk-stopper (white round field at the lower pole), *Z* yolk-cells of the entoderm (blackish "granular embryo"), *A* primitive gut cavity (protoplast or blastoderm blastoderm cavity). The primitive mouth (proctoderm) is closed by the yolk-stopper. *P* a partition between the primitive gut cavity (*N*) and the mesenteric cavity (*F*). *K* *K'* sections of the large circular lip-border of the primitive mouth (the blastoderm). The line of dots between *K* and *K'* ends at the outer connection of the yolk-stopper (*P*) with the central mass of the yolk-cells (*Z*). In Fig. 44 the dot has turned up, so that the back of the embryo is uppermost and the ventral side down. (I from Schröder.)

(*G*). Soon, however, the latter also sub-divide into sixteen, a third or "meridian of latitude" appearing, this time in the southern hemisphere: this makes thirty-two cells altogether (*H*). Then eight new longitudinal lines are formed at the north pole, and these proceed to divide, first the darker cells above and afterwards the lighter southern cells, and finally reach the south pole. In this way we get

of the ovum 128 small cells in the upper half and thirty-two large ones in the lower half, or 160 altogether. The dissimilarity of the two halves increases: while the northern breaks up into a great number of small cells, the southern consists of a much smaller number of larger cells. Finally, the dark cells of the upper half grow almost over the surface of the ovum, leaving only a small circular spot

at the south pole, where the large cells of the lower half are visible. This white region at the south pole corresponds, as we shall see afterwards, to the primitive mouth of the gastrula. The



FIG. 41.—Gastrula of the water-salamander (*Triton*). *P*, segmentation-cavity, the yolk-cells, on border-line. (From Hertwig.)

..... of the inner and larger and clearer cells (including the white polar region) belongs to the ontoderm layer. The outer envelope of dark smaller cells forms the ectoderm or skin-layer.

In the meantime, a large cavity, full of fluid, has been formed within the globular body—the segmentation-cavity or embryonic cavity (*blastocoel*, Figs. 41-44 *P*). It extends considerably as the cleavage proceeds, and afterwards assumes an almost semi-circular form (Fig. 42 *P*). The frog-embryo now represents a modified embryonic vesicle or *blastula*, with hollow animal half and solid vegetal half.

Now a second, narrower but longer, cavity arises by a process of folding at the lower pole, and by the falling away from each other of the white ontoderm-cells (Figs. 41-44 *N*). This is the primitive gut-cavity or the gastric cavity of the gastrula, *progut* or *archenteron*. It was first observed in the ovum of the amphibia by Rusconi, and so called the Rusconian cavity. The reason of its peculiar narrowness here is that it is, for the most part, full of yolk-cells of the ectoderm. These also stop up the whole of the wide opening of the primitive mouth, and form what is known as the "yolk-stopper," which is seen freely at the white round spot at the south pole (*P*). Around it the ectoderm is much thicker, and forms the border of the primitive mouth, the

most important part of the embryo (Fig. 44 *K*, *P*). Soon the primitive gut-cavity stretches further and further at the expense of the segmentation-cavity (*P*), until at last the latter disappears altogether. The two cavities are only separated by a thin partition (Fig. 43 *A*). With the formation of the primitive gut our frog-embryo has reached the *gastrula* stage, though it is clear that this cenogenetic amphibian gastrula is very different from the real palingogenetic gastrula we have considered (Figs. 30-36).

In the growth of this hooded gastrula we cannot sharply mark off the various stages which we distinguish successively in the bell-gastrula, *amphibula* and gastrula. Nevertheless, it is not difficult to reduce the whole cenogenetic or disturbed development of this amphigastrula to the true palingogenetic formation of the archigastrula of the amphioxus.

This reduction becomes easier if, after considering the gastrulation of the tailless amphibia (frogs and toads), we glance for a moment at that of the tailed amphibia, the salamanders. In some of the latter, that have only recently been carefully studied, and that are phylogenetically older, the process is much simpler and clearer than in the case with the former and longer known. Our common water-salamander (*Triton lacustris*) is a particularly good subject for observation. Its nutritive yolk is much smaller and its formative yolk less obscured with black pigment-cells than in the case of the frog; and its gastrulation has better retained the original palingogenetic character. It was first described by Scott and Osborn (1879), and Oscar Hertwig especially made



FIG. 42.—Embryo vesicle of *Triton* (*blastula*). *P*, *A*, same, with the transverse fold of the primitive mouth (*P*). (From Hertwig.)

..... *seriatim* development. Its globular blastula (Fig.) is of loosely-aggregated, yolk-

filled entodermic cells or yolk-cells (*dy*) in the lower vegetal half; the upper, animal half encloses the hemispherical segmentation-cavity (*sh*), the curved roof of which is formed of two or three




FIG. 41.—Sagittal section of a herring-embryo (*depictus*) at the commencement of gastrulation. *sh*, outer germinal layer; *md*, inner germinal layer; *sh*, segmentation-cavity; *dy*, primitive gut; *st*, primitive mouth; *dl*, end of dorsal; and ventral lip of the mouth, *dy*, yolk-cells. (From Hertwig.)

have the border zone? *ca*). The folding which leads to the formation of the gastrula takes place at a spot in this border zone, the primitive mouth (Fig. 46 a).

Unequal segmentation takes place in some of the cyclostoma and in the oldest fishes in just the same way as in most of the amphibia. Among the cyclostoma ("round-mouthed") the familiar lampreys

half-way between the acrania (I

were formerly associated, and I then a special class of vertebrate ovum-segmentation in our common river-lamprey (*Petromyzon fluviatilis*) was described by Max Schultz in 1856, and afterwards by Scott (1882) and Goette (1890).

Unequal total segmentation follows the same lines in the oldest fishes, the selachii and ganoids, which are directly desc. from the cyclostoma. The primitive fish (Selachii), which are the ancestral group of the true fishes,

generally considered, until a short time ago, to be discoblastic. It was not until the beginning of the twentieth century that Bashford Down made the important discovery in Japan that one of the oldest living fishes of the shark type (*Coelacanth japonicus*) has the same total unequal segmentation as the amphiblastic plated fishes (*ganoides*).¹ This is particularly interesting in connection with our subject, because the few remaining survivors of this division, which was so numerous in paleozoic times, exhibit three different types of gastrulation. The oldest and most conservative forms of the modern ganoids are the scaly sturgeons (*Sturiones*), plated fishes of great evolutionary importance, the eggs of which are eaten as caviare; their cleavage is not essentially different from that of the lampreys and the amphibia. On the other hand, the most modern of the plated fishes, the beautifully scaled bony pike of the North American rivers (*Lepidosteus*), approaches the osseous fishes, and is discoblastic like them. A third genus (*Araia*) is midway between the sturgeons and the latter.

The group of the lung-fishes (*Dipnoeae* or *Dipnoi*) is closely connected with the older ganoids. In respect of their whole




FIG. 42.—Sagittal section of the gastrula of the water-scorpion (*Tribolium*). (From Hertwig.)
Labels: *sh*, segmentation-cavity; *dy*, yolk-cells; *md*, mesoderm; *dl*, dorsal lip of the mouth; *st*, stomach; *mb*, mouth border.

organization they are midway between the gill-breathing fishes and the lung-breathing amphibia, they share with the former the shape of the body and fins, with the latter the form of the heart.

¹ Bashford Down, *Embryonic Cleavage in the Egg of a Shark, Coelacanth japonicus* (Macleay). *Annals of the Entomological Society of America*, vol. 17, 1924.

and lungs. Of the older dipnoi (*Psalidipneuste*) we have now only one specimen, the remarkable *Ceratodus* of East Australia; its amphiblastic gastrulation has been recently explained by Richard Semon (cf. Chapter XXI.). That of the two

and batrachia, belong to the old, conservative groups of our stem. Their unequal cross-segmentation and gastrulation have many peculiarities in detail, but can always be reduced with comparative ease to the original cleavage and gastrulation




FIG. 20.—Gastrulation of the lamprey (*Petromyzon Australis*). A blastula, with wide embryonic cavity (blastocoel) & isomeric segmentation. B cepula, with advanced invagination from the primitive mouth (g). C gastrula, with complete primitive gut. The embryonic cavity has almost disappeared.

America, is not materially different. (Cf. Fig. 51.)

All these amphiblastic vertebrates, *Petromyzon* and *Ceratodus*, *Accipenser* and *Ceratodus*, and also the salamanders

of the lowest vertebrate, the amphioxus; and this is little removed, as we have seen, from the very simple archegastrula of the *Sagittæ* and *Monoxenus* (see Figs. 29-35). All these and many other classes of animals generally agree in the circumstances that in segmentation their

ovum divides into a large number of cells by repeated cleavage. All such ova have been called, after Kernak, "whole-cleaving" (*Aleoblasts*), because their division into cells is complete or total.

In a great many other classes of animals this is not the case, as we find (in the vertebrate stem) among the birds, reptiles, and most of the fishes; among the insects and most of the spiders and

of the ovum; this alone divides in segmentation, and produces the numerous cells which make up the embryo. On the other hand, the nutritive yolk is merely a passive part of the contents of the ovum, a subordinate element which contains nutritive material (albumin, fat, etc.) and so represents in a sense the provision-store of the developing embryo. The latter takes a quantity of food out of this




FIG. 3.—Gastrulation of *ceratodus* (from Semov). *A* and *C* stage with four cells, *B* and *D* with sixteen cells. *A* and *B* are seen from above, *C* and *D* sideways. *E* stage with thirty-five cells, *F* blastula. *G* gastrula in longitudinal section. *fb* blastoderm cavity. *zh* primitive gut or gastric cavity.

crabs (of the arthropods); and the cephalopods (of the molluscs). In all these animals the mature ovum, and the stem-cell that arises from it in fertilisation, consist of two different and separate parts, which we have called formative yolk and nutritive yolk. The formative yolk alone consists of living protoplasm, and is the active, evolutionary, and nucleated part

store, and finally consumes it all. Hence the nutritive yolk is of great indirect importance in embryonic development, though it has no direct share in it. It either does not divide at all, or only later on, and does not generally consist of cells. It is sometimes large and sometimes small, but generally many times larger than the formative yolk; and hence it is

that it was formerly thought the more important of the two. As the respective significance of these two parts of the ovum is often wrongly described, it must be

minimal
secondary

real appendages. All ova that have independent nutritive yolk are called,

blast). Their segmentation is complete or partial.

There are many difficulties of understanding this part

ing these difficulties, and reducing the ontogenetic form of gastrula to the original palingnetic type.




FIG. 52.—Ovum of a deep-sea hony fish. *a*, protoplasm of the ovum; *n*, nucleus of ovum; *y*, nutritive yolk; *e*, outer membrane of the ovum, or vitelline membrane.

comparatively easy in the small mem-

found them joined together in lumps of jelly, floating on the surface of the sea, and, as the little ovula were completely

he nucleus (*n*), this is the formative yolk of the stem-cell, or the germinal disk. The nutritive yolk (*y*) and the

elk not dividing at all!

formative yolk (*y*) proceeds quite inde-

like a watch-glass, with thick borders at the upper and lower poles, and the nutritive yolk like the

and the border,

tion, or a kind of turning-up of the edge of the blastoderm. In this process the segmentation-cavity disappears.

The space underneath the entoderm corresponds to the primitive gut-cavity, and is filled with the decreasing food-yolk (*y*). Thus the formation of the gastrula of our fish is complete. In contrast to the two chief forms of gastrula we considered previously, we give the name of discoid gastrula (*discogastrula*, Fig. 54) to this third principal type.

Very similar to the discoid gastrulation of the bony fishes is that of the hags or myxini, the remarkable cyclostomes that live parasitically in the body-cavity of fishes, and are distinguished by several notable peculiarities from their nearest

membrane (*protoplasm*, Fig. 51 *c*) and a large, quite clear, and transparent globule of albumin (*d*). At both poles of

le (which is turned down) (Fig. 51 *a*) there is a bi-convex lens composed of protoplasm, and this encloses

chief for considered are determined by the large food-yolk. This takes no direct part in the

city of the opening. If we imagine the original gastrula (Figs. 30-36) trying to swallow a

ball of food which is much bigger than itself, it would spread out round it in discoid shape in the attempt, just as we find to be the case here (Fig. 54). Hence we may derive the discoid gastrula from the original ball-gastrula, through the intermediate stage of the hooded gastrula. It has arisen through the accumulation of a store of food-stuff at the vegetal pole, a "nutritive yolk" being thus formed in contrast to the "formative yolk." Nevertheless, the gastrula is formed here, as in the previous cases, by the folding or invagination of the blastula. We can, therefore, reduce this cenogenetic form of the discoid segmentation to the paligenetic form of the primitive cleavage.

This reduction is tolerably easy and confident in the case of the small ovum of our deep-sea bony fish, but it becomes

embryonic development and consumed by the embryo. The latter develops solely from the living formative yolk of the stem-cell. This is equally true of the ova of our small bony fishes and of the colossal ova of the primitive fishes, reptiles, and birds.

The gastrulation of the primitive fishes or selachii (sharks and rays) has been carefully studied of late years by Ruckert, Rabl, and H. E. Ziegler in particular, and is very important in the sense that this group is the oldest among living fishes, and their gastrulation can be derived directly from that of the cyclostoma by the accumulation of a large quantity of food-yolk. The oldest sharks (*Cestracion*) still have the unequal segmentation inherited from the cyclostoma. But while in this case, as in the case of




FIG. 55. Ovum-segmentation of a bony fish. A first cleavage of the stem-cell (yolk). B division of nutritive yolk into two parts (a and b). C the germinal disk (border) and the blastoderm (B) and the primitive (P) of nutritive yolk (a) and (b). a and b, a space between the blastoderm and the ovum, filled with a clear fluid.

difficult and uncertain in the case of the large ova that we find in the majority of the other fishes and in all the reptiles and birds. In these cases the food-yolk is, in the first place, comparatively colourless, the formative yolk being almost invisible beside it; and, in the second place, the food-yolk contains a quantity of different elements, which are known as "yolk-granules, yolk-globules, yolk-plates, yolk-bubbles, yolk-vesicles," and so on. Frequently these definite elements in the yolk have been described as real cells, and it has been wrongly stated that a portion of the embryonic body is built up from these cells. This is by no means the case. In every case, however large it is—and even when cell-nuclei travel into it during the cleavage of the border—the nutritive yolk remains a dead accumulation of food, which is taken into the gut during

the amphibian, the small ovum completely divides into cells in segmentation, this is no longer so in the great majority of the selachii (or *Elaemobranchii*). In these the contractility of the active protoplasm no longer suffices to break up the huge mass of the passive deutoplasm completely into cells; this is only possible in the upper or dorsal part, but not in the lower or ventral section. Hence we find in the primitive fishes a blastula with a small eccentric segmentation-cavity (Fig. 55 B), the wall of which varies greatly in composition. The circular border of the germinal disk which connects the roof and floor of the segmentation-cavity corresponds to the border-zone at the equator of the amphibian ovum. In the middle of its hinder border we have the beginning of the invagination of the primitive gut

(Fig. 56 *inf*); it extends gradually from this spot (which corresponds to the Russian anus of the amphibia) forward and around, so that the primitive mouth becomes first crescent-shaped and then




FIG. 55.—Discoid gastrula (*discogastrula*) of a lung fish. *a*, ectoderm; *b*, endoderm; *c*, food-yolk; *d*, primitive mouth; *e*, albuminous globule of the nutritive yolk; *f*, the globule of mucus; *g*, anterior coelomic cavity; *h*, partition between endoderm and ectoderm (anterior to the segmentation-cavity).

circular, and, as it opens wider, surrounds the half of the larger food-yolk.

Essentially different from the wide-

velachis is the narrow-mouthed discoid gastrula (*discogastrula*) of the *Amphioxus*, the reptiles, birds, and mon... between the two—as an intermediate stage—we have the *amphigastrea* of the amphibia. The latter has developed from the amphigastrea of the ganoids and dipneusts, whereas the discoid amniote gastrula has been evolved from the amphibian gastrula by the addition of food-yolk. This change of gastrulation is still found in the remarkable ophidia (*Gymnophiona*, *Cecilia*, or *Plesiosaurs*), serpent-like amphibia that live in moist soil in the tropics, and in many respects represent the transition from the gill-breathing amphibia to the lung-breathing reptiles. Their embryonic development has been explained by the fine studies of the brothers Sarasin of *Isothyris plumosa* at Ceylon (1887), and those of August Brauer of the

The bird's egg is particularly important for our purpose, because most of the chief studies of the development of the vertebrates are based on observations of the hen's egg, during

sch n obtain and study, and for this practical and obvious reason very rarely thoroughly investigated. But we can get hens' eggs in any quantity at any time, and, by means of artificial incubation, follow the development of the embryo step by step. The bird's egg differs considerably from the tiny mammal ovum in size, a large quantity of food-yolk accumulating within the original yolk or the protoplasm of the ovum. This is the yellow ball which we commonly call the yolk of the egg. In order to understand the bird's egg aright—for it is very often quite wrongly explained—we

and follow it from the very beginning of its development in the bird's ovary. We then see that the original ovum is a quite small, naked, and simple cell with a nucleus, not differing in either size or shape from the original ovum of the mammals and other animals (cf. Fig. 13 *E*). As in the case

the original or primitive ovum (*protoplast*) is covered with a continuous layer of cells. This... follicle, from which the ovum afterwards issues. Immediately underneath it the structureless yolk-membrane is secreted... in the yolk.

The small primitive ovum of the bird begins very early to take up into itself a quantity of food-stuff through the yolk-membrane, and work it up into the "yellow yolk." In this way the ovum




FIG. 56.—Longitudinal section through the blastula of

historical and comparative study of those that we can understand the difficult and obscure gastrulation of the amniotes.

enters on its second stage (the *morula*), which is many times larger than the first, but still only a single enlarged cell. Through the accumulation of the store of yellow yolk within the ball of protoplasm the nucleus it contains (the germinal vesicle) is forced to the surface of the ball. Here it is surrounded by a small quantity of protoplasm, and with this forms the lens-shaped formative yolk (Fig. 15 *b*). This is seen on the yellow yolk-ball, at a certain point of the surface, as a small round white spot—the "tread" (*exeatricule*). From this point a thread-like column of white nutritive yolk (*d*), which contains no yellow yolk-granules, and is softer than the yellow food-yolk, proceeds to the middle of the yellow yolk-ball, and forms there a small central globe of white yolk (Fig. 15 *d*). The whole of this white yolk is not sharply separated from the yellow yolk, which

(Fig. 5). First two equal segmentation-cells (*A*) are formed from the ovum. These divide into four (*B*), then into eight, sixteen (*C*), thirty-two, sixty-four, and so on. The cleavage of the cells is always preceded by a division of their nuclei. The cleavage surfaces between the segmentation-cells appear at the free surface of the tread as clefts. The first two divisions are vertical to each other, in the form of a cross (*B*). Then there are two more divisions, which cut the former at an angle of forty-five degrees. The tread, which thus becomes the germinal disk, now has the appearance of an eight-rayed star. A circular cleavage next taking place round the middle, the eight triangular cells divide into sixteen, of which eight are in the middle and eight distributed around (*C*). Afterwards circular clefts and radial clefts, directed towards the centre, alternate more or less



FIG. 16.—Longitudinal section of the blastula of a shark (*Prionace*) at the beginning of gastrulation (From Hildebrand). (Drawn from the left.) *P* fore end, *H* hind end, *B* segmentation-cavity, *A* blast. trace the primitive gut, *C* yolk-capsule, *D* the germinal yolk, *E* the nutritive yolk, *F* the nutritive yolk.

shows a slight trace of concentric layers in the hard-boiled egg (Fig. 15 *c*). We also find in the hen's egg, when we break the shell and take out the yolk, a round small white disk at its surface which corresponds to the tread. But this small white "germinal disk" is now further developed, and is really the gastrula of the chick. The body of the chick is formed from it alone. The whole white and yellow yolk-mass is without any significance for the formation of the embryo, it being merely used as food by the developing chick. The clear, glistening mass of albumin that surrounds the yellow yolk of the bird's egg, and also the hard chalky shell, are only formed within the oviduct round the impregnated ovum.

When the fertilisation of the bird's ovum has taken place within the mother's body, we find in the lens-shaped stem-cell the progress of flat, discoid segmentation

irregularly (*D*, *E*). In most of the amniotes the formation of concentric and radial clefts is irregular from the very first; and so also in the hen's egg. But the final outcome of the cleavage-process is once more the formation of a large number of small cells of a similar nature. As in the case of the fish-ovum, these segmentation-cells form a round, lens-shaped disk, which corresponds to the morula, and is embedded in a small depression of the white yolk. Between the lens-shaped disk of the morula-cells and the underlying white yolk a small cavity is now formed by the accumulation of fluid, as in the fishes. Thus we get the peculiar and not easily recognisable blastula of the bird (Fig. 16). The small segmentation-cavity (*A*) is very flat and much compressed. The upper or dorsal wall (*d*_w) is formed of a single layer of clear, distinctly separated cells; this

corresponds to the upper or animal hemisphere of the triton-blastula (Fig. 45). The lower or ventral wall of the flat dividing space (*zw*) is made up of larger and darker segmentation-cells; it corresponds to the lower or vegetal hemisphere of the blastula of the water-salamander (Fig. 45 *str*). The nuclei of the yolk-cells, which are in this case especially numerous at the edge of the lens-shaped blastula, travel into the white yolk, increase by cleavage, and contribute even to the further growth of the germinal disk by furnishing it with food-stuff.

The invagination or the folding inwards of the bird-blastula takes place in this

which was described for a long time as the "primitive groove." If we make a vertical section through this part, we see that a flat and broad cleft stretches under the germinal disk forwards from the primitive mouth; this is the primitive gut (Fig. 60 *md*). Its roof or dorsal wall is formed by the folded upper part of the blastula, and its floor or ventral wall by the white yolk (*wd*), in which a number of yolk-nuclei (*dk*) are distributed. There is a brisk multiplication of these at the edge of the germinal disk, especially in the neighbourhood of the sickle-shaped primitive mouth.

We learn from sections through later stages of this discoid bird-gastrula that




FIG. 57.—Diagram of discoid segmentation in the bird's ovum (enlarged about ten times). Only the formative yolk (the troch) is shown in these six figures (A-F); however, cleavage only takes place in them. The much larger food-yolk, which does not share in the cleavage, is left out and merely indicated by the dark ring without.

case also at the hinder pole of the subsequent chief axis, in the middle of the hind border of the round germinal disk (Fig. 59 *r*). At this spot we have the most brisk cleavage of the cells, hence the cells are more numerous and smaller here than in the fore-half of the germinal disk. The border-swelling or thick edge of the disk is less clear but whiter behind, and is more sharply separated from contiguous parts. In the middle of its hind border there is a white, crescent-shaped groove—Koller's sickle-groove (Fig. 59 *r*); a small projecting process in the centre of it is called the sickle-knob (*sk*). This important cleft is the primitive mouth,

the primitive gut-cavity, extending forward from the primitive mouth as a flat pouch, underlining the whole region of the round flat lens-shaped blastula (Fig. 61 *md*). At the same time, the segmentation-cavity gradually disappears altogether, the folded inner germinal layer (*sk*) placing itself from underneath on the overlying outer germinal layer (*sk*). The typical process of invagination, though greatly disguised, can thus be clearly seen in this case, as Goette and Rauber, and more recently Duval (Fig. 61), have shown.

The older embryologists (Pander, Baer, Ramak), and, in recent times especially,

His, Kollmer, and others, said that the two primary germinal layers of the hen's ovum—the oldest and most frequent subject of observation!—arose by horizontal cleavage of a simple germinal disk. In opposition to this accepted view, I affirmed in my *Gastrula Theory* (1873)

its surface. I endeavoured to establish this view by the derivation of the vertebrates from one source, and especially by proving that the birds descend from the reptiles, and thence from the amphibia. If this is correct, the discoid gastrula of the amniotes must have been formed by



FIG. 41.



FIG. 42.



FIG. 43.



FIG. 44.

FIG. 41.—Vertical section of the blastula of a hen (*Gallus gallus*). f.h. fertilized hen; yolk, central yolk, passing directly into the white yolk (yolk). (From *Journal*.)

FIG. 42.—The germinal disk of the hen (*Gallus gallus*) at the beginning of gastrulation: A before incubation, B at the first hour of incubation. (From *Journal*.) f.h. fertilized hen; yolk, central yolk; cell wall, outer cell wall; cell, inner cell wall; cell, inner cell wall.

FIG. 43.—Longitudinal section of the germinal disk of a chick (*Gallus gallus*) at gastrulation. yolk, central yolk; cell wall, outer cell wall; cell, inner cell wall; cell, inner cell wall; cell, inner cell wall.



FIG. 44.—Longitudinal section of the discoid gastrula of the nightingale. (From *Journal*.) yolk, central yolk; cell wall, outer cell wall; cell, inner cell wall; cell, inner cell wall; cell, inner cell wall.

that the discoid bird-gastrula, like that of all other vertebrates, is formed by folding (or invagination), and that this typical process is merely altered in a peculiar way and disguised by the immense accumulation of food-yolk and the flat spreading of the discoid blastula at one part of

its surface. I endeavoured to establish this view by the derivation of the vertebrates from one source, and especially by proving that the birds descend from the reptiles, and thence from the amphibia. If this is correct, the discoid gastrula of the amniotes must have been formed by

recently for the birds; and the same has been done for the reptiles by the fine studies of Kupffer, Bonnet, Wenckebach, and others. In the shield-shaped germinal disk of the lizard (Fig. 6a), the crocodile, the tortoise, and other reptiles, we find in the middle of the hind border (at the same spot as the sickle-groove in the bird) a transverse furrow (*w*), which leads into a flat, pouch-like, blind sac, the primitive gut. The fore (dorsal) and hind (ventral) lips of the transverse furrow correspond exactly to the lips of the primitive mouth (or sickle-groove) in the birds.

The gastrulation of the mammals must

I first advanced this fundamental principle in my essay *On the Gastrulation of Mammals* (1877), and sought to show in this way that I assumed a gradual degeneration of the food-yolk and the yolk-sac on the way from the proreptiles to the mammals. The ontogenetic process of adaptation," I said, "which has occasioned the atrophy of the rudimentary yolk-sac of the mammal, is perfectly clear. It is due to the fact that the young of the mammal, whose ancestors were certainly oviparous, now remain a long time in the womb. As the great store of food-yolk, which the oviparous ancestors gave to the egg, became superfluous in their descendants

owing to the long carrying in the womb, and the maternal blood in the wall of the uterus made itself the chief source of nourishment, the now useless yolk-sac was bound to atrophy by embryonic adaptation."

My opinion met with little

vigorous attack by Kolliker, Hensen, and His in particular. However, it has been gradually accepted, and has recently been firmly established by a large number of excellent studies of mammal gastrulation, especially by Edward Van Beneden's studies of the rabbit and bat, Selenka's on the marsupials and rodents, Heape's and Leberkuhn's on the mole, Kupffer and Kolbel's on the rodents, Bonnet's on the ruminants, etc. From the general comparative point of view, Carl Rabl in his theory of

the mesoderm, Oscar Hertwig in the latest edition of his *Mammal* (1902), and Hubrecht in his *Studies in Mammalian Embryology* (1891), have supported the opinion, and sought to derive the peculiarly modified gastrulation of the mammal from that of the reptile.

In the meantime (1884) the studies of Wilhelm Haacke and Caldwell provided a proof of the long-suspected and interesting fact, that the lowest mammals, the monotremes, lay eggs, like the birds and reptiles, and are not viviparous like the other mammals. Although the gastrulation of the monotremes was not really known until studied by Richard




FIG. 6a.—Germinal disk of the lizard (*Lacerta agilis*). (From Kupffer) is primitive mouth, *s* sickle, *st* embryonic shield, *h* and *h'* hind and dark gastrulative area.

be derived from this special embryonic development of the reptiles and birds. This latest and most advanced class of the vertebrates has, as we shall see afterwards, evolved at a comparatively recent date from an older group of reptiles, and all these amniotes must have come originally from a common stem-form. Hence the distinctive embryonic process of the mammal must have arisen by ontogenetic modifications from the older gastrulation of the reptiles and birds.

Until we admit this, we cannot demand the formation of the germinal layers in the mammal, and therefore

Semon in 1894, there could be little doubt, in view of the great size of their food-yolk, that their ovum-segmentation was discoid, and led to the formation of a sickle-mouthed discogastrula, as in the



1 opossum (*Didelphys*)
on *Selenka's* 2 the four
3 cleavage body, 4 unsegmented
albumen-membrane.

case of the reptiles and birds. Hence I had, in 1875 (in my essay on *The Gastrula and Ovum-segmentation of Animals*), counted the monotremes among the discoblastic vertebrates. This hypothesis

afterwards by the careful observations of Semon, he gave in the second volume of his great work, *Zoological Journeys in Australia* (1894), the first description and correct explanation of the discoid gastrulation of the monotremes. The fertilized ovum of the two living monotremes *Ecchinus* and *Ornithorhynchus* are balls of one-fifth of an inch in diameter, enclosed in a stiff shell; but they grow considerably during development, so that when laid the egg is three times as large. The structure of the plentiful yolk, and especially the relation of the yellow and the white yolk, are just the same as in the reptiles and birds. As with these, partial cleavage takes place at a spot on the surface at which the small formative yolk and the nucleus it encloses are found. First is formed a lens-shaped circular germinal disk. This is made up of several strata of cells, but it spreads over the yolk-ball, and thus becomes a one-layered blastula. If we then imagine the yolk it contains to be dissolved and replaced by a clear liquid, we have the characteristic blastula of the

higher mammals. In these the gastrulation proceeds in two phases, as Semon rightly observes: firstly, formation of the endoderm by cleavage at the centre and further growth at the edge; secondly, invagination. In the monotremes more primitive conditions have been retained better than in the reptiles and birds. In the latter, before the commencement of the gastrula-folding, we have, at least at the periphery, a two-layered embryo forming from the cleavage. But in the monotremes the formation of the coelogenous endoderm does not precede the invagination, hence in this case the construction of the germinal layers is less modified than in the other amniota.

The marsupials, a second sub-class, come next to the oviparous monotremes, the oldest of the mammals. But as in their case the food-yolk is already atrophied, and the little ovum develops within the mother's body, the partial cleavage has been reconverted into total. One section of the marsupials still show points of agreement with the monotremes, while another section of them, according to the splendid investigations of Selenka, form a connecting-link between these and the placental....

The fertilized ovum of the opossum (*Didelphys*) divides, according to Selenka, first into two, then four, then eight equal cells; hence the segmentation is at first



FIG. 4.—Blastula of the opossum (*Didelphys*). (From Selenka.) a general view of the blastula, a vaginal pole, b mother-cell of the endoderm, c endoderm cells, d nucleus of the blastula, e unsegmented yolk-balls (interior of the food-yolk), f albumen membrane.

1 or homogeneous. But in the case of the cleavage a larger cell, distinguished by its less clear plasma and its containing more yolk-granules (the mother cell of the endoderm, Fig. 64 vs),

separates from the others; the latter multiply more rapidly than the former. As, further, a quantity of fluid gathers in the morula, we get a round blastula, the wall of which is of varying thickness, like that of the amphioxus (Fig. 38 B) and the amphibia (Fig. 43). The upper or animal hemisphere is formed of a large number of small cells; the lower or vegetal hemisphere of a small number of large cells. One of the latter, distinguished by its size (*P*), lies at the vegetal pole of the *a*-axis, at the point where the primitive mouth afterwards appears; it is the mother-cell of the entoderm; it now begins to multiply by cleavage, and the daughter-cells (Fig. 65 *d*) spread out from this spot

gastrula (Fig. 66) gradually changes into globular, a larger quantity of fluid accumulating in the vesicle. At the same time, the entoderm spreads further and further over the inner surface of the ectoderm (*e*). A globular form, the wall of which consists of two thin simple strata of cells; cells of the outer germinal layer rounder, and those of the inner layer flatter. In the region of the primitive mouth (*p*) the cells are less flattened, and multiply briskly. From this point—the hind (ventral) lip of the primitive mouth, which extends in a central cleft, the primitive groove—the construction of the mesoderm proceeds.

Gastrulation is still more modified and



FIG. 65.



FIG. 66.

FIG. 65.—Blastula of the opossum (*Didelphis*) at the beginning of gastrulation. (From Schmidt.) *a* ectoderm; *b* entoderm; *c* animal pole; *d* primitive mouth at the vegetal pole; *e* gastrulation-cleft; *f* nucleus of yellow ball (pole of the reduced egg-cell) embedded cleft (without yellow ball).

FIG. 66.—Oval gastrula of the opossum (*Didelphis*) about eight hours old. (From Schmidt.) (enlarged view).

over the inner surface of the blastula, though at first only over the vegetal hemisphere. The less clear entodermic cells (*b*) are distinguished at first by their rounder shape and darker nuclei from the higher, clearer, and longer entodermic cells (*c*); afterwards both are greatly flattened, the inner blastodermic cells more than the outer.

The un-nucleated yellow-balls and cleft (Fig. 65 *d*) that we find in the fluid of the blastula in these marsupials are very remarkable; they are the relics of the atrophied food-yolk, which was developed in their ancestors, the monotremes, and in the reptiles.

In the further course of the gastrulation of the opossum the oval shape of the

curtailed cenogametocyte in the placentals than in the marsupials. It was first accurately known to us by the distinguished investigations of Edward Van Beneden in 1875. The first object of study being the ovum of the rabbit. But as man also belongs to this sub-class, and as his as yet unstudied gastrulation cannot be materially different from that of the other placentals, it merits the closest attention. We have, in the first place, the peculiar feature that the two first segmentation-cells that proceed from the cleavage of the fertilized ovum (Fig. 68) are of different sizes and natures; the difference is sometimes greater, sometimes less (Fig. 66). One of these first daughter-cells of the ovum is a little

larger, clearer, and more transparent than the other. Further, the smaller cell takes a colour in carmine, osmium, etc., more strongly than the larger. By repeated cleavage of it a morula is formed, and from this a blastula, which changes in a very characteristic way into the greatly modified gastrula. When the number of the segmentation-cells in the morula reaches a hundred or more

six (in the rabbit, about seventy hours after impregnation) the fetus assumes a form very like the archigastrea (Fig. 72). The spherical embryo consists of a central mass of thirty-two soft, round cells with dark nuclei, which are flattened into polygonal shape by mutual pressure, and colour dark-brown with osmic acid (Fig. 72 *b*). This dark central group of cells is surrounded by a lighter spherical membrane, consisting of sixty-four cube-shaped, small, and fine-grained cells which lie close together in a single stratum, and only colour slightly in osmic acid (Fig. 72 *c*). The authors who regard this embryonic form as the primary gastrula of the placental conceive the outer layer as the ectoderm and the inner as the entoderm. The entodermic membrane is only interrupted at one spot, one, two, or three of the ectodermic cells being loose there. These form the yolk-stopper, and fill up the mouth of the gastrula (*a*). The central primitive gut-cavity (*d*) is full of entodermic cells. The uni-axial type of the mammal gastrula is accentuated in this way. However, opinions still differ considerably as to the real nature of this "provisional gastrula" of the placental and its relation to the blastula into which it is converted.

As the gastrulation proceeds a large spherical blastula is formed from this peculiar solid amphigastrea of the placental, as we saw in the case of the marsupial. The accumulation of fluid in the solid gastrula (Fig. 73 *a*) leads to the formation of an eccentric cavity, the group of the darker entodermic cells (*ay*) remaining directly attached at one spot with the round enveloping stratum of the lighter ectodermic cells (*cp*). This spot corresponds to the original primitive mouth (prostoma or blastoporus). From this important spot the inner germinal layer spreads all round on the inner surface of the outer layer, the cell-stratum of which forms the wall of the hollow sphere; the extension proceeds from the vegetal towards the animal pole.

The ontogenetic gastrulation of the placental has been greatly modified by secondary adaptation in the various groups of this most advanced and youngest sub-class of the mammals. Thus, for instance, we find in many of the rodents (guinea-pigs, mice, etc.) apparently a temporary inversion of the two germinal layers. This is due to a folding of the blastodermic wall by what is called the "girdler," a plug-shaped growth of Rauber's "roof-layer." It is a thin layer of flat epithelial cells, that is freed from the surface of the blastoderm in some of the rodents; it has no more significance in connection with the general course of placental gastrulation than the conspicuous departure from



FIG. 72.—Longitudinal section through the oral gastrula of the opossum (Fig. 72). (From Wilson.) *a* primitive mouth, *b* ectoderm, *c* entoderm.

globular shape in the blastula of some of the unguulates. In some pigs and ruminants it grows into a thread-like, long and thin tube.

Thus the gastrulation of the placental of the amphibious, the primitive form, is reduced to the original type, the invagination of a modified blastula. Its character is that the folded part of the blastoderm does not form a completely closed (only open at the primitive mouth) blind sac, as is usual; but this blind sac has a wide opening at the ventral curve (opposite to the dorsal mouth); and through this opening the primitive gut communicates from the first with the embryonic cavity of the blastula. The folded crest-shaped

emboderm grows with a free circular border on the inner surface of the entoderm towards the vegetal pole; when it has reached this, and the inner surface of the blastula is completely grown over, the primitive gut is closed. This remarkable

their ancestors (the reptiles), is atrophied. This proves the essential unity of gastrulation in all the vertebrates, in spite of the striking differences in the various classes.

In order to complete our consideration of the important processes of segmenta-




FIG. 64.—Stem-cell of the mammalian ovum (from the rabbit). *a* dead sperm-cells, *b* zona pellucida, *c* protoplasm of the stem-cell, *d* nucleus.




FIG. 65.—The first four segmentation-cells of the rabbit ovum (and lighter).




FIG. 66.—Indifferent cleavage of the mammalian ovum (from the rabbit). The stem-cell has divided into two unequal cells, one lighter (*a*) and one darker (*b*). *a* zona pellucida, *b* outer blastomeres, *c* protoplasm, *d* dead sperm-cells.




FIG. 67.—Mammalian ovum with eight segmentation-cells (from the rabbit). *a* four larger and lighter cells, *b* four smaller and darker cells, *c* zona pellucida, *d* outer blastomeres, *e* inner blastomeres.

direct transition of the primitive gut-cavity into the segmentation-cavity is explained simply by the assumption that in most of the mammals the yolk-mass, which is still possessed by the eldest forms of the class (the monotremes) and

tion and gastrulation, we will, in conclusion, cast a brief glance at the fourth chief type—superficial segmentation. In the vertebrates this form is not found at all. But it plays the chief part in the large stem of the articulates—the insects,

spiders, myriapods, and crabs. The distinctive form of gastrula that comes of it is the "vesicular gastrula" (*Perrinitia*).

In the ova which undergo this superficial cleavage the formative yolk is sharply divided from the nutritive yolk, as in the preceding cases of the ova of birds, reptiles, fishes, etc.; the formative yolk alone undergoes cleavage. But while in the ova with discoid gastrulation the formative yolk is not in the centre, but at one pole of the uni-axial ovum, and the food-yolk gathered at the other pole, in the ova with superficial cleavage we find the formative yolk spread over the whole surface of the ovum; it encloses spherically the food-yolk, which is accumulated in the middle of the ova. Gastrulation only affects the formative yolk, it is bound to be

achieved by it. As a rule, it proceeds in regular geometrical progression. In the end the whole of the formative yolk divides into a number of small and homogeneous cells, which lie close together in a single stratum on the entire surface of the ovum, and form a superficial blastoderm. This blastoderm is a simple, completely closed vesicle, the internal cavity of which is entirely full of food-yolk. This real blastula only differs from that of the primitive ova in its chemical composition. In the latter the content is water or a watery jelly, in the former it is food-yolk.

begins, there is no difference between the morula. The two stages rather

When the blastula is fully formed, we find the important folding or invagination that determines gastrulation. The space between the skin-layer and the gut-layer (the remainder of the segmentation-cavity) remains full of food-yolk, which is gradually used up. This is the only material difference between our vesicular gastrula (*Perrinitia*) and the original form of the bell-gastrula (*Archegastrula*). Clearly the one has been developed from the other in the course of time, owing to the accumulation of food-yolk in the centre of the ovum.¹

We must count it an important advance that we are thus in a position to reduce the embryonic phenomena in the different groups of animals to these four principal forms of segmentation and gastrulation. Of these four forms we must regard one only as the original palingenetic, and the other three as anagenetic and derivative. The unequal, the discoid, and the superficial segmentation have all clearly arisen by secondary adaptation from the primary segmentation, and the chief cause of their development has been the gradual formation of the food-yolk, and the increasing antithesis between animal and vegetal halves of the ovum, or between ectoderm (skin-layer) and entoderm (gut-layer).




FIG. 14.—Gastrula of the planula stage (see p. 14). Longitudinal section through the axis. 1. ectodermic cells (slightly larger and smaller), 2. entodermic cells (thirty-two, darker and larger), 3. central entodermic cells filling the primitive gut-cavity, 4. spherical ectodermic cells, stopping up the opening of the primitive mouth (yolk-stopper in the *Rhipidura* genus).

The numbers of careful studies of animal gastrulation that have been made in the last decades have completely established the

hitherto first advanced in the years 1872-76. For a time they were greatly disputed by many embryologists. Some said that the original embryonic form of the metazoa was not the gastrula, but the "planula"—a double-walled vesicle with closed cavity and without mouth-aperture; the latter was supposed to pierce through gradually. It was afterwards shown that this planula (found in several sponges, etc.) was a later evolution from the gastrula.

¹ On the reduction of all forms of gastrulation to the original palingenetic form see especially the lucid treatment of the subject in Arnold Lang's *Manual of Comparative Zoology* (1898), Part I.

¹ On the reduction of all forms of gastrulation to

It was also shown that what is called delamination—the rise of the two primary germinal layers by the folding of the surface of the blastoderm (for instance, in the *Geryoneis* and other rindans)—was a secondary formation, due to ontogenetic

they attach themselves to the inner wall of the blastula, and form a second external epithelial layer—that is to say, the entoderm. In these and many other contraventions of modern embryology the first requisite for clear and natural explanation




FIG. 35.—Gastrula of the rabbit. A as a solid spherical cluster of cells, B changing into the embryo. *A*₁ primitive mouth, *A*₂ ectoderm, *A*₃ endoderm.

variations from the original invagination of the blastula. The same may be said of what is called "immigration," in which certain cells or groups of cells are detached from the simple layer of the blastoderm, and travel into the interior of the blastula.

is a careful and discriminative distinction between palingenetic (hereditary) and ontogenetic (adaptive) processes. If this is properly attended to, we find evidence everywhere of the biogenetic law.

CHAPTER X.

THE CŒLOM THEORY

THE two "primary germinal layers" which the gastrula theory has shown to be the first foundation in the construction of the body are found in this simplest form throughout life only in animals of the lowest grade—in the gastruæda, olynthus (the stem-form of the sponges), hydra, and simple animals. In all the other animals new strata are formed subsequently between these two primary body-layers, and these are generally comprehended under the title of the middle layer, or *mesoderm*. As a rule, the

various products of this middle layer afterwards constitute the great bulk of the animal frame, while the original entoderm, or internal germinal layer, is restricted to the clothing of the alimentary canal and its glandular appendages; and, on the other hand, the ectoderm, or external germinal layer, furnishes the outer clothing of the body, the skin and nervous system.

so large groups of the lower such as the sponges, corals, and is, the middle germinal layer

as a single connected mass, and of the body is developed from it; these have been called the three-layered metazoa, in opposition to the layered animals described. Like the two

animals, they have no body-cavity—that is to say, no cavity distinct from the alimentary system. On the other hand,

they have this real body-cavity (*coeloma*), and so are called *coelomates*; we can distinguish them from the three-layered animals, which

develop from the two primary layers. To the same class belong all true vermians (excepting the platodes), and also

those evolved from them—molluscs, echinoderms, arthropods, tunicates, and vertebrates.

multicellular animal-body, is the oldest and most important organ of all the metazoa, and, together with the primitive mouth, is formed in

the earliest stages of development. In the gut, it is only at a much later stage that the body-cavity, which is entirely wanting

in the coelenterata, is developed in the metazoa between the body wall. The two cavities are entirely different in content and purpose.

The alimentary cavity (*enteron*) serves the purpose of digestion; it contains water and food taken from without, as well as the pulp (*chymus*) formed from

the body-cavity, quite distinct from the gut and closed externally, has nothing to do with digestion; it encloses the gut




FIG. 74




FIG. 75

FIG. 74 and 75—Diagram of the four-spore stage of a metazoan, showing a cross-section through the alimentary cavity. FIG. 74 is an animal, FIG. 75 is a vegetative. Ar, outermost layer; Am, skin-sense-layer; Gd, gut-gland-layer; Co, coelomic layer. Ar shows the beginning of the coelomic cavity, a beginning of the metazoan.

The body-cavity (*coeloma*) is therefore a new acquisition of the animal body, much younger than the alimentary system, and of great importance. I first pointed out this fundamental significance of the coelom in my *Monograph on the Sponges* (1872), in the section which draws a distinction between the body-cavity and the gut-cavity, and which follows immediately on the germ-layer theory and the ancestral tree of the animal kingdom (the first sketch of the gastraea theory). Up to that time these two principal cavities of the animal body had been confused, or very imperfectly distinguished; chiefly because Leuckart, the founder of the coelenterata group (1848), has attributed a body-cavity, but not a gut-cavity, to these lowest metazoa. In reality, the truth is just the other way about.

itself and its glandular appendages, and also contains the sexual products and a certain amount of blood or lymph, a fluid that is transmitted through the ventral wall.

As soon as the body-cavity appears, the ventral wall is found to be separated from the enclosing body-wall, but the two continue to be directly connected at various points. We can also then always distinguish a number of different layers of tissue in both walls—at least two in each. These tissue-layers are formed originally from four different simple cell-layers, which are the much-discussed four secondary germinal layers. The outermost of these, the skin-sense-layer (FIG. 74, 75 Ar), and the innermost, the gut-gland-layer (*Gd*), remain at first simple epithelia or covering-layers. The one covers the outer

surface of the ventral wall; hence they are called confining or limiting layers. Between them are the two middle-layers, somata, which enclose the body-cavity.



Fig. 76—Coelomaria of sagitta, (sagitta with a permanent mouth).

no permanent mouth.

The four secondary germinal layers are so distributed in the structure of the body in all the coelomaria (or all metazoa that have a body-cavity) that the outer two, joined fast together, constitute the body-wall, and the inner two the ventral wall. Each of the walls is a limiting layer.

tissues, and glands and nerves, middle layers form the great bulk of the fibrous tissue, muscles, and connective matter. Hence the latter have been called fibrous or muscular layers. The outer middle layer, which lies on the inner side of the skin-somite-layer, is the skin fibre-layer, the inner middle layer, which attaches from without to the ventral glandular layer, is the ventral fibre layer. The former is usually called briefly the parietal, and the latter the visceral layer or mesoderm. Of the many different names that have been given to the four secondary germinal layers, the following are the most common:

- | | | | |
|---|--------------------------------|--|---|
| I. Skin-somite-layer (outer limiting layer). | II. Parietal layer (muscular). | III. Epithelial layer (visceral). | IV. Gut-gland-layer (inner limiting layer). |
| The two secondary germinal layers of the body-wall. | | The two secondary germinal layers of the ventral wall. | |

The first scientist to recognise and clearly distinguish the four secondary germinal layers was Baer. It is true he was not quite clear as to their origin and further.

several mistakes in detail in explaining them. But, on the whole, their great importance did not escape him. However, in later years his view had to be given up in consequence of more accurate observations. Remak then propounded a three-layer theory, which was generally accepted. These theories of cleavage, however, began to give way thirty years when Kowalevsky (1871) showed in the case of *Sagitta* (a very clear typical subject of gastrulation) the middle germinal layers and the two outer layers arise not by cleavage, but by folding—by a secondary invagination of the primary inner germ-layer. This invagination or folding proceeds from the primitive mouth, at the two sides of which (right and left) a couple of pouches are formed. As these coelom-pouches or coelom-walls detach themselves from the primitive gut, a double body-cavity is formed (Figs 74-6).

The same kind of coelom-formation as in *Sagitta* was afterwards found by Kowalevsky in *Brachiopod*.




Fig. 77—Coelomaria of sagitta, in section. (From Haeckel.) D dorsal side, V ventral side, of outer germinal layer, two ventral somites, B body-cavity, M parietal somites, A outer germinal layer.

In ontogeny—E. Ray-Lankester and F. Balfour. On the strength of these and other studies, as well as most extensive research of their own, the brothers Oscar and Richard Hertwig constructed in 1881

the Coelom Theory. In order to appreciate fully the great merit of this illuminating and helpful theory, one must remember what a chaos of contradictory views was then represented by the "problem of the mesoderm," or the much-disputed "question of the origin of the middle germinal layer." The coelom theory brought some light and order into this infinite confusion by establishing the following points: 1. The body-cavity originates in the great majority of animals (especially in all the vertebrates) in the same way as in *Ascaris*: a couple of pouches or sacs are formed by folding inwards at the primitive mouth, between the two primary germinal layers; as these pouches detach from the primitive gut, a pair of coelom-sacs (right and left) are formed, the coalescence of these produces simple body-cavity. 2. A

coelom-embryo develop, not as a pair of hollow pouches, but as solid layers of cells (in the shape of a pair of mesodermal streaks)—as happens in the higher vertebrates—we have a secondary (ontogenetic) modification of the primary (palingenetic) structure, the two walls of the pouches, inner and outer, have been pressed together by the expansion of the large food-yolk. 3. Hence the mesoderm from the first of two genetically distinct layers, which do not originate by the cleavage of a primary simple middle layer (as Kernak supposed). 4. These two middle layers have, in all vertebrates, and the great majority of the invertebrates, the same radical significance for the construction of the animal body: the inner middle layer, or the visceral mesoderm (gut-fibre layer), attaches itself to the original ectoderm, and forms the fibrous, muscular, and connective part of the visceral wall, the outer middle layer, or the parietal mesoderm (skin-fibre-layer), attaches itself to the original ectoderm, and forms the fibrous, muscular, and connective part of the body-wall. 5. It is only at the point of origination, the primitive mouth and its vicinity, that the four secondary germinal layers are directly connected; from this point the two middle layers advance forward separately between the two primary germinal layers, to which they severally attach themselves. 6. The further separation or differentiation of the four secondary germinal layers and their division into the various tissues and organs take place especially in the later fore-part or head of the embryo, and

extend backwards from there towards the primitive mouth.

All animals in which the body-cavity demonstrably arises in this way from the primitive gut (vertebrates, tunicates, echinoderms, articulates, and a part of the Vermalia) were comprised by the Hertwigs under the title of *enterocoela*, and were contrasted with the other groups of the *pseudocoela* (with false body-cavity) and the *acoelenterata* (with no body-cavity). However, this radical distinction and the views as to classification which it occasioned have been shown to be untenable. Further, the absolute differences in transformation which the Hertwigs set up between the enterocoela and pseudocoela cannot be sustained.

For these and other reasons their coelom-theory has been much criticised and partly



FIG. 11.—Section of a young *Ascaris*. (From Hertwig.) *gut*, visceral cavity, *ect* and *ectoderm* and outer limiting layers, *vis* and *visceral* and inner middle layers, *par* body-cavity, *par* and *parietal* and visceral mesoderm.

abandoned. Nevertheless, it has rendered a great and lasting service in the solution of the difficult problem of the mesoderm, and a material part of it will certainly be retained. I consider it an especial merit of the theory that it has established the identity of the development of the two middle layers in all the vertebrates, and has traced them as ontogenetic modifications back to the original palingenetic form of development that we still find in the amphioxus. Carl Rabl comes to the same conclusion in his able *Theory of the Mesoderm*, and so do Ray-Lankester, Rauber, Kupffer, Ruckert, Selenska, Hatschek, and others. There is a general agreement in these and many other recent writers that all the different forms of coelom-construction, like those of gastrulation, follow one and the same strict hereditary law in the vast vertebrate stem; in spite of their apparent differences, they

are all only coelogenetic modifications of one paligenetic type, and this original type has been preserved for us down to the present day by the invaluable amphioxus.

But before we go into the regular colonisation of the amphioxus, we will glance at that of the arrow-worm (*Sagitta*), a remarkable deep-sea worm that is interesting in many ways for comparative anatomy and ontogeny. On the one hand, the transparency of the body and the embryo, and, on the other hand, the typical simplicity of its embryonic development, make the *sagitta* a most instructive object in connection with various problems. The class of the *cheilognathus*, which is only represented by the cognate genera of *Sagitta* and *Spadella*, is in another respect

afterwards arises). The two sacs are at first separated by a couple of folds of the ectoderm (Fig. 76 *fm*), and are still connected with the primitive gut by wide apertures; they also communicate for a short time with the dorsal side (Fig. 77 *d*). Soon, however, the coelom-pouches completely separate from each other and from the primitive gut, at the same time they enlarge so much that they close round the primitive gut (Fig. 78). But in the middle line of the dorsal and ventral sides the pouches remain separated, their approaching walls joining here to form a thin vertical partition, the mesentery (*dm* and *vm*). Thus *Sagitta* has throughout life a double body-cavity (Fig. 78 *bc*), and the gut is fastened to the body-wall both above and below by a mesentery—below



FIG. 76.




FIG. 78.

FIGS. 76 AND 78.—Transverse section of amphioxus-larva. (From Metchnikoff.) FIG. 76 at the commencement of coelom formation (and without segmentation); FIG. 78 at the stage with four primitive coelomic sacs. *sk*, *sk*, *mp*, *mp* outer and inner, and medullary plates; *ch*, chorda; "—" and "—" disjunctions of the ectoderm-pouches, *bc* body-cavity.

also a most remarkable branch of the extensive vermetaria stem. It was therefore very gratifying that Oscar Hertwig (1886) fully explained the anatomy, classification, and evolution of the cheilognathus in his careful monograph.

The spherical blastula that arises from the impregnated ovum of the *sagitta* is converted by a folding at one pole into a typical archigastrea, entirely similar to that of the *Amoebocera* which I described (Chapter VIII., Fig. 29). This oval, uniaxial cup-larva (circular in section) becomes bilateral (or tri-axial) by the growth of a couple of coelom-pouches from the primitive gut (Figs. 76, 77). To the right and left-shaped fold appears towards the top pole (where the permanent mouth, as

by the ventral mesentery (*vm*), and above by the dorsal mesentery (*dm*). The inner layer of the two coelom-pouches (*mp*) attaches itself to the ectoderm (*sk*), and forms with it the visceral wall. The outer layer (*mp*) attaches itself to the ectoderm (*sk*), and forms with it the outer body-wall. Thus we have in *Sagitta* a perfectly clear and simple illustration of the original colonisation of the enterocoela. This paligenetic fact is the more important, as the greater part of the two body-cavities in *Sagitta* changes afterwards into sexual glands—the fore or female part into a pair of ovaries, and the hind or male part into a pair of testicles.

Colonisation takes place with equal clearness and transparency in the case of

the amphioxus, the lowest vertebrate, and its nearest relatives, the invertebrate tunicates, the sea-squirts. However, in these two stems, which we class together as *Chordata*, this important process is more complex, as two other processes are associated with it—the development of the chorda from the entoderm and the separation of the medullary plate or nervous centre from the ectoderm. Here again the skullless amphioxus has preserved to our own time by tenacious heredity the

while it has been more or less modified by embryonic adaptation in all the other vertebrates (with skulls). Hence we must once more thoroughly understand the paleogenetic embryonic features of the lancelet before we go on to consider the congenetic forms of the craniota.

border of the concave medullary plate fold towards each other and grow underneath the horny-plate, a cylindrical tube is formed, the medullary tube (Fig. 82 a); this quickly detaches itself altogether from the horny-plate. At each side of the medullary tube, between it and the alimentary tube (Figs. 79–82 *ab*), two parallel longitudinal folds grow out of the dorsal wall of the alimentary tube, and form the two coelom-pouches (Figs. 80 and 81 *A*). This part of the entoderm,

of the middle germinal layer, is shown darker than the rest of the inner germinal layer in Figs. 79–82. The edges of the folds meet, and thus form closed tubes (Fig. 81 in section).

During this interesting process the outline of a third very important organ,



FIG. 81.



FIG. 82.

FIGS. 81 AND 82.—Transverse section of amphioxus embryo. FIG. 81 at the stage with the coelom, while FIG. 82 at the stage with closed coelom. (From Haeckel's) *ab* outer germinal layer, *mp* medullary plate, *gut* dorsal wall of alimentary tube, *ch* coelom-pouch, *md* middle germinal layer (dark), *ect* ectoderm.

The coelomation of the amphioxus, which was first observed by Kowalevsky in 1867, has been very carefully studied since by Hatschek (1881). According to him, there are first formed on the bilateral gastrula we have already considered (Figs. 36, 37) three parallel longitudinal folds—one single ectodermal fold in the central line of the dorsal surface, and a pair of entodermal folds at the two sides of the former. The broad ectodermal fold that first appears in the middle line of the flattened dorsal surface, and forms a shallow longitudinal groove, is the beginning of the central nervous system, the medullary tube. Thus the primary outer germinal layer divides into two parts, the middle medullary plate (Fig. 81 *mp*) and the horny-plate (*ah*), the beginning of the outer skin or epidermis. As the parallel

the chorda or axial rod, is being formed between the two coelom-pouches. This first foundation of the skeleton, a solid cylindrical cartilaginous rod, is formed in the middle line of the dorsal primitive gut-wall, from the ectodermal cell-streak that remains here between the two coelom-pouches (Figs. 79–82 *ch*). The chorda appears at first in the shape of a flat longitudinal fold or a shallow groove (Figs. 80, 81); it does not become a solid cylindrical cord until after separation from the primitive gut (Fig. 84). Hence we might say that the dorsal wall of the primitive gut forms three parallel longitudinal folds at this important period—one single fold and a pair of folds. The single middle fold becomes the chorda, and lies immediately below the groove of the ectoderm, which becomes the medullary

tube; the pair of folds to the right and left lie at the sides between the former and the latter, and form the coelom-pouches. The part of the primitive gut that remains after the cutting off of these three dorsal primitives organs is the per-

(Figs. 83, 84, in the third period of development according to Hatachek). (Strabo and Plinius give the name of *cordula* or *cordula* to young fish larvae.) I ascribe the utmost phylogenetic significance to it, as it is found in all the chorda-



FIG. 83.



FIG. 84.

Figs. 83 and 84.—Chordula of the amphioxus. FIG. 83 shows longitudinal section (seen from the left). FIG. 84 transverse section. (From Hatachek.) In FIG. 84 the coelom-pouches are omitted, in order to show the chordula more clearly. FIG. 84 is rather diagrammatic. a hardy-plate, b medullary tube, c wall of nerve (c dorsal, c' ventral), d chorda, e notochord, f coelom, g coelom-pouch, h gut-cavity, i gut dorsal wall, j gut ventral wall, k yolk-cells in the latter, l primitive mouth, m mouth-let, n pharynx, o pharynx, p pharynx, q pharynx, r pharynx, s pharynx, t pharynx, u pharynx, v pharynx, w pharynx, x pharynx, y pharynx, z pharynx, aa pharynx, ab pharynx, ac pharynx, ad pharynx, ae pharynx, af pharynx, ag pharynx, ah pharynx, ai pharynx, aj pharynx, ak pharynx, al pharynx, am pharynx, an pharynx, ao pharynx, ap pharynx, aq pharynx, ar pharynx, as pharynx, at pharynx, au pharynx, av pharynx, aw pharynx, ax pharynx, ay pharynx, az pharynx, ba pharynx, bb pharynx, bc pharynx, bd pharynx, be pharynx, bf pharynx, bg pharynx, bh pharynx, bi pharynx, bj pharynx, bk pharynx, bl pharynx, bm pharynx, bn pharynx, bo pharynx, bp pharynx, bq pharynx, br pharynx, bs pharynx, bt pharynx, bu pharynx, bv pharynx, bw pharynx, bx pharynx, by pharynx, bz pharynx, ca pharynx, cb pharynx, cc pharynx, cd pharynx, ce pharynx, cf pharynx, cg pharynx, ch pharynx, ci pharynx, cj pharynx, ck pharynx, cl pharynx, cm pharynx, cn pharynx, co pharynx, cp pharynx, cq pharynx, cr pharynx, cs pharynx, ct pharynx, cu pharynx, cv pharynx, cw pharynx, cx pharynx, cy pharynx, cz pharynx, da pharynx, db pharynx, dc pharynx, dd pharynx, de pharynx, df pharynx, dg pharynx, dh pharynx, di pharynx, dj pharynx, dk pharynx, dl pharynx, dm pharynx, dn pharynx, do pharynx, dp pharynx, dq pharynx, dr pharynx, ds pharynx, dt pharynx, du pharynx, dv pharynx, dw pharynx, dx pharynx, dy pharynx, dz pharynx, ea pharynx, eb pharynx, ec pharynx, ed pharynx, ee pharynx, ef pharynx, eg pharynx, eh pharynx, ei pharynx, ej pharynx, ek pharynx, el pharynx, em pharynx, en pharynx, eo pharynx, ep pharynx, eq pharynx, er pharynx, es pharynx, et pharynx, eu pharynx, ev pharynx, ew pharynx, ex pharynx, ey pharynx, ez pharynx, fa pharynx, fb pharynx, fc pharynx, fd pharynx, fe pharynx, ff pharynx, fg pharynx, fh pharynx, fi pharynx, fj pharynx, fk pharynx, fl pharynx, fm pharynx, fn pharynx, fo pharynx, fp pharynx, fq pharynx, fr pharynx, fs pharynx, ft pharynx, fu pharynx, fv pharynx, fw pharynx, fx pharynx, fy pharynx, fz pharynx, ga pharynx, gb pharynx, gc pharynx, gd pharynx, ge pharynx, gf pharynx, gg pharynx, gh pharynx, gi pharynx, gj pharynx, gk pharynx, gl pharynx, gm pharynx, gn pharynx, go pharynx, gp pharynx, gq pharynx, gr pharynx, gs pharynx, gt pharynx, gu pharynx, gv pharynx, gw pharynx, gx pharynx, gy pharynx, gz pharynx, ha pharynx, hb pharynx, hc pharynx, hd pharynx, he pharynx, hf pharynx, hg pharynx, hh pharynx, hi pharynx, hj pharynx, hk pharynx, hl pharynx, hm pharynx, hn pharynx, ho pharynx, hp pharynx, hq pharynx, hr pharynx, hs pharynx, ht pharynx, hu pharynx, hv pharynx, hw pharynx, hx pharynx, hy pharynx, hz pharynx, ia pharynx, ib pharynx, ic pharynx, id pharynx, ie pharynx, if pharynx, ig pharynx, ih pharynx, ii pharynx, ij pharynx, ik pharynx, il pharynx, im pharynx, in pharynx, io pharynx, ip pharynx, iq pharynx, ir pharynx, is pharynx, it pharynx, iu pharynx, iv pharynx, iw pharynx, ix pharynx, iy pharynx, iz pharynx, ja pharynx, jb pharynx, jc pharynx, jd pharynx, je pharynx, jf pharynx, jg pharynx, jh pharynx, ji pharynx, jj pharynx, jk pharynx, jl pharynx, jm pharynx, jn pharynx, jo pharynx, jp pharynx, jq pharynx, jr pharynx, js pharynx, jt pharynx, ju pharynx, jv pharynx, jw pharynx, jx pharynx, jy pharynx, jz pharynx, ka pharynx, kb pharynx, kc pharynx, kd pharynx, ke pharynx, kf pharynx, kg pharynx, kh pharynx, ki pharynx, kj pharynx, kk pharynx, kl pharynx, km pharynx, kn pharynx, ko pharynx, kp pharynx, kq pharynx, kr pharynx, ks pharynx, kt pharynx, ku pharynx, kv pharynx, kw pharynx, kx pharynx, ky pharynx, kz pharynx, la pharynx, lb pharynx, lc pharynx, ld pharynx, le pharynx, lf pharynx, lg pharynx, lh pharynx, li pharynx, lj pharynx, lk pharynx, ll pharynx, lm pharynx, ln pharynx, lo pharynx, lp pharynx, lq pharynx, lr pharynx, ls pharynx, lt pharynx, lu pharynx, lv pharynx, lw pharynx, lx pharynx, ly pharynx, lz pharynx, ma pharynx, mb pharynx, mc pharynx, md pharynx, me pharynx, mf pharynx, mg pharynx, mh pharynx, mi pharynx, mj pharynx, mk pharynx, ml pharynx, mm pharynx, mn pharynx, mo pharynx, mp pharynx, mq pharynx, mr pharynx, ms pharynx, mt pharynx, mu pharynx, mv pharynx, mw pharynx, mx pharynx, my pharynx, mz pharynx, na pharynx, nb pharynx, nc pharynx, nd pharynx, ne pharynx, nf pharynx, ng pharynx, nh pharynx, ni pharynx, nj pharynx, nk pharynx, nl pharynx, nm pharynx, nn pharynx, no pharynx, np pharynx, nq pharynx, nr pharynx, ns pharynx, nt pharynx, nu pharynx, nv pharynx, nw pharynx, nx pharynx, ny pharynx, nz pharynx, oa pharynx, ob pharynx, oc pharynx, od pharynx, oe pharynx, of pharynx, og pharynx, oh pharynx, oi pharynx, oj pharynx, ok pharynx, ol pharynx, om pharynx, on pharynx, oo pharynx, op pharynx, oq pharynx, or pharynx, os pharynx, ot pharynx, ou pharynx, ov pharynx, ow pharynx, ox pharynx, oy pharynx, oz pharynx, pa pharynx, pb pharynx, pc pharynx, pd pharynx, pe pharynx, pf pharynx, pg pharynx, ph pharynx, pi pharynx, pj pharynx, pk pharynx, pl pharynx, pm pharynx, pn pharynx, po pharynx, pp pharynx, pq pharynx, pr pharynx, ps pharynx, pt pharynx, pu pharynx, pv pharynx, pw pharynx, px pharynx, py pharynx, pz pharynx, qa pharynx, qb pharynx, qc pharynx, qd pharynx, qe pharynx, qf pharynx, qg pharynx, qh pharynx, qi pharynx, qj pharynx, qk pharynx, ql pharynx, qm pharynx, qn pharynx, qo pharynx, qp pharynx, qq pharynx, qr pharynx, qs pharynx, qt pharynx, qu pharynx, qv pharynx, qw pharynx, qx pharynx, qy pharynx, qz pharynx, ra pharynx, rb pharynx, rc pharynx, rd pharynx, re pharynx, rf pharynx, rg pharynx, rh pharynx, ri pharynx, rj pharynx, rk pharynx, rl pharynx, rm pharynx, rn pharynx, ro pharynx, rp pharynx, rq pharynx, rr pharynx, rs pharynx, rt pharynx, ru pharynx, rv pharynx, rw pharynx, rx pharynx, ry pharynx, rz pharynx, sa pharynx, sb pharynx, sc pharynx, sd pharynx, se pharynx, sf pharynx, sg pharynx, sh pharynx, si pharynx, sj pharynx, sk pharynx, sl pharynx, sm pharynx, sn pharynx, so pharynx, sp pharynx, sq pharynx, sr pharynx, ss pharynx, st pharynx, su pharynx, sv pharynx, sw pharynx, sx pharynx, sy pharynx, sz pharynx, ta pharynx, tb pharynx, tc pharynx, td pharynx, te pharynx, tf pharynx, tg pharynx, th pharynx, ti pharynx, tj pharynx, tk pharynx, tl pharynx, tm pharynx, tn pharynx, to pharynx, tp pharynx, tq pharynx, tr pharynx, ts pharynx, tt pharynx, tu pharynx, tv pharynx, tw pharynx, tx pharynx, ty pharynx, tz pharynx, ua pharynx, ub pharynx, uc pharynx, ud pharynx, ue pharynx, uf pharynx, ug pharynx, uh pharynx, ui pharynx, uj pharynx, uk pharynx, ul pharynx, um pharynx, un pharynx, oo pharynx, op pharynx, oq pharynx, or pharynx, os pharynx, ot pharynx, ou pharynx, ov pharynx, ow pharynx, ox pharynx, oy pharynx, oz pharynx, va pharynx, vb pharynx, vc pharynx, vd pharynx, ve pharynx, vf pharynx, vg pharynx, vh pharynx, vi pharynx, vj pharynx, vk pharynx, vl pharynx, vm pharynx, vn pharynx, vo pharynx, vp pharynx, vq pharynx, vr pharynx, vs pharynx, vt pharynx, vu pharynx, vv pharynx, vw pharynx, vx pharynx, vy pharynx, vz pharynx, wa pharynx, wb pharynx, wc pharynx, wd pharynx, we pharynx, wf pharynx, wg pharynx, wh pharynx, wi pharynx, wj pharynx, wk pharynx, wl pharynx, wm pharynx, wn pharynx, wo pharynx, wp pharynx, wq pharynx, wr pharynx, ws pharynx, wt pharynx, wu pharynx, wv pharynx, ww pharynx, wx pharynx, wy pharynx, wz pharynx, xa pharynx, xb pharynx, xc pharynx, xd pharynx, xe pharynx, xf pharynx, xg pharynx, xh pharynx, xi pharynx, xj pharynx, xk pharynx, xl pharynx, xm pharynx, xn pharynx, xo pharynx, xp pharynx, xq pharynx, xr pharynx, xs pharynx, xt pharynx, xu pharynx, xv pharynx, xw pharynx, xx pharynx, xy pharynx, xz pharynx, ya pharynx, yb pharynx, yc pharynx, yd pharynx, ye pharynx, yf pharynx, yg pharynx, yh pharynx, yi pharynx, yj pharynx, yk pharynx, yl pharynx, ym pharynx, yn pharynx, yo pharynx, yp pharynx, yq pharynx, yr pharynx, ys pharynx, yt pharynx, yu pharynx, yv pharynx, yw pharynx, yx pharynx, yy pharynx, yz pharynx, za pharynx, zb pharynx, zc pharynx, zd pharynx, ze pharynx, zf pharynx, zg pharynx, zh pharynx, zi pharynx, zj pharynx, zk pharynx, zl pharynx, zm pharynx, zn pharynx, zo pharynx, zp pharynx, zq pharynx, zr pharynx, zs pharynx, zt pharynx, zu pharynx, zv pharynx, zw pharynx, zx pharynx, zy pharynx, zz pharynx.



FIG. 85.



FIG. 86.

Figs. 85 and 86.—Chordula of the amphioxus (the rapid-sider). (From Gortz.) FIG. 85 shows longitudinal section (seen from the left). FIG. 86 transverse section (slightly diagrammatic). Labeled as in Figs. 83 and 84.

manent gut; its endoderm is the gut-gland-layer or enteric layer.

I give the name of *chordula* or *chordula* to the embryonic stage of the vertebrate organism which is represented by the amphioxus larva at this period

animals (unicates as well as vertebrates) in essentially the same form. Although the accumulation of food-yolk greatly modifies the form of the chordula in the higher vertebrates, it remains the same in its main features throughout. In all

a discoid expansion. These cenogenetic modifications seem to be so great that until twenty years ago these important processes were totally misunderstood. It was generally believed that the body-cavity in man and the higher vertebrates was due to the division of a simple middle layer, and that the latter arose by cleavage from one or both of the primary germinal layers. The truth was brought to light at last by the comparative embryological research of the Hertwigs. They showed in their *Celom Theory* (1881) that all vertebrates are true enterocoela, and that in every case a pair of coelom-pouches are developed from the primitive gut by folding. The cenogenetic chondula-forms of the craniate adult therefore be derived from the paligenetic embryology of the amphioxus in the same way as I had previously proved for their gastrula-forms.

The chief difference between the coelomation of the acrania (*amphioxus*) and the other vertebrates (with skulls - craniates) is that the two coelom-folds of the primitive gut in the former are from the first hollow vesicles, filled with fluid, but in the latter are empty pouches, the layers of which (inner and outer) close with each other. In common parlance we still call a pouch or pocket by that name, whether it is full or empty. It is different in ontogeny, in some of our embryological literature ordinary logic does not count for very much. In many of the manuals and large treatises on this science it is proved that vesicles, pouches, or sacs deserve that name only when they are inflated and filled with a clear fluid. When they are not so filled (for instance, when the primitive gut of the gastrula is filled with yolk, or when the walls of the empty coelom-pouches are pressed together), these vesicles must not be cavities any longer, but "solid structures."

The accumulation of food-yolk in the ventral wall of the primitive gut (Figs 85, 86) is the simple cause that converts the sac-shaped coelom-pouches of the acrania into the leaf-shaped coelom-streaks of the craniates. To convince ourselves of this we need only compare, with Hertwig, the paligenetic coelomula of the amphioxus (Figs. 80, 81) with the corresponding cenogenetic form of the amphibia (Figs. 85-90), and construct the simple diagram that connects the two (Figs. 87, 88). If we

imagine the ventral half of the primitive gut-wall in the amphioxus embryo (Figs. 79-84) distended with food-yolk, the vascular coelom-pouches (*pb*) must be pressed together by this, and forced to extend in the shape of a thin double plate between the gut-wall and body-wall (Figs. 86, 87). This expansion follows a downward and forward direction. They are not directly connected with these two walls. The real unbroken connection between the two middle layers and the primary germ-layers is found right at the back, in the region of the primitive mouth (Fig. 87 *u*). At this important spot we have the source of embryonic development (*blastopore*), or "zone of growth," from which the coelomation (and also the gastrulation) originally proceeds.

Hertwig even succeeded in showing, in the coelomula-embryo of the water salamander (*Triton*), between the first structures of the two middle layers, the relic of



Fig. 88.—Transverse section of the chordate-embryo of a bird (from a hen's egg at the close of the first day of incubation). (From *Amphioxus*). *A* hyp-plate (notochord), *B* ventral plate, *pb* dorsal folds of same, *rb* ventral furrow, *ch* chorda, *wp* median (inner) part of the middle layer (median wall of the coelom-pouches), *lp* lateral (outer) part of same, or lateral plates, *ms* structure of the body-cavity, *st* gut-limb-layer.

the body-cavity, which is represented in the diagrammatic transitional form (Figs. 87, 88). In sections both through the primitive mouth itself (Fig. 89) and in front of it (Fig. 90) the two middle layers (*pb* and *rb*) diverge from each other, and disclose the two body-cavities as narrow clefts. At the primitive mouth itself (Fig. 90 *u*) we can penetrate into them from without. It is only here at the border of the primitive mouth that we can show the direct transition of the two middle layers into the two limiting layers or primary germinal layers.

The structure of the chorda also shows the same features in these coelomula-embryos of the amphibia (Fig. 91) as in the amphioxus (Figs. 79-84). It arises from the endodermic cell-streak, which forms the middle dorsal line of the primitive gut, and occupies the space between the flat coelom-pouches (Fig. 91 *A*).

While the nervous centre is formed here in the middle line of the back and separated from the ectoderm as "modul-
lary tube," there takes place at the same
time, directly underneath, the severance
of the chorda from the entoderm (Fig. 91
A, B, C). Under the chorda is formed
(out of the ventral entodermic half of the
gastrula) the permanent gut or visceral
cavity (*enteron*) (Fig. 91 B, *ab*). This
is done by the confluence, under the
chorda in the median line, of the two
dorsal side-borders of the gut-gland-layer
(*sk*), which were previously separated by
the chorda-plate (Fig. 91 A, *sk*); those
now alone form the clothing of the visceral
cavity (*sk*) (enteroderm, Fig. 91 C). All
these important modifications take place
at first in the fore or head-part of the
embryo, and spread backwards from
there; here at the hinder end, the region
of the primitive mouth, the important
border of the mouth (or *propharynx*),

possible as a matter of fact; even the older
illustrations showed an essential identity
of features. Thus forty years ago Kölliker
gave, in the first edition of his *Human
Embryology* (1861), some sections of the
chicken-embryo, the features of which
could at once be reduced to those already
described and explained in the sense of
Hertwig's coelom-theory. A section
through the embryo in the hatched hen's
egg towards the close of the first day of
incubation shows in the middle of the
dorsal surface a broad ectodermic modul-
lary groove (Fig. 92 B f), and under-
neath the middle of the chorda (*sk*) and
at each side of it a couple of broad meso-
dermic layers (*sp*). These enclose a
narrow space or cleft (*war*), which is
nothing else than the structure of the
body-cavity. The two layers that enclose
it—the upper parietal layer (*hpl*) and the
lower visceral layer (*dl*)—are pressed
together from without, but clearly distin-



FIG. 92.—Transverse section of the vertebrate-embryo of a bird (from a hen's egg on the second day of incubation). (From Kölliker.) A, *hystrophila*, not modul-lary tube, *sk*, chorda, *war*, primitive mesenteron, *hpl*, parietal layer, *dl*, gut-gland-layer, *sp*, primitive-lary passage, *ab*, primitive mouth, *sk*, gut-gland-layer.

remains for a long time the space of
development or the zone of fresh construc-
tion, in the further building-up of the
organism. One has only to compare care-
fully the illustrations given (Figs. 85-91)
to see that, as a fact, the ontogenetic
coelomation of the amphibia can be
deduced directly from the palingenetic
form of the neurula (Figs. 79-84).

The same principle holds good for
the amniotes, the reptiles, birds, and
mammals, although in this case the pro-
cesses of coelomation are more modified
and more difficult to identify on account
of the colossal accumulation of food-yolk
and the corresponding notable flattening
of the germinal disk. However, as the
whole group of the amniotes has been
developed at a comparatively late date
from the class of the amphibia, their
coelomation must also be directly trace-
able to that of the latter. This is really

guishable. This is even clearer a little
later, when the medullary furrow is closed
into the nerve-tube (Fig. 93 *nr*).

Special importance attaches to the fact
that here again the four secondary ger-
minal layers are already sharply distinct,
and easily separated from each other.
There is only one very restricted area in
which they are connected, and actually
pass into each other; this is the region of
the primitive mouth, which is contracted
in the amniotes into a dorsal longitudinal
cleft, the primitive groove. Its two lateral
lip-borders form the *propharyngeal* streak, which
has long been recognised as the most
important embryonic source and starting-
point of further processes. Sections
through this primitive streak (Figs. 94
and 95) show that the two primary ger-
minal layers grow at an early stage (in
the discoid gastrula of the chick, a few
hours after incubation) into the primitive

streak (*x*), and that the two middle layers extend outward from this thickened axial plate (*y*) to the right and left between the former. The plates of the celom-layers, the parietal stem-fibre-layer (*w*) and the visceral gut-fibre-layer (*z*), are seen to be still pressed close together, and only diverge later to form the body-cavity. Between the inner borders of the two flat celom-pouches lies the chorda (Fig. 95 *x*), which here again develops from the middle line of the dorsal wall of the primitive gut.

Colonization takes place in the vertebrates in just the same way as in the birds and reptiles. This was to be ex-

posed; the four secondary germinal layers consists of a single stratum of cells.

Finally, we must point out, as a fact of the utmost importance for our anthropogeny and of great general interest, that the four-layered celomula of man has just the same construction as that of the rabbit (Fig. 96). A vertical section that Count Spen made through the primitive mouth or streak of a very young human germinal disk (Fig. 97) clearly shows that here again the four secondary germ-layers are inseparably connected only at the primitive streak, and that here also the two flattened celom-pouches (*w* and *z*) extend outwards to right and left from



FIG. 94



FIG. 95

FIGS. 94 AND 95.—Transverse section of the primitive streak (primitive mouth) of the chick. FIG. 94 a few hours after the commencement of incubation. FIG. 95 a little later (From Hildebrand). A horizontal line is not a plate, *w* stem-fibre-layer, *z* gut-fibre-layer, *d* parietal-layer, *y* primitive streak or axial plate, in which all four germinal layers meet, *x* structure of the chorda, *v* nucleus of the later primitive kidneys.

posed, as the characteristic gastrulation of the mammal has descended from that of the reptiles. In both cases a discoid gastrula with primitive streak arises from the segmented ovum, a two-layered germinal disk with large and small hinder primitive mouth. Here again the two primary germinal layers are only directly connected (Fig. 95 *y*) along the primitive streak (at the folding-point of the blastula), and from this spot (the border of the primitive mouth) the middle germinal layer (*mb*) grow out to right and left between the preceding. In the fine illustration of the celomula of the rabbit which Van Beneden has given us (Fig. 96) one can clearly see that each of the

the primitive mouth between the outer and inner germinal layers. In this case, too, the middle germinal layer consists from the first of two separate strata of cells, the parietal (*mp*) and visceral (*mv*) mesoblasts.

These concordant results of the best recent investigations (which have been confirmed by the observations of a number of scientists I have not enumerated) prove the unity of the vertebrate-stem in point of colonization, no less than of gastrulation. In both respects the invaluable amphioxus—the sole survivor of the acrania—is found to be the original model that has preserved for us in paleogenetic form by a tenacious heredity these

most important embryonic processes. From this primary model of construction we can ontogenetically deduce all the embryonic forms of the other vertebrates, the craniota, by secondary modifications. My thesis of the universal formation of the gastrula by folding of the blastula has now been clearly proved for all the vertebrates; so also has been Hertwig's thesis of the origin of the middle germinal layers by the folding of a couple of coloma-pouches which appear at the border of

typical, unarticulated, worm-like form, which has an axial chorda between the dorsal nerve-tube and the ventral gut-tube. This instructive chordula (Figs. 83-86) provides a valuable support of our phylogeny; it indicates the important moment in our stem-history at which the stem of the chordonia (tunicates and vertebrate) parted far ever from the divergent stems of the other metazoa (articulars, echinoderms, and molluscs).

I may express here my opinion, in the




FIG. 66.—Transverse section of the primitive groove (or primitive mouth) of a rabbit. (From *Van Beneden*). *pr* primitive mouth at top of gastrula (primitive fold); *ad* and *al* dorsal and ventral folds; *mp* and *ms* middle germinal layers; *la* lateral layer of the notochord.




FIG. 67.—Transverse section of the primitive mouth (or groove) of a human embryo (at the notochord stage). (From *Chapt. Shaw*). *pr* primitive mouth; *ad* top of gastrula (primitive fold); *al* and *al* dorsal and ventral folds; *mp* middle germinal layers; *ms* middle germinal layers; *la* lateral layer of the notochord.

the primitive mouth. Just as the gastrula-theory explains the origin and identity of the two primary layers, so the coloma-theory explains those of the four secondary layers. The point of origin is always the properistoma, the border of the original primitive mouth of the gastrula, at which the two primary layers pass directly into each other.

Moreover, the coloma is important as the immediate source of the chordula, the embryonic reproduction of the ancient

form of a chordula-theory, that the characteristic chordula-larva of the chordonia has in reality this great significance—it is the typical reproduction (preserved by heredity) of the ancient common stem-form of all the vertebrates and tunicates, the long-extinct *Chordula*. We will return in the twentieth chapter to these worm-like ancestors, which stand out as luminous points in the obscure stem-history of the invertebrate ancestors of our race.

CHAPTER XI.

THE VERTEBRATE CHARACTER OF MAN

We have now secured a number of firm standing-places in the labyrinthine course of our individual development by our study of the important embryonic forms which we have called the *ovula*, *morula*, *blastula*, *gastrula*, *coelomula*, and *chordula*. But we have still in front of us the difficult task of deriving the complicated scheme of the human body, with all its different parts, organs, members, etc., from the simple form of the *chordula*. We have previously considered the origin of this four-layered embryonic form from the two-layered *gastrula*. The two primary germinal layers, which form the entire body of the *gastrula*, and the two middle layers of the *coelomula* that develop between them, are the four simple cell-strata, or *epithelia*, which alone go to the formation of the complex body of man and the higher animals. It is so difficult to understand this construction that we will first seek a companion who may help us out of many difficulties.

This helpful associate is the science of comparative anatomy. Its task is, by comparing the fully-developed bodily forms in the various groups of animals, to learn the general laws of organization according to which the body is constructed, at the same time, it has to determine the affinities of the various groups by critical appreciation of the degrees of difference between them. Formerly, this work was conceived in a teleological sense, and it was sought to find traces of the plan of the Creator in the actual purposeful organization of animals. But comparative anatomy has gone much deeper since the establishment of the theory of descent; its philosophical aim now is to explain the variety of organic forms by adaptation, and their similarity by heredity. At the same time, it has to recognise in the shades of difference in form the degree of blood-relationship, and make an effort to construct the ancestral tree of the animal world. In this way, comparative anatomy enters into the closest relations with comparative

embryology on the one hand, and with the science of classification on the other.

Now, when we ask what position man occupies among the other organisms according to the latest teaching of comparative anatomy and classification, and how man's place in the zoological system is determined by comparison of the mature bodily forms, we get a very definite and significant reply, and this reply gives us extremely important conclusions that enable us to understand the embryonic development and its evolutionary purport. Since Cuvier and Beer, since the immense progress that was effected in the early decades of the nineteenth century by these two great zoologists, the opinion has generally prevailed that the whole animal kingdom may be distributed in a small number of great divisions or types. They are called types because a certain typical or characteristic structure is constantly preserved within each of these large sections. Since we applied the theory of descent to this doctrine of types, we have learned that this common type is an outcome of heredity; all the animals of one type are blood-relatives, or members of one stem, and can be traced to a common ancestral form. Cuvier and Beer set up four of these types: the *vertebrates*, *articulates*, *molluscs*, and *radiates*. The first three of these are still retained, and may be conceived as natural phylogenetic units, as *stems* or *phyla* in the sense of the theory of descent. It is quite otherwise with the fourth type—the *radiata*. These animals, hitherto known as yet at the beginning of the nineteenth century, were made to form a sort of lumber-room, into which were cast all the lower animals that did not belong to the other three types. As we obtained a closer acquaintance with them in the course of the last sixty years, it was found that we must distinguish among them from four to eight different types. In this way the total number of animal stems or *phyla* has been raised to eight or twelve (cf. Chapter XX).

These twelve stems of the animal kingdom are, however, by no means co-ordinate and independent types, but have definite relations, partly of subordination, to each other, and a very different phylogenetic meaning. Hence they must not be arranged simply in a row one after the other, as was generally done until thirty years ago, and is still done in some manuals. We must distribute them in three subordinate principal groups of very different value, and arrange the various stems phylogenetically on the principles which I laid down in my *Diagramm der Sponges*, and developed in the *Study of the Gastraea Theory*. We have first to distinguish the unicellular animals (*Protozoa*) from the multicellular tissue-forming (*metazoa*). Only the latter exhibit the important processes of segmentation and gastrulation; and they alone have a primitive gut, and form germinal layers and tissues.

The metazoa, the tissue-animals or gut-animals, then sub-divide into two main sections, according as a body-cavity is or is not developed between the primary germinal layers. We may call these the *coelenterata* and *celenterata*, the former are often also called *amphioxys* or *celenterata*, and the latter *blastozoa*. This division is the more important as the coelenterata (without coelom) have no blood and blood-vessels, nor an anus. The celenterata (with body-cavity) have generally an anus, and blood and blood-vessels. There are four stems belonging to the coelenterata: the gastricals ("primitive-gut animals"), sponges, cnidaria, and platodes. Of the celenterata we can distinguish six stems: the vermalia at the bottom represent the common stem-group (derived from the platodes) of these, the other five typical stems of the celenterata—the molluscs, echinoderms, articulates, tunicates, and vertebrates—being evolved from them.

Man is, in his whole structure, a true vertebrate, and develops from an impregnated ovum in just the same characteristic way as the other vertebrates. There can no longer be the slightest doubt about this fundamental fact, nor of the fact that all the vertebrates form a natural phylogenetic unity, a single stem. The whole of the members of this stem, from the amphioxys and the cyclostoma to the ape and man, have the same characteristic disposition, connection, and development of the central organs, and arise in the

same way from the common embryonic form of the chordula. Without going into the difficult question of the origin of this stem, we must emphasize the fact that the vertebrate stem has no direct affinity whatever to five of the other ten stems; these five isolated phyla are the sponges, cnidaria, molluscs, articulates, and echinoderms. On the other hand, there are important and, to an extent, close phylogenetic relations to the other five stems—the protozoa (through the amebae), the gastricals (through the blastula and gastrula), the platodes, and vermalia (through the celenterata), and the tunicates (through the chordula).

How we are to explain these phylogenetic relations in the present state of our knowledge, and what place is assigned to the vertebrates in the animal kinship-tree, will be considered later (Chapter XX). For the present our task is to make plainer the vertebrate character of man, and especially to point out the chief peculiarities of organization by which the vertebrate stem is profoundly separated from the other eleven stems of the animal kingdom. Only after these comparative-anatomical considerations shall we be in a position to attack the difficult question of our embryology. The development of even the simplest and lowest vertebrate from the simple chordula (Figs. 83-86) is so complicated and difficult to follow that it is necessary to understand the organic features of the fully-formed vertebrate in order to grasp the course of its embryonic evolution. But it is equally necessary to confine our attention, in this general anatomic description of the vertebrate-body, to the essential facts, and pass by all the unessential. Hence, in giving now an anatomic description of the chief features of the vertebrate and its external organization, I omit all the subordinate points, and restrict myself to the most important characteristics.

Much, of course, will seem to the reader to be essential that is only of subordinate and secondary interest, or even not essential at all, in the light of comparative anatomy and embryology. For instance, the skull and vertebral column and the extremities are non-essential in this sense. It is true that these parts are very important *physiologically*; but for the *morphological* conception of the vertebrate they are not essential, because they are only found in the higher, not the lower, vertebrates. The lowest vertebrates have

neither skull nor vertebrae, and no extremities or limbs. Even the human embryo passes through a stage in which it has no skull or vertebrae; the trunk is quite simple, and there is yet no trace of arms and legs. At this stage of development man, like every other higher vertebrate, is essentially similar to the simplest vertebrate form, which we now find in only one living specimen. This one lowest vertebrate that merits the closest study—undoubtedly the most interesting of all the vertebrates after man—is the famous lancelet or amphioxus, to which we have already often referred. As we are going to study it more closely later on (Chapters XVI and XVII), I will only make one or two passing observations on it here.

The amphioxus lives buried in the sand of the sea, is about one or two inches in length, and has, when fully developed, the shape of a very simple, longish, lancelet-like leaf; hence its name of the lancelet. The narrow body is compressed on both sides, almost equally pointed at the fore and hind ends, without any trace of external appendages or articulation of the body into head, neck, breast, abdomen, etc. Its whole shape is so simple that its first discoverer thought it was a naked snail. It was not until much later—half a century ago—that the tiny creature was studied more carefully, and was found to be a true vertebrate. More recent investigations have shown that it is of the greatest importance in connection with the comparative anatomy and ontogeny of the vertebrates, and therefore with human phylogeny. The amphioxus reveals the great secret of the origin of the vertebrates from the invertebrate vermiform, and in its development and structure connects directly with certain lower tunicates, the ascidia.

When we make a number of sections of the body of the amphioxus, firstly vertical longitudinal sections through the whole body from end to end, and secondly transverse sections from right to left, we get anatomic pictures of the utmost instructiveness (cf. Figs. 98-102). In the main they correspond to the ideal which we form, with the aid of comparative anatomy and ontogeny, of the primitive type or build of the vertebrate—the long extinct form to which the whole stem owes its origin. As we take the phylogenetic unity of the vertebrate stem to be beyond dispute, and assume a common

origin from a primitive stem-form for all the vertebrates, from amphioxus to man, we are justified in forming a definite morphological idea of this primitive vertebrate (*Protophyllus* or *Vertebra*). We need only imagine a few slight and unessential changes in the real sections of the amphioxus in order to have this ideal anatomic figure or diagram of the primitive vertebrate form, as we see in Figs. 98-102. The amphioxus departs so little from this primitive form that we may, in a certain sense, describe it as a modified "primitive vertebrate."

The outer form of our hypothetical primitive vertebrate was at all events very simple, and probably more or less similar to that of the lancelet. The bilateral or bilateral-symmetrical body is stretched out lengthways and compressed at the sides (Figs. 98-102), oval in section (Figs. 101, 102). There are no external articulation and no external appendages, in the shape of limbs, legs, or fins. On the other hand, the division of the body into two sections, head and trunk, was probably clearer in *Protophyllus* than it is in its little-changed ancestor, the amphioxus. In both animals the fore or head-half of the body contains different organs from the trunk, and different on the dorsal from on the ventral side. As this important division is found even in the sea-squirt, the remarkable invertebrate stem-relative of the vertebrates, we may assume that it was also found in the prochordoma, the common ancestor of both stems. It is also very pronounced in the young larva of the cyclostoma; this fact is particularly interesting, as this palæogenetic larva-form is in other respects also an important connecting-link between the higher vertebrates and the acrania.

The head of the acrania, or the anterior half of the body (both of the real amphioxus and the ideal *protophyllus*), contains the branchial (gill) gut and heart in the ventral section and the brain and sense-organs in the dorsal section. The trunk, or posterior half of the body, contains the hepatic (liver) gut and sexual

* The ideal figure of the vertebrate as given in Figs. 98-102 is a hypothetical scheme or diagram. It has been chiefly constructed on the basis of the amphioxus, but with a certain attention to the comparative anatomy and ontogeny of the acrania and appendicularia on the one hand, and of the cyclostoma and teleostei on the other. This diagram has no pretensions whatever to be an "exact picture," but merely an attempt to construct hypothetically the unknown and long extinct vertebrate stem-form, an ideal "ancestry."




FIG. 91.



FIG. 92.



FIG. 93.




FIG. 94.




FIG. 95.

FIG. 91-95.—The lampbrush (Pisces 91-95). FIG. 91, of anterior (from the left). FIG. 92, of posterior. FIG. 93, of transverse. FIG. 94, of transverse section through the head (to the left through the pharynx, to the right through the gill-chamber). FIG. 95, of transverse section of the trunk (to the right a pharyngeal bone is visible). a, aorta; sh, scales; na, notch; r, ray; b, brain; ma, muscle; l, liver; o, operculum; f, faint; p, pharynx; ha, heart; g, gill; s, scales; t, tail; d, dorsal; v, vertebra; a, artery; f, faint.

FIG. 96-97.—The lampbrush (Pisces 96-97). FIG. 96, of anterior (from the left). FIG. 97, of posterior. FIG. 98, of transverse. FIG. 99, of transverse section through the head (to the left through the pharynx, to the right through the gill-chamber). FIG. 100, of transverse section of the trunk (to the right a pharyngeal bone is visible). a, aorta; sh, scales; na, notch; r, ray; b, brain; ma, muscle; l, liver; o, operculum; f, faint; p, pharynx; ha, heart; g, gill; s, scales; t, tail; d, dorsal; v, vertebra; a, artery; f, faint.

glands in the ventral part, and the spinal marrow and most of the muscles in the dorsal part.

In the longitudinal section of the ideal vertebrate (Fig. 98) we have in the middle of the body a thin and flexible, but stiff, cylindrical rod, pointed at both ends (*ca*). It goes the whole length through the middle of the body, and forms, as the central skeletal axis, the original structure of the later vertebral column. This is the axial rod, or *chorda dorsalis*, also called *chorda vertebralis*, vertebral cord, axial cord, dorsal cord, *notochorda*, or, briefly, *chorda*. This solid, but flexible and elastic, axial rod consists of a cartilaginous mass of cells, and forms the inner axial skeleton or central frame of the body; it is only found in vertebrates and tunicates, not in any other animals. As the first structure of the spinal column it has the same radical significance in all vertebrates, from the amphioxus to man. But it is only in the amphioxus and the cyclostoma that the axial rod retains its simplest form throughout life. In man and all the higher vertebrates it is found only in the earlier embryonic period, and is afterwards replaced by the articulated vertebral column.

The axial rod or *chorda* is the real solid chief axis of the vertebrate body, and at the same time corresponds to the ideal long-axis, and serves to direct us with some confidence in the orientation of the principal organs. We therefore take the vertebrate-body in its original, natural disposition, in which the long-axis lies horizontally, the dorsal side upward and the ventral side downward (Fig. 98). When we make a vertical section through the whole length of this long axis, the body divides into two equal and symmetrical halves, right and left. In each half we have *originally* the same organs in the same disposition and connection; only their disposal in relation to the vertical plane of section, or median plane, is exactly reversed: the left half is the reflection of the right. We call the two halves *anatomers* (opposed-parts). In the vertical plane of section that divides the two halves the sagittal ("arrow") axis, or "dorsoventral axis," goes from the back to the belly, corresponding to the sagittal suture of the skull. But when we make a horizontal longitudinal section through the *chorda*, the whole body divides into a dorsal and a ventral half. The line of section that passes through

the body from right to left is the transverse, frontal, or lateral axis.

The two halves of the vertebrate body that are separated by this horizontal transverse axis and by the *chorda* have quite different characters. The dorsal half is mainly the animal part of the body, and contains the greater part of what are called the animal organs, the nervous system, muscular system, osseous system, etc.—the instruments of movement and sensation. The ventral half is essentially the vegetative half of the body, and contains the greater part of the vertebrate's vegetative organs, the visceral and vascular systems, sexual system, etc.—the instruments of nutrition and reproduction. Hence in the construction of the dorsal half it is chiefly the outer, and in the construction of the ventral half chiefly the inner, germinal layer that is engaged. Each of the two halves develops in the shape of a tube, and encloses a cavity in which another tube is found. The dorsal half contains the narrow spinal-column cavity or vertebral canal above the *chorda*, in which lies the tube-shaped central nervous system, the medullary tube. The ventral half contains the much more spacious visceral cavity or body-cavity *underneath* the *chorda*, in which we find the alimentary canal and all its appendages.

The medullary tube, as the central nervous system or psychic organ of the vertebrate is called in its first stage, consists, in man and all the higher vertebrates, of two different parts: the large brain, contained in the skull, and the long spinal cord which stretches from there over the whole dorsal part of the trunk. Even in the primitive vertebrate this composition is plainly indicated. The fore half of the body, which corresponds to the head, encloses a knob-shaped vesicle, the brain (*gb*), this is prolonged backwards into the thin cylindrical tube of the spinal marrow (*sj*). Hence we find here this very important psychic organ, which accomplishes sensation, will, and thought, in the vertebrates, in its simplest form. The thick wall of the nerve-tube, which runs through the long axis of the body immediately over the axial rod, encloses a narrow central canal filled with fluid (Figs. 98-100 +). We still find the medullary tube in this very simple form for a time in the embryo of all the vertebrates, and it retains this form in the amphioxus throughout life;

only in the latter case the cylindrical medullary tube barely indicates the separation of brain and spinal cord. The lancelet's medullary tube runs nearly the whole length of the body, above the chorda, in the shape of a long thin tube of almost equal diameter throughout, and there is only a slight swelling of it right at the front to represent the rudiment of a cerebral lobe. It is probable that this peculiarity of the amphioxus is connected with the partial atrophy of its head, as the ascidian larva on the one hand and the young cyclostoma on the other clearly show a division of the vesicular brain, or head marrow, from the thinner, tubular spinal marrow.

Probably we must trace to the same phylogenetic cause the defective nature of the sense organs of the amphioxus, which we will describe later (Chapter XVI). Prosopodius, on the other hand, probably had three pairs of sense-organs, though of a simple character, a pair of, or a single olfactory depression, right in front (Figs. 98, 99, 100), a pair of eyes (*an*) in the lateral walls of the brain, and a pair of simple auscultatory vesicles (*g*) behind. There was also, perhaps, a single parietal or "pincal" eye at the top of the skull (*epiphysus*, *e*).

In the median plane (or middle plane, dividing the bilateral body into right and left halves) we have in the scranium, underneath the chorda, the mesenteric and visceral tube, and above it the medullary tube; and above the latter a membranous partition of the two halves of the body. With this partition is connected the mass of connective tissue which acts as a sheath both for the medullary tube and the underlying chorda, and is, therefore, called the chord-sheath (*parichorda*). It originates from the dorsal and median part of the colon-pouches, which we shall call the skeleton plate or "sclerotom" in the craneo-embryo. In the latter the chief part of the skeleton—the vertebral column and skull—develops from this chord-sheath; in the scranium it remains in simple form as a soft connective matter, from which are formed the membranous partitions between the various muscular plates or myotomes (Figs. 98, 99, 100).

To the right and left of the cord-sheath, at each side of the medullary tube and the underlying axial rod, we find in all the vertebrates the large masses of muscle that constitute the musculature of the

trunk and effect its movements. Although these are very elaborately differentiated and connected in the developed vertebrate (corresponding to the various parts of the bony skeleton), in our ideal primitive vertebrate we can distinguish only two pairs of these principal muscles, which run the whole length of the body parallel to the chorda. These are the upper (dorsal) and lower (ventral) lateral muscles of the trunk. The upper (dorsal) muscles, or the original dorsal muscles (Fig. 100 *ms*), form the thick mass of flesh on the back. The lower (ventral) muscles, or the original muscles of the belly, form the fleshy wall of the abdomen. Both sets are segmented, and consist of a double row of muscular plates (Figs. 98, 99, 100), the number of these myotomes determines the number of joints in the trunk, or metamera. The myotomes are also developed from the thick wall of the colon-pouches (Fig. 100 *h*).

Outside this muscular tube we have the external envelope of the vertebrate body, which is known as the corium or cutis. This strong and thick envelope consists, in its deeper strata, chiefly of fat and loose connective tissue, and in its upper layers of cutaneous muscles and firmer connective tissue. It covers the whole surface of the fleshy body, and is of considerable thickness in all the craniota. But in the scranium the corium is merely a thin plate of connective tissue, an insignificant "corium-plate" (*lamella coria*, Figs. 98-100 *i*).

Immediately above the corium is the outer skin (*epidermis*, *e*), the general covering of the whole outer surface. In the higher vertebrates the hairs, nails, feathers, claws, scales, etc., grow out of the epidermis. It consists, with all its appendages and products, of simple cells, and has no blood-vessels. Its cells are connected with the terminations of the sensory nerves. Originally, the outer skin is a perfectly simple covering of the outer surface of the body, composed only of homogeneous cells—a permanent horn-plate. In this simplest form, as a one-layered epithellium, we find it, at first, in all the vertebrates, and throughout life in the scranium. It afterwards grows thicker in the higher vertebrates, and divides into two strata—an outer, firmer corneous (horn) layer and an inner, softer mucous layer; also a number of external and internal appendages grow out of it: outwards, the hairs, nails, claws, etc., and

inwardly, the sweat-glands, fat-glands, etc.

It is probable that in our primitive vertebrate the skin was raised in the middle line of the body in the shape of a vertical fin border (*f*). A similar fringe, going round the greater part of the body, is found to-day in the amphioxus and the cyclostoma; we also find one in the tail of fish-larvæ and tadpoles.

Now that we have considered the external parts of the vertebrate and the animal organs, which mainly lie in the dorsal half, above the chorda, we turn to the vegetal organs, which lie for the most part in the ventral half, below the axial rod. Here we find a large body-cavity or visceral cavity in all the craniotom. The spacious cavity that encloses the greater part of the viscera corresponds to only a part of the original coeloma, which we considered in the tenth chapter; hence it may be called the *metacoeloma*. As a rule, it is still briefly called the coeloma; formerly it was known in anatomy as the pleuropersonal cavity. In man and the other mammals (but only in these) this coeloma divides, when fully developed, into two different cavities, which are separated by a transverse partition—the muscular diaphragm. The fore or pectoral cavity (pleura-cavity) contains the oesophagus (gullet), heart, and lungs; the hind or peritoneal or abdominal cavity contains the stomach, small and large intestine, liver, pancreas, kidneys, etc. But in the vertebrate embryo, before the diaphragm is developed, the two cavities form a single continuous body-cavity, and we find it thus in all the lower vertebrates throughout life. This body-cavity is clothed with a delicate layer of cells, the coelom-epithelium. In the acornia the coelom is segmented both dorsally and ventrally, as their muscular pouches and primitive genital organs plainly show (Fig. 100).

The chief of the viscera in the body-cavity is the alimentary canal, the organ that represents the whole body in the gastrula. In all the vertebrates it is a long tube, enclosed in the body-cavity and more or less differentiated in length, and has two apertures—a mouth for taking in food (Figs. 98, 100 *md*) and an anus for the ejection of unusable matter or excrementa (*af*). With the alimentary canal a number of glands are connected which are of great importance for the vertebrate body, and which all grow out of the

canal. Glands of this kind are the salivary glands, the lungs, the liver, and many smaller glands. Nearly all these glands are wanting in the acornia; probably there were merely a couple of simple hepatic tubes (Figs. 98, 100 *f*) in the vertebrate stem-form. The wall of the alimentary canal and all its appendages consists of two different layers; the inner, cellular clothing is the gut-gland-layer, and the outer, fibrous envelope consists of the gut-fibre-layer; it is mainly composed of muscular fibres which accomplish the digestive movements of the canal, and of connective-tissue fibres that form a firm envelope. We have a continuation of it in the mesentery, a thin, bundle-like layer, by means of which the alimentary canal is fastened to the ventral side of the chorda, originally the dorsal partition of the two coelom-pouches. The alimentary canal is variously modified in the vertebrates both as a whole and in its several sections, though the original structure is always the same, and is very simple. As a rule, it is longer (often several times longer) than the body, and (therefore folded and winding within the body-cavity, especially at the lower end. In man and the higher vertebrates it is divided into several sections, often separated by valves—the mouth, pharynx, oesophagus, stomach, small and large intestine, and rectum. All these parts develop from a very simple structure, which originally (throughout life in the amphioxus) runs from end to end under the chorda in the shape of a straight cylindrical canal.

As the alimentary canal may be regarded morphologically as the oldest and most important organ in the body, it is interesting to understand its essential features in the vertebrate more fully, and distinguish them from unessential features. In this connection we must particularly note that the alimentary canal of every vertebrate shows a very characteristic division into two sections—a fore and a hind chamber. The fore chamber is the head-gut or branchial gut (Figs. 98–100 *p. f*), and is chiefly occupied with respiration. The hind section is the trunk-gut or hepatic gut, which accomplishes digestion (*see, d*). In all vertebrates there are formed, at an early stage, to the right and left in the fore-part of the head-gut, certain special clefts that have an intimate connection with the original respiratory apparatus of

the vertebrate—the branchial (gill) clefts (*br*). All the lower vertebrates, the lancelets, lampreys, and fishes, are constantly taking in water at the mouth, and letting it out again by the lateral clefts of the gullet. This water serves for breathing. The oxygen contained in it is inspired by the blood-canals, which spread out on the parts between the gill-clefts, the gill-arches (*gr*). These very characteristic branchial clefts and arches are found in the embryo of man and all the higher vertebrates at an early stage of development, just as we find them throughout life in the lower vertebrates. However, these clefts and arches never act as respiratory organs in the mammals, birds, and reptiles, but gradually develop into quite different parts. Still, the fact that they are found at first in the same form as in the fishes is one of the most interesting proofs of the descent of these three higher classes from the fishes.

Not less interesting and important is an organ that develops from the ventral wall in all vertebrates—the gill-groove or hypobranchial groove. In the acrania and the ascidus it consists throughout life of a glandular ciliated groove, which runs down from the mouth in the ventral middle line of the gill-gut, and takes small particles of food to the stomach (Fig. 101 *a*). But in the craniota the thyroid gland (*thyroides*) is developed from it, the gland that lies in front of the larynx, and which, when pathologically enlarged, forms goitre (*strueme*).

From the head-gut we get not only the gills, the organs of water-breathing in the lower vertebrates, but also the lungs, the organs of atmospheric breathing in the five higher classes. In these cases a vesicular fold appears in the gullet of the embryo at an early stage, and gradually takes the shape of two spacious sacs, which are afterwards filled with air. These sacs are the two air-breathing lungs, which take the place of the water-breathing gills. But the vesicular invagination, from which the lungs arise, is merely the familiar air-filled vesicle, which we call the floating-bladder of the fish, and which alters its specific weight, acting as hydrostatic organ or floating apparatus. This structure is not found in the lowest vertebrate classes—the acrania and cyclostoma. We shall see more of it in Vol. II.

The second chief section of the vertebrate-gut, the trunk or liver-gut, which

accomplishes digestion, is of very simple construction in the acrania. It consists of two different chambers. The first chamber, immediately behind the gill-gut, is the expanded stomach (*st*); the second, narrower and longer chamber, is the straight small intestine (*d*): it issues behind on the ventral side by the anus (*af*). Near the limit of the two chambers in the visceral cavity we find the liver in the shape of a simple tube or blind sac (*l*); in the amphioxus it is single; in the prosopodius it was probably double (Figs. 98, 100 *f*).

Closely related morphologically and physiologically to the alimentary canal is the vascular system of the vertebrate, the chief sections of which develop from the fibrous gut-layer. It consists of two different but directly connected parts, the system of blood-vessels and that of lymph-vessels. In the passages of the one we find red blood, and in the other colourless lymph. To the lymphatic system belong, first of all, the lymphatic canals proper or absorbent veins, which are distributed among all the organs, and absorb the used-up juices from the tissues, and conduct them into the venous blood; but besides these there are the chyle-vessels, which absorb the white chyle, the milky fluid prepared by the alimentary canal from the food, and conduct this also to the blood.

The blood-vessel system of the vertebrate has a very elaborate construction, but seems to have had a very simple form in the primitive vertebrate, as we find it to-day particularly in the annelids (for instance, earth-worms) and the amphioxus. We accordingly distinguish first of all as essential, original parts of it two large single blood-canals, which lie in the fibrous wall of the gut, and run along the alimentary canal in the median plane of the body, one above and the other underneath the canal. These principal canals give out numerous branches to all parts of the body, and pass into each other by arches, before and behind; we will call them the primitive artery and the primitive vein. The first corresponds to the dorsal vessel, the second to the ventral vessel, of the worms. The primitive or principal artery, usually called the aorta (Fig. 98 *a*), lies above the gut in the middle line of its dorsal side, and conducts oxidized or arterial blood from the gills to the body. The primitive or principal vein (Fig. 100 *e*) lies below the

gut, in the middle line of its ventral side, and is therefore also called the *vena sub-intestinalis*; it conducts carbonized or venous blood back from the body to the gills. At the branchial section of the gut in front the two canals are connected by a number of branches, which rise in arches between the gill-slits. These "branchial vascular arches" (*ky*) run along the gill-arches, and have a direct share in the work of respiration. The anterior continuation of the principal vein which runs on the ventral wall of the gill-gut, and gives off these vascular arches upwards, is the branchial artery (*ky*). At the border of the two sections of the ventral vessel it enlarges into a contractile spindle-shaped tube (Figs 98, 100 A). This is the first outline of the heart, which afterwards becomes a four-chambered pump in the higher vertebrates and man. There is no heart in the amphioxus, probably owing to degeneration. In *protophyllus* the ventral gill-heart probably had the simple form in which we still find it in the *ascidia* and the embryo of the *craniota* (Figs 98, 100 B).

The kidneys, which act as organs of excretion or urinary organs in all vertebrates, have a very different and elaborate construction in the various sections of this stem; we will consider them further in the twenty-ninth chapter. Here I need only mention that in our hypothetical primitive vertebrate they probably had the same form as in the actual amphioxus—the primitive kidneys (*protonephridia*). These are originally made up of a double row of little canals, which directly convey the used-up juices or the urine out of the body-cavity (Fig. 102 a). The inner aperture of these protonephridial canals opens with a ciliated funnel into the body-cavity, the external aperture opens in lateral grooves of the epidermis, a couple of longitudinal grooves in the lateral surface of the outer skin (Fig. 102 b). The protonephridial duct is formed by the closing of this groove to the right and left at the sides. In all the *craniota* it develops at an early stage in the horny plate; in the amphioxus it seems to be converted into a wide cavity, the atrium, or peribranchial space.

Next to the kidneys we have the sexual organs of the vertebrates. In most of the members of this stem the two are united in a single urogenital system; it is only in a few groups that the urinary and sexual organs are separated (in the

amphioxus, the cyclostoma, and some sections of the fish-class). In man and all the higher vertebrates the sexual apparatus is made up of various parts, which we will consider in the twenty-ninth chapter. But in the two lowest classes of our stem, the *ascidia* and *cyclostoma*, they consist merely of simple sexual glands or gonads, the ovaries of the female sex and the testicles (*spermatia*) of the male; the former provide the ova, the latter the sperm. In the *craniota* we always find only one pair of gonads; in the amphioxus several pairs, arranged in succession. They must have had the same form in our hypothetical *protophyllus* (Figs 98, 100 A). These segmental pairs of gonads are the original ventral halves of the colon-pouches.

The organs which we have now enumerated in this general survey, and of which we have noted the characteristic disposition, are those parts of the organism that are found in all vertebrates without exception in the same relation to each other, however much they may be modified. We have chiefly had in view the transverse section of the body (Figs. 101, 102), because in this we see most clearly the distinctive arrangement of them. But to complete our picture we must also consider the segmentation or metamereformation of them, which has yet been hardly noticed, and which is seen best in the longitudinal section. In man and all the more advanced vertebrates the body is made up of a series or chain of similar members, which succeed each other in the long axis of the body—the segments or metamers of the organism. In man these homogeneous parts number thirty-three in the trunk, but they run to several hundred in many of the vertebrates (such as serpents or eels). As this internal articulation or metamereformation is mainly found in the vertebral column and the surrounding muscles, the sections or metamers were formerly called *pro-vertebrae*. As a fact, the articulation is by no means chiefly determined and caused by the skeleton, but by the muscular system and the segmental arrangement of the kidneys and gonads. However, the composition from these *pro-vertebrae* or internal metamers is usually, and rightly, put forward as a prominent character of the vertebrate, and the manifold division or differentiation of them is of great importance in the various groups of the vertebrates. But as far as our present

task—the derivation of the simple body of the primitive vertebrate from the chordula—is concerned, the articulation parts or metamers, are of secondary interest, and we need not go into them just now.

The characteristic composition of the vertebrate body develops from the embryonic structure in the same way in man,

that this answer is just as certain and precise in the case of the origin of man from the mammals. This advanced vertebrate class is also monophyletic, or has evolved from one common stem-group of lower vertebrates (reptiles, and, earlier still, amphibia). This follows from the fact that the mammals are



FIGS. A, B, C, D.—Instances of rudimentary mammary glands and nipples (*hypermastia*). A a pair of small rudimentary breasts (with two nipples on the left) above the large normal ones; from a 45-year-old Berlin woman, who had had children 17 times (twice since). (From *Hilsmann*.) B the highest member has nipples (all young males, three pairs above, one pair below, the large normal breasts), from a 45-year-old normal at Wenzlau. (From *Nagelschke*.) C three pairs of nipples, two pairs on the normal glands and one pair above, from a 35-year-old Japanese girl. D four pairs of nipples, one pair above the normal and two pairs of small rudimentary nipples underneath, from a 40-year-old Bavarian soldier. (From *Nagelschke*.)

as in all the other vertebrates. As all competent experts now admit the monophyletic origin of the vertebrates on the strength of this significant agreement, and this "common descent of all the vertebrates from one original stem-form" is admitted as an historical fact, we have found the answer to "the question of question." We may, moreover, point out

clearly distinguished from the other classes of the stem, not merely in one striking particular, but in a whole group of distinctive characters.

It is only in the mammals that we find the skin covered with hair, the breast-cavity separated from the abdominal cavity by a complete diaphragm, and the larynx provided with an epiglottis. The

mammals alone have three small auscultatory bones in the tympanic cavity—a feature that is connected with the characteristic modification of their maxillary joint. Their red blood-cells have no nucleus, whereas this is retained in all other vertebrates. Finally, it is only in the mammals that we find the remarkable function of the breast structure which has given its name to the whole class—the feeding of the young by the mother's milk. The mammary glands which serve this purpose are interesting in so many ways that we may devote a few lines to them here.

As is well known, the lower mammals, especially those which beget a number of young at a time, have several mammary glands at the breast. Hedgehogs and cows have five pairs, mice four or five pairs, dogs and squirrels four pairs, cats and bears three pairs, most of the ruminants and many of the rodents two pairs, each provided with a teat or nipple (*maria*). In the various genera of the half-apes, (lemurs) the number varies a good deal. On the other hand, the bats and apes, which only beget one young at a time as a rule, have only one pair of mammary glands, and these are found at the breast, as in man.

These variations in the number or structure of the mammary apparatus (*mammarium*) have become doubly interesting in the light of recent research in comparative anatomy. It has been shown that in man and the apes we often find redundant mammary glands (*hyper-marium*) and corresponding teats (*hyper-thelium*) in both sexes. Fig. 103 shows four cases of this kind—A, B, and C of three women, and D of a man. They prove that all the above-mentioned numbers may be found occasionally in man. Fig. 103 A shows the breast of a Berlin woman who had had children seventeen times, and who has a pair of small accessory breasts (with two nipples on the left one) above the two normal breasts; this is a common occurrence, and the small soft pad above the breast is not infrequently represented in ancient statues of Venus. In Fig. 103 C we have the same phenomenon in a Japanese girl of nineteen, who has two nipples on each breast besides (three pairs altogether). Fig. 103 D is a man of twenty-two with four pairs of nipples (as in the dog), a small pair above and two small pairs beneath the large normal teats. The

maximum number of five pairs (as in the sow and hedgehog) was found in a Polish servant of twenty-two who had had several children; milk was given by each nipple; there were three pairs of redundant nipples above and one pair underneath the normal and very large breasts (Fig. 103 B).

A number of recent investigations (especially among recruits) have shown that these things are not uncommon in the male as well as the female sex. They can only be explained by evolution, which attributes them to atavism and latent heredity. The earlier ancestors of all the primates (including man) were lower placentals, which had, like the hedgehog (one of the oldest forms of the living placentals), several mammary glands (five or more pairs) in the abdominal skin. In the apes and man only a couple of them are normally developed, but from time to time we get a development of the atrophied structures. Special notice should be taken of the arrangement of these accessory mammae; they form, as is clearly seen in Fig. 103 B and D, two long rows, which diverge forward (towards the arm-pit), and converge behind in the middle line (towards the clitoris). The milk-glands of the polymastic lower placentals are arranged in similar lines.

The phylogenetic explanation of polymastism, as given in comparative anatomy, has lately found considerable support in ontogeny. Hans Strahl, E. Schmitt, and others, have found that there are always in the human embryo at the sixth week (when it is three-fifths of an inch long) the microscopic traces of five pairs of mammary glands, and that they are arranged at regular distances in two lateral and divergent lines, which correspond to the mammary lines. Only one pair of them—the central pair—are normally developed, the others atrophying. Hence there is for a time in the human embryo a normal hyperthelism, and this can only be explained by the descent of man from lower primates (lemurs) with several pairs.

But the milk-gland of the mammal has a great morphological interest from another point of view. This organ for feeding the young in man and the higher mammals is, as is known, found in both sexes. However, it is usually active only in the female sex, and yields the valuable "mother's milk"; in the male sex it is

small and inactive, a real rudimentary organ of no physiological interest. Nevertheless, in certain cases we find the breast as fully developed in man as in woman, and it may give milk for feeding the young.

We have a striking instance of this *gynecomastia* (large milk-giving breasts in a male) in Fig. 104. I owe the photograph (taken from life) to the kindness of Dr. Grassein, of Athens, a German physician, who has rendered service by a number of anthropological observations (for instance, in several cases of failed

my stay in Ceylon (at Belligemma) in 1881. A young Cinghalese in his twenty-fifth year was brought to me as a curious hermaphrodite, half-man and half-woman. His large breasts gave plenty of milk; he was employed as "male nurse" to suckle a new-born infant whose mother had died at birth. The outline of his body was softer and more feminine than in the Greek shown in Fig. 104. As the Cinghalese are small of stature and of graceful build, and as the men often resemble the women in clothing (upper part of the body naked, female dress on the lower




FIG. 104.—A Greek gynecomast.

men). The gynecomast in question is a Greek recruit in his twentieth year, who has both normally developed male organs and very pronounced female breasts. It is noteworthy that the other features of his structure are in accord with the softer form of the female sex. It reminds us of the marble statues of hermaphrodites which the ancient Greek and Roman sculptors often produced. But the man would only be a real hermaphrodite if he had ovaries internally besides the (externally visible) testicles.

I observed a very similar case during

part) and the dressing of the hair (with a comb), I first took the beardless youth to be a woman. The illusion was greater, as in this remarkable case gynecomastia was associated with *cryptorchism*—that is to say, the testicles had kept to their original place in the visceral cavity, and had not travelled in the normal way down into the scrotum. (Cf. Chapter XXIX.) Hence the latter was very small, soft, and empty. Moreover, one could feel nothing of the testicles in the inguinal canal. On the other hand, the male organ was very small, but normally developed. It was

clear that this apparent hermaphrodite also was a real male.

Another case of practical gynecomastia has been described by Alexander von Humboldt. In a South American forest he found a solitary settler whose wife had died in child-birth. The man had laid the new-born child on his own breast in despair; and the continuous stimulus of the child's sucking movements had revived the activity of the mammary glands. It is possible that nervous suggestion had some share in it. Similar cases have been often observed in recent years, even among other male mammals (such as sheep and goats).

The great scientific interest of these facts is in their bearing on the question of heredity. The stem-history of the mammarium rests partly on its embryology (Chapter XXIV.) and partly on the facts of comparative anatomy and physio-

logy. As in the lower and higher mammals (the monotremes, and most of the marsupials) the whole lactiferous apparatus is only found in the female; and as there are traces of it in the male only in a few younger marsupials, there can be no doubt that these important organs were originally found only in the female mammal, and that they were acquired by these through a special adaptation to habits of life.

Later, these female organs were communicated to both sexes by heredity; and they have been maintained in all persons of either sex, although they are not physiologically active in the males. This normal permanence of the female lactiferous organs in both sexes of the higher mammals and man is independent of any selection, and is a fine instance of the much-disputed "inheritance of acquired characters."

CHAPTER XII.

EMBRYONIC SHIELD AND GERMINATIVE AREA

THE three higher classes of vertebrates which we call the amniotes—the mammals, birds, and reptiles—are notably distinguished by a number of peculiarities of their development from the five lower classes of the stem—the animals without an amnion (the anamniotes). All the amniotes have a distinctive embryonic membrane known as the amnion (or "water-membrane"), and a special embryonic appendage—the allantois. They have, further, a large yolk-sac, which is filled with food-yolk in the reptiles and birds, and with a corresponding clear fluid in the mammals. In consequence of these later-acquired structures, the original features of the development of the amniotes are so much altered that it is very difficult to reduce them to the pallogenic embryonic processes of the lower amnion-less vertebrates. The gastræa theory shows us how to do this, by representing the embryology of the lowest vertebrate, the skull-less amphioxus, as

the original form, and deducing from it, through a series of gradual modifications, the gastrulation and celomization of the craniota.

It was somewhat fatal to the true conception of the chief embryonic processes of the vertebrates that all the older embryologists, from Malpighi (1667) and Wolff (1750) to Baer (1828) and Remak (1850), always started from the investigation of the hen's egg, and transferred to man and the other vertebrates the impressions they gathered from this. This classical object of embryological research is, as we have seen, a source of dangerous errors. The large round food-yolk of the bird's egg causes, in the first place, a flat discoid expansion of the small gastrula, and then so distinctive a development of this thin round embryonic disk that the controversy as to its significance occupies a large part of embryological literature.

One of the most unfortunate errors that this led to was the idea of an original

antithesis of germ and yolk. The latter was regarded as a foreign body, extrinsic to the real germ, whereas it is properly a part of it, an embryonic organ of nutrition. Many authors said there was no trace of the embryo until a later stage, and outside the yolk; sometimes the two-layered embryonic disk itself, at other times only the central portion of it (as distinguished from the germinative area, which we will describe presently), was taken to be the first outline of the embryo.

primitive gut. This is clearly shown by the ova of the amphibia and cyclostoma, which explain the transition from the yolk-less ova of the amphioxus to the large yolk-filled ova of the reptiles and birds.

It is precisely in the study of these difficult features that we see the incalculable value of phylogenetic considerations in explaining complex ontogenetic facts, and the need of separating ontogenetic phenomena from pangenetic.




FIG. 109.—Development of the discoid mammalian embryo from the yolk-mass, in transverse section (Mammalian). A The germinal disk (*a*, *b*) has set on one side of the branching vein (*ab*). B In the middle of the germinal disk we find the medullary groove (*mr*), and underneath it the chorda (*ch*). C The gut-bladder (*g*) has been enclosed by the gut-gland-layer (*gd*). D The skin-fibro-layer (*sf*) and gut-fibro-layer (*gf*) divide in the periphery, the gut (*g*) begins to separate from the yello-mass or umbilical vessels (*uv*). E The medullary tube (*mr*) is closed, the body-cavity (*c*) begins to form. F The pre-embryo (*pe*) begins to grow round the medullary tube (*mr*) and the chorda (*ch*); the gut (*g*) is cut off from the umbilical vessels (*uv*). G The umbilical vessels (*uv*) have grown round the medullary tube (*mr*) and chorda, the body-cavity is closed, and the umbilical vessels have disappeared. The ectoderm and mesoderm membranes are visible. H The embryo has the same structure throughout: *a* bone-plate, *mr* medullary tube, *sf* skin-fibro-layer, *w* pre-embryo, *ch* chorda, *c* body-cavity or coelom, *gf* gut-fibro-layer, *gd* gut-gland-layer, *g* gut-cavity, *uv* umbilical vessels.

The embryo has the same structure throughout: *a* bone-plate, *mr* medullary tube, *sf* skin-fibro-layer, *w* pre-embryo, *ch* chorda, *c* body-cavity or coelom, *gf* gut-fibro-layer, *gd* gut-gland-layer, *g* gut-cavity, *uv* umbilical vessels.

In the light of the gastraea theory it is hardly necessary to dwell on the defects of this earlier view and the erroneous conclusions drawn from it. In reality, the first segmentation-cell, and even the stem-cell itself and all that issues therefrom, belong to the embryo. As the large original yolk-mass in the undivided egg of the bird only represents an inclosure in the greatly enlarged ovum, so the later contents of its embryonic yolk-mass (whether yet segmented or not) are only a part of the ectoderm which forms the

This is particularly clear as regards the comparative embryology of the vertebrates, because here the phylogenetic unity of the stem has been already established by the well-known facts of paleontology and comparative anatomy. If this unity of the stem, on the basis of the amphioxus, were always borne in mind, we should not have these errors constantly recurring.

In many cases the ontogenetic relation of the embryo to the food-yolk has until now given rise to a quite wrong idea of

the first and most important embryonic processes in the higher vertebrates, and has occasioned a number of false theories in connection with them. Until thirty years ago the embryology of the higher vertebrates always started from the position that the first structure of the embryo is a flat, leaf-shaped disk; it was for this reason that the cell-layers that compose this germinal disk (also called *germinative area*) are called "germinal layers." This flat germinal disk, which is round at first and then oval, and which is often described as the tread or cicatrice in the laid hen's egg, is found at a certain part of the surface of the large globular food-yolk. I am convinced that it is nothing else than the discoid, flattened gastrula of the birds. At the beginning of germination the flat embryonic disk curves outwards, and separates on the inner side from the underlying large yolk-ball. In this way the flat layers are converted into tubes, their edges folding and joining together (Fig. 105). As the embryo grows at the expense of the food-yolk, the latter becomes smaller and smaller, it is completely surrounded by the germinal layers. Later still, this remainder of the food-yolk only forms a small round sac, (the yolk-sac or umbilical vesicle (Fig. 105 ab). This is enclosed by the visceral layer, is connected by a thin stalk, the yolk-duct, with the central part of the gut-tube, and is finally, in most of the vertebrates, entirely absorbed by this (H). The point at which this takes place, and where the gut finally closes, is the vascular navel. In the mammals, in which the remainder of the yolk-sac remains without and atrophies, the yolk-duct at length penetrates the outer ventral wall. At birth the umbilical cord proceeds from here, and the point of closure remains throughout life in the skin as the navel.

As the older embryology of the higher vertebrates was mainly based on the chick, and regarded the antithesis of embryo (or formative-yolk) and food-yolk (or yolk-sac) as original, it had also to look upon the flat leaf-shaped structure of the germinal disk as the primitive embryonic form, and emphasize the fact that hollow grooves were formed of these flat layers by folding, and closed tubes by the joining together of their edges.

This idea, which dominated the whole treatment of the embryology of the higher vertebrates until thirty years ago, was

totally false. The gastrula theory, which has its chief application here, teaches us that it is the very reverse of the truth. The cup-shaped gastrula, in the body-wall of which the two primary germinal layers appear from the first as closed tubes, is the original embryonic form of all the vertebrates, and all the multicellular invertebrates; and the flat germinal disk with its superficially expanded germinal layers is a later, secondary form, due to the cenogenetic formation of the large food-yolk and the gradual spread of the germ-layers over its surface. Hence the actual folding of the germinal layers and their conversion into tubes is not an original and primary, but a much later and tertiary, evolutionary process. In the phylogeny of the vertebrate embryonic process we may distinguish the following three stages.—

A. First Stage Primary (embryonic) embryo- proton.	B. Second Stage Secondary (cenogenetic) embryo- proton.	C. Third Stage Tertiary (metagenetic) embryo- proton.
The germinal layer forms from the first closed tubes, the embryonic blastula being converted into the two-layered gastrula by invagination. No food-yolk (<i>Amphioxus</i> .)	The germinal layers spread out leaf-shaped over the food-yolk, gathering in the ventral midline, and a large yolk-sac being formed from the middle of the gut-tube. (<i>Amphioxus</i> .)	The germinal layer forms a flat germinal disk, the borders of which join to gather and form closed tubes separating from the central yolk-sac. (<i>Ammonia</i> .)

As this theory, a logical conclusion from the gastrula theory, has been fully substantiated by the comparative study of gastrulation in the last few decades, we must exactly reverse the hitherto prevalent mode of treatment. The yolk-sac is not to be treated, as was done formerly, as if it were originally antithetic to the embryo, but as an essential part of it, a part of its visceral tube. The primitive gut of the gastrula line, on this view, been divided into two parts in the higher animals as a result of the cenogenetic formation of the food-yolk—the permanent gut (*metagastron*), or permanent alimentary canal, and the yolk-sac (*lecithosoma*), or umbilical vesicle. This is very clearly shown by the comparative ontogeny of the fishes and amphibia. In these cases the whole yolk undergoes cleavage at first, and forms a yolk-gland, composed of yolk-cells, in the ventral wall

of the primitive gut. But it afterwards becomes so large that a part of the yolk does not divide, and is used up in the yolk-sac that is cut off outside.

When we make a comparative study of the embryology of the amphioxus, the frog, the chick, and the rabbit, there cannot, in my opinion, be any further doubt as to the truth of this position, which I have held for thirty years. Hence in the light of the gastrula theory we must regard the features of the amphioxus as the only and real primitive structure among all the vertebrates, departing very little from the palæogenetic embryonic form. In the cyclostoma and the frog these features are, on the whole, not much altered cenogenetically, but

The oldest, oviparous mammals, the monotremes, behave in the same way as the reptiles and birds. But the corresponding embryonic processes in the viviparous mammals, the marsupials and placentals, are very elaborate and distinctive. They were formerly quite misinterpreted; it was not until the publication of the studies of Edward van Beneden (1875) and the later research of Selenka, Kupffer, Rabi, and others, that light was thrown on them, and we were in a position to bring them into line with the principles of the gastrula theory and trace them to the embryonic forms of the lower vertebrates. Although there is no independent food-yolk, apart from the formative yolk, in the mammal ovum, and although its augmentation is total on this account, nevertheless a large yolk-sac is formed in their embryos, and the "embryo proper" spreads leaf-wise over its surface, as in the reptiles and birds, which have a large food-yolk and partial segmentation. In the mammals, as well as in the latter, the flat, leaf-shaped germinal disk separates from the yolk-sac, and its edges join together and form tubes.




FIG. 106.




FIG. 107.

FIG. 106.—The visceral embryonic vesicle (*blastogastrula* or *gastrula*) of a rabbit (the "blastula" or *vesicula blastodermica* of other writers: a center vesicle is retained). 2 also-layer of ectoderm forming the entire wall of the yolk-sac, a group of dark cells representing the visceral layer or endoderm.

FIG. 107.—The same in section. Letters as above. *a* ectoderm, *b* endoderm.

they are very much so in the chick, and most of all in the rabbit. In the blastogastrula of the amphioxus and in the hooded gastrula of the lamprey and the frog the germinal layers are found to be closed tubes or vesicles from the first. On the other hand, the chick-embryo (in the new laid, but not yet hatched, egg) is a flat circular disk, and it was not easy to recognize this as a real gastrula. Rauber and Goette have, however, achieved this. As the discoid gastrula grows round the large globular yolk, and the permanent gut then separates from the outlying yolk-sac, we find all the processes which we have shown (diagrammatically) in Fig. 106—processes that were hitherto regarded as principal ones, whereas they are merely secondary.

How can we explain this curious anomaly? Only as a result of very characteristic and peculiar cenogenetic modifications of the embryonic process, the real causes of which must be sought in the change in the rearing of the young on the part of the viviparous mammals. These are clearly connected with the fact that the ancestors of the viviparous mammals were oviparous ancestors like the present monotremes, and only gradually became viviparous. This can no longer be questioned now that it has been shown (1884) that the monotremes, the lowest and oldest of the mammals, still lay eggs, and that these develop like the ova of the reptiles and birds. Their nearest descendants, the marsupials, formed the habit of retaining the eggs, and developing them in the

oviduct; the latter was thus converted into a womb (uterus). A nutritive fluid that was secreted from its wall, and passed through the wall of the blastula, now served to feed the embryo, and took the place of the food-yolk. In this way the original food-yolk of the monotremes gradually atrophied, and at last disappeared so completely that the partial ovum-segmentation of their descendants, the rest of the mammals, once more became total. From the therapsentalis of the former was evolved the distinctive *epiblastula* of the latter.

It is only by this phylogenetic explanation that we can understand the formation and development of the peculiar, and hitherto totally misunderstood, blastula of the mammal. The vesicular condition of the mammalian embryo was discovered 200 years ago (1677) by *Ragner de Graaf*. He found in the uterus of a rabbit four days after insemination small, round, loose, transparent vesicles, with a double envelope. However, *Graaf's* discovery passed without recognition. It was not until 1827 that these vesicles were rediscovered by *Baer*, and then more closely studied in 1842 by *Bischoff* in the rabbit (Figs. 106, 107). They are found in the womb of the rabbit, the dog, and other small mammals, a few days after copulation. The mature ova of the mammal, when they have left the ovary, are fertilized either here or in the oviduct immediately afterwards by the invading sperm-cells. (As to the womb and oviduct see Chapter XXIX.) The cleavage and formation of the gastrula take place in the oviduct. Either here in the oviduct or after the mammalian gastrula has passed into the uterus it is converted into the globular vesicle which is shown externally in Fig. 106, and in section in Fig. 107. The thick, outer, structureless envelope that encloses it is the original *coelomema* or *zona pellucida*, modified, and clothed with a layer of albumin that has been deposited on the outside. From this stage the envelope is called the external membrane, the *primary chorion* or *pro-chorion* (a). The real wall of the vesicle

enclosed by it consists of a simple layer of entodermic cells (b), which are fastened by mutual pressure, and generally hexagonal; a light nucleus shines through their fine-grained protoplasm (Fig. 106). At one part (c) inside this hollow ball we find a circular disk, formed of darker, softer, and rounder cells, the dark-grained entodermic cells (Fig. 106).

The characteristic embryonic form that the developing mammal now exhibits has up to the present usually been called the "blastula" (*Bischoff*), "one-aimed embryo" (*Baer*), "vesicular embryo" (*vertebrate blastodermica*, or, briefly, *blastopora*). The wall of the hollow vesicle, which consists of a single layer of cells, was called the "blastoderm," and was supposed to be equivalent to the cell-layer of the same name that forms the wall of the real blastula of the amphioxus and



FIG. 106.

FIG. 107.

FIG. 106.—Four entodermic cells from the embryonic vesicle of the rabbit.

FIG. 107.—Two blastulae from the embryonic vesicle of the rabbit.

many of the invertebrates (such as *Monaxozoa*, Fig. 29 F. G). Formerly this real blastula was generally believed to be equivalent to the embryonic vesicle of the mammal. However, this is by no means the case. What is called the "blastula" of the mammal and the real blastula of the amphioxus and many of the invertebrates are totally different embryonic structures. The latter (blastula) is palliogenic, and precedes the formation of the gastrula. The former (blastodermic vesicle) is oögenetic, and follows gastrulation. The globular wall of the blastula is a real blastoderm, and consists of homogeneous (blastodermic) cells; it is not yet differentiated into the two primary germinal layers. But the globular wall of the mammalian vesicle is the differentiated ectoderm, and at one point in it we find a circular disk of quite different cells—the entoderm. The round

* In this and the other mammals the fertilization of the ova probably takes place, as a rule, in the oviduct, here the ova, which come from the female ovary in the shape of the *Graafian* follicle, and enter the lower aperture of the oviduct, encounter the mobile sperm-cells of the male seed, which pass into the uterus of copulation, and from there into the external aperture of the oviduct. Insemination rarely takes place in the ovary or in the vagina.

cavity, filled with fluid, inside the real blastula is the segmentation-cavity. But the similar cavity within the mammal vesicle is the yolk-sac cavity, which is

which we have considered previously (Chapter IX.). For these reasons it is very necessary to recognise the secondary embryonic vesicle in the mammal (*gastrula*



FIG. 110.




FIG. 111.



FIG. 112.




FIG. 113.

FIG. 110.—Ovum of a rabbit from the uterus, one week of an inch in diameter. The embryonic vesicle (*b*) has withdrawn a little from the nucleus (*a*). (*c*) is the middle of the cell-membrane, or middle germ-layer (blastodermis) at the edge of which (*d*) the outer layer of the embryonic vesicle is already beginning to expand. (Figs. 110-114 from *Randall*.)

FIG. 111.—The same ovum, same as profile. Later as in Fig. 110.

FIG. 112.—Ovum of a rabbit from the uterus, one fourth of an inch in diameter. The blastoderm is already for the most part two-layered (*b*). The cell-membrane, or outer perhaps, is defined (*c*).

FIG. 113.—The same ovum, same as profile. Later as in Fig. 112.

FIG. 114.—Ovum of a rabbit from the uterus, one third of an inch in diameter. The embryonic vesicle is now nearly everywhere two-layered (*b*), only remaining one-layered below (*a*).



FIG. 114.

connected with the incipient gut-cavity. This primitive gut-cavity passes directly into the segmentation-cavity in the mammal. In consequence of the peculiar ontogenetic changes in their gastrulation,

cycles or blastocycles) as a characteristic structure peculiar to this class, and distinguish it carefully from the primary blastula of the amphioxus and the invertebrates.

The small, circular, whitish, and opaque spot which the gastric disk (Fig. 106) forms at a certain part of the surface of the clear and transparent embryonic vesicle has long been known to science,



FIG. 103.—Broad germinative area of the rabbit, divided into the central light area (inner germinal spot) and the peripheral dark area (outer germinal spot). The light area forms darker on account of the dark ground appearing through it.

and compared to the germinal disk of the birds and reptiles. Sometimes it has been called the germinal disk, sometimes the germinal spot, and usually the germinative area. From the area the further development of the embryo proceeds. However, the larger part of the embryonic vesicle of the mammal is not directly used for building up the later body, but for the construction of the temporary umbilical vesicle. The embryo separates from this in proportion as it grows at its expense; the two are only connected by the yolk-duct (the stalk of the yolk-sack), and thus maintains the direct communication between the cavity of the umbilical vesicle and the forming visceral cavity (Fig. 105).

The germinative area or gastric disk of the mammal consists at first (like the germinal disk of birds and reptiles) merely of the two primary germinal layers, the ectoderm and endoderm. But soon there appears in the middle of the circular disk between the two a third stratum of cells, the rudiment of the middle layer or fibrous layer (*mesoderm*). This middle germinal layer consists from the first, as we have seen in the tenth Chapter, of two separate epithelial plates, the two layers of the coelom-pouches (parietal and visceral). However, in all the amniotes (on account of the large formation of yolk) these thin middle plates are so firmly pressed together that they seem to represent a

single layer. It is thus peculiar to the mammals that the middle of the germinative area is composed of four germinal layers, the two limiting (or primary) layers and the middle layers between them (Figs. 96, 97). These four secondary germinal layers can be clearly distinguished as soon as what is called the saddle-groove (or "embryonic saddle") is seen at the hind border of the germinative area. At the borders, however, the germinative area of the mammal only consists of two layers. The rest of the wall of the embryonic vesicle consists at first (but only for a short time in most of the mammals) of a single layer, the outer germinal layer.

From this stage, however, the whole wall of the embryonic vesicle becomes two-layered. The middle of the germinative area is much thickened by the growth of the cells of the middle layers, and the inner layer expands at the same time, and increases at the border of the disk all round. Lying close on the outer layer throughout, it grows over its inner surface at all points, covers first the upper and then the lower hemisphere, and at last closes in the middle of the inner layer (Figs. 110-114). The wall of the embryonic vesicle now consists throughout of two layers of cells, the ectoderm without and the endoderm within. It is only in the centre of the circular area, which



FIG. 105.—Oral area, with the opaque whitish border of the dark area without.

becomes thicker and thicker through the growth of the middle layers, that it is made up of all four layers. At the same time, small structureless tufts or warts are deposited on the surface of the outer

ovolemma or prochorion, which has been raised above the embryonic vesicle (Figs. 1112-1114 a).

We may now disregard both the outer ovolemma and the greater part of the



FIG. 1113.—Oval germinal disk of the rabbit, magnified about ten times. As the disk is half-transparent disk lies on a black ground, the lighted area looks like a dark ring, and the opaque area (lying outside of it) like a white ring. The oval shield in the center also looks whitish, and in its area we see the dark embryonic groove. (From Burdick.)

vesicle, and concentrate our attention on the germinative area and the four-layered embryonic disk. It is here alone that we find the important changes which lead to the differentiation of the first organs. It is immaterial whether we examine the germinative area of the mammal (the rabbit, for instance) or the germinal disk of a bird or a reptile (such as a lizard or tortoise). The embryonic processes we are now going to consider are essentially the same in all members of the three higher classes of vertebrates which we call the amniotes. Man is found to agree in this respect with the rabbit, dog, ox, etc.; and in all these mammals the germinative area undergoes essentially the same changes as in the birds and reptiles. They are most frequently and accurately studied in the chick, because we can have incubated hens' eggs in any quantity at any stage of development. Moreover, the round germinal disk of the chick passes immediately after the beginning of incubation (within a few hours) from the two-

layered to the four-layered stage, the two-layered mesoderm developing from the median primitive groove between the ectoderm and endoderm (Figs. 82-95).

The first change in the round germinal disk of the chick is that the cells at its edges multiply more briskly, and form darker nuclei in their protoplasm. This gives rise to a dark ring, more or less sharply set off from the lighter center of the germinal disk (Fig. 1115). From this point the latter takes the name of the "light area" (*area pellucida*), and the darker ring is called the "dark area" (*area opaca*). (In a strong light, as in Figs. 1115-1117, the light area seems dark, because the dark ground is seen through it; and the dark area seems whiter). The circular shape of the area now changes into elliptic, and then immediately into oval (Figs. 1116, 1117). One end seems to be broader and blunter, the other narrower and more pointed, the former corresponds to the anterior and the latter to the posterior section of the subsequent body. At the same time, we can already trace the characteristic bilateral form of the body, the antithesis of right and left, before and



FIG. 1116.—Pear-shaped germinal shield of the rabbit (eight days old), magnified twenty times. *a*, embryonic groove, *b*, primitive groove (primitive streak). (From Burdick.)

behind. This will be made clearer by the "primitive streak," which appears at the posterior end.

At an early stage an opaque spot is seen in the middle of the clear germinative

area, and this also passes from a circular to an oval shape. At first this shield-shaped marking is very delicate and barely perceptible; but it soon becomes clearer, and now stands out as an oval shield, surrounded by two rings or areas (Fig. 117). The inner and brighter ring is the remainder of the pallid area, and the dark outer ring the remainder of the opaque area; the opaque shield-like spot itself is the first rudiment of the dorsal part of the embryo. We give it briefly

"shield" and "germinative area" are used in many different senses—and this has led to a fatal confusion in embryonic literature—we must explain very clearly the real significance of these important embryonic parts of the amniote. It will be useful to do so in a series of formal principles:—

1. The so-called "first trace of the embryo" in the amniotes, or the embryonic shield, in the centre of the pallid area, consists merely of an early differen-




FIG. 117.—Median longitudinal section of the gastrula of four vertebrates. (From Rüd.) A, *Gastrula* of a shark (*Alopias*). B, *Gastrula* of a murex (*Littorina*). C, *Gastrula* of an amphioxus (*Tunicata*). D, *Gastrula* of an ascidian (*Stolidia*). A, ventral; B, dorsal; C, dorsal; D, ventral.

the name of embryonic shield or dorsal shield. In most works this embryonic shield is described as "the first rudiment or trace of the embryo," or "primitive embryo." But this is wrong, though it rests on the authority of Baer and Bischoff. As a matter of fact, we already have the embryo in the stem-cell, the gastrula, and all the subsequent stages. The embryonic shield is simply the first rudiment of the dorsal part, which is the earliest telodermis. As the older names of "embryonic rudiment" and "germinative area" are used

in many different senses—and this has led to a fatal confusion in embryonic literature—we must explain very clearly the real significance of these important embryonic parts of the amniote. It will be useful to do so in a series of formal principles:—

2. Hence the best name for it is "the dorsal shield," as I proposed long ago.

3. The germinative area, in which the first embryonic blood-vessels appear at an early stage, is not opposed as an external area to the "embryo proper," but is a part of it.

4. In the same way, the yolk-mass or the umbilical vesicle is not a foreign external

appendage of the embryo, but an outlying part of its primitive gut.

5. The dorsal shield gradually separates from the germinative area, and the yolk-sac, its edges growing downwards, and folding together to form ventral plates.

6. The yolk-sac and vessels of the germinative area, which soon spread over its whole surface, are, therefore, real embryonic organs, or temporary parts of the embryo, and have a transitory importance in connection with the nutrition of the growing later body; the latter may be called the "permanent body" in contrast to them.

The relation of these ontogenetic features of the amniotes to the palaeogenetic structures of the older non-amniotic vertebrates may be expressed in the following theses. The original gastrula, which completely passes into the embryonic body in the acrochord, cyclostomes, and amphibia, is early divided into two parts in the amniotes—the embryonic shield, a fuch represents the dorsal outline of the permanent body, and the temporary embryonic organs of the germinative area and its blood-vessels, which soon grow over the whole of the yolk-sac. The differences which we find in the various classes of the vertebrate stem in these important particulars can only be fully understood when we bear in mind their phylogenetic relations on the one hand, and, on the other, the ontogenetic modifications of structure that have been brought about by changes in the rearing of the young and the variation in the mass of the food-yolk.

We have already described in the ninth chapter the changes which this increase and decrease of the nutritive yolk causes in the form of the gastrula, and especially in the situation and shape of the primitive mouth. The primitive mouth or proboscis is originally a simple round aperture at the lower pole of the long axis, its dorsal lip is above and ventral lip below. In the amphioxus this primitive mouth is a little eccentric, or shifted to the dorsal side (Fig. 39). The aperture increases with the growth of the food-yolk in the cyclostomes and ganoids; in the sturgeon it lies almost on the equator of the round ovum,

the ventral lip (*a*) in front and the dorsal lip (*b*) behind (Fig. 119 *δ*). In the wide-mouthed, circular discoid gastrula of the selachii or primitive fishes, which spreads quite flat on the large food-yolk, the anterior semi-circle of the border of the disk is the ventral, and the posterior semi-circle the dorsal lip (Fig. 119 *A*). The amphiblastic amphibia are directly connected with their earlier fish-ancestors, the dipnoans and ganoids, and further the old-fashioned selachii (*Cestracanth*), they have retained their total unequal segmentation, and their small primitive mouth (Fig. 119 *C*, *δδ*), blocked up by the yolk-stopper, lies at the limit of the dorsal and ventral surface of the embryo (at the lower pole of its equatorial axis), and there again has an upper dorsal and a lower ventral lip (*a*, *b*). The formation of a large food-yolk followed again in the stem-forms of the tunicates, the protobranchs or procephala, descended from the amphibia (Fig. 119 *D*). But here the accumulation of the food-yolk took place only in the ventral wall of the peristome-gut, so that the narrow primitive mouth lying behind was forced upwards, and came to be on the back of the discoid "epigastrula" in the shape of the "primitive groove", thus (in contrast to the case of the selachii, Fig. 119 *A*) the dorsal lip (*b*) had to be in front, and the ventral lip (*a*) behind (Fig. 119 *D*). This feature was transmitted to all the amniotes, whether they retained the large food-yolk (reptiles, birds, and monotremes), or lost it by atrophy (the viviparous mammals).

This phylogenetic explanation of gastrulation and cleftation, and the comparative study of them in the various vertebrates, throw a clear and full light on many ontogenetic phenomena, as to which the most obscure and confused opinions were prevalent thirty years ago. In this we see especially the high scientific value of the biogenetic law and the careful separation of palaeogenetic from ontogenetic processes. To the opponents of this law the real explanation of these remarkable phenomena is impossible. Here, and in every other part of embryology, the true key to the solution lies in phylogeny.

CHAPTER XIII.

DORSAL BODY AND VENTRAL BODY

THE earliest stages of the human embryo are, for the reasons already given, either quite unknown or only imperfectly known to us. But as the subsequent embryonic forms in man behave and develop just as they do in all the other mammals, there cannot be the slightest doubt that the preceding stages also are similar. We have been able to see in the coelomula of the human embryo (Fig. 97), by transverse sections (through its primitive mouth, that its two coelom-pouches are developed in just the same way as in the

it is in the middle line of this that the primitive streak appears (Fig. 121 *pe*). The narrow longitudinal groove in it—the so-called “primitive groove”—is, as we have seen, the primitive mouth of the gastrula. In the gastrula-embryo of the mammals, which are much modified ontogenetically, this cleft-shaped prostoma is lengthened so much that it soon traverses the whole of the hinder half of the dorsal shield; as we find in a rabbit-embryo of six to eight days (Fig. 222 *pr*). The two swollen parallel borders that



FIG. 122.—Embryonic vesicle of a seven-days-old rabbit with oral embryonic shield (*sz*). *A* seen from above, *B* from the back. (From Kükenthal.) *sz*, dorsal shield or embryonic spot. In *A* the upper half of the shield is made up of the two primary germinal layers, the lower (up to *pr*) only from the coelom layer.

rabbit (Fig. 95); moreover, the peculiar course of the gastrulation is just the same.

The germinal area forms in the human embryo in the same way as in the other mammals, and in the middle part of this we have the embryonic shield, the purport of which we considered in the previous chapter. The next changes in the embryonic disk, or the “embryonic spot,” take place in corresponding fashion. These are the changes we are now going to consider more closely.

The chief part of the oral embryonic shield is at first the narrow hinder end;

limit this median furrow are the side lips of the primitive mouth, right and left. In this way the bilateral-symmetrical type of the vertebrate becomes pronounced. The subsequent head of the annulate is developed from the broader and rounder fore-half of the dorsal shield.

In this fore-half of the dorsal shield a median furrow quickly makes its appearance (Fig. 123 *cf*). This is the broader dorsal furrow or medullary groove, the first beginning of the central nervous system. The two parallel dorsal or medullary swellings that enclose it grow

together over it afterwards, and form the medullary tube. As is seen in transverse sections, it is formed only of the outer germinal layer (Figs. 95, 136). The lips of the primitive mouth, however, Ea, as

we know, at the important point where the outer layer bends over the inner, and from which the two coelomic pouches grow between the primary germinal layers.

Thus the median primitive furrow (*pf*)



FIG. 137.—Dorsal aspect of the rabbit. (A of 18 days eighteen hours. B of eight days). (From Kalkbrenner.) *pf* primitive furrow, *pr* primitive groove, *my* myel. groove, *sc* schisto-shaped eye spot, *g* gut.




FIG. 138.



FIG. 139.

FIG. 138.—Dorsal aspect (*pf*) and primitive furrow of a rabbit-embryo of eight days. (From Kalkbrenner.) *pf* primitive furrow, *pr* primitive groove.

FIG. 139.—Ventral aspect of a rabbit of eight days. (From Van Beneden.) *pf* primitive furrow, *sc* schisto-shaped eye spot, *g* gut, *my* myel. groove, *sc* schisto-shaped eye spot, *g* gut.

in the hind-half and the median medullary furrow (*mf*) in the fore-half of the oval shield are totally different structures, although the latter seems to a superficial observer to be merely the forward continuation of the former. Hence they

plainly in the amphioxus in a longitudinal section of the oösomula, as soon as the primitive mouth is completely closed at its hinder end. The medullary tube has still at this stage an opening at the forward end, the neurocrus (Fig. 83 *np*).

This opening also is afterwards closed. There are then two completely closed canals over each other—the medullary tube above and the gastric tube below, the two being separated by the chorda. The same features as in the acrania are exhibited by the related tunicates, the ascidians.

Again, we find the neurenteric canal in just the same form and situation in the amphibia. A longitudinal section of a young tadpole (Fig. 125) shows how we may penetrate from the still open primitive mouth (*x*) either into the wide primitive gut-cavity (*al*) or the narrow overlying nerve-tube. A little later, when the

primitive mouth is closed, the narrow neurenteric canal (Fig. 126 *nc*) represents the arched connection between the dorsal medullary canal (*mc*) and the ventral gastric canal.

In the annelotes this original curved form of the neurenteric canal cannot be found at first, because here the primitive mouth travels completely over to the



FIG. 124.—Longitudinal section of the oösomula of amphioxus (from the left). *a*, notochord; *d*, primitive gut; *ce*, medullary tube; *ne*, nerve tube; *mc*, medullary canal; *st*, first primitive segment; *c*, cellular pouches. (From Haeckel.)

were formerly always confused. This error was the more pardonable as immediately afterwards the two grooves do actually pass into each other in a very remarkable way. The point of transition is the remarkable neurenteric canal (Fig. 124 *nc*). But the direct connection which is thus established does not last long; the two are soon definitely separated by a partition.

The enigmatic *neurenteric canal* is a very old embryonic organ, and of great phylogenetic interest, because it arises in the same way in all the chordonia (both tunicates and vertebrates). In every case it touches or embraces like an arch the posterior end of the chorda, which has been developed here in front out of the middle line of the primitive gut (between the two coelom-folds of the middle groove) ("head-process," Fig. 123 *h*). These very ancient and strictly hereditary structures, which have no physiological significance to-day, deserve (as "rudimentary organs") our closest attention. The tenacity with which the useless neurenteric canal has been transmitted down to man (through the whole series of vertebrates is of equal interest for the theory of descent in general, and the phylogeny of the chordonia in particular.

The connection which the neurenteric canal (Fig. 123 *nc*) establishes between the dorsal nerve-tube (*n*) and the ventral gut-tube (*d*) is seen very




FIG. 125.—Longitudinal section of the oösomula of a frog. (From Zabor.) *nc*, neurenteric canal; *mc*, medullary canal; *st*, first primitive segment; *c*, cellular pouches.

dorsal surface of the gastrula, and is converted into the longitudinal furrow we call the primitive groove. Hence the primitive groove (Fig. 125 *pr*), examined from above, appears to be the straight

continuation of the fore-lying and younger medullary furrow (*mf*). The divergent hind legs of the latter embrace the anterior end of the former. Afterwards we have the complete closing of the primitive mouth, the dorsal swellings

While these important processes are taking place in the axial part of the dorsal shield, its external form also is changing. The oval form (Fig. 117) becomes like the sole of a shoe or sandal, lyre-shaped or finger-bisuit shaped (Fig. 120). The middle third does not grow in width as quickly as the posterior, and still less than the anterior third; thus the shape of the permanent body becomes somewhat narrow at the waist. At the same time, the oval form of the generative area returns to a circular shape, and the inner pellucid area separates more clearly from the opaque outer area (Fig. 121 a). The completion of the circle in the area marks the limit of the formation of blood-vessels in the mesoderm.

The characteristic sandal-shape of the dorsal shield, which is



FIG. 116. Longitudinal section of a frog-embryo. (From *Smith*.) *mf*, medullary furrow; *ch*, chorda; *pr*, paraxial body; *st*, stomodaeum.



FIG. 117

FIGS. 117 and 118. Dorsal shield of the chick. (From *Slater*.) The medullary furrow (*mf*), which is not yet visible in Fig. 120, encloses with its border end the fore end of the primitive groove (*pr*) in Fig. 121.




FIG. 118

joining to form the medullary tube and growing over it. The neuraxonic canal then leads directly, in the shape of a narrow arch-shaped tube (Fig. 129 *ae*), from the medullary tube (*mf*) to the gastric tube (*ga*). Directly in front of it is the latter end of the chorda (*ch*).

determined by the narrowness of the middle part, and which is compared to a violin, lyra, or shoe-sole, persists for a long time in all the amniotes. All mammals, birds, and reptiles have substantially the same construction at this stage, and even for a longer or shorter

It is all the more notable that the early separation of dorsal and ventral halves takes place in the same rigidly hereditary fashion in all the vertebrates. In both the acrania and the craniota the dorsal body is about this period separated from the ventral body. In the middle part of the body this division has already taken place by the constriction of the chorda between the dorsal nerve-tube and the ventral canal. But in the outer or lateral

proceed step by step with interesting changes in the ectoderm, while the entoderm changes little at first. We can study these processes best in transverse sections, made vertically to the surface through the saucer-shaped embryonic shield. Such a transverse section of a chick-embryo, at the end of the first day of incubation, shows the gut-gland layer as a very simple epithelium, which is spread like a leaf over the outer surface of the




FIG. 131.



FIG. 132.

FIG. 131.—Embryo of the quail, sixty hours old, cross-section of an inch in diameter. (From *Wetzel*.)
a the pharynx; *b* the dorsal epidermis; *c* the dorsal epidermis; *d* the dorsal epidermis; *e* the dorsal epidermis; *f* the dorsal epidermis; *g* the dorsal epidermis; *h* the dorsal epidermis; *i* the dorsal epidermis; *j* the dorsal epidermis; *k* the dorsal epidermis; *l* the dorsal epidermis; *m* the dorsal epidermis; *n* the dorsal epidermis; *o* the dorsal epidermis; *p* the dorsal epidermis; *q* the dorsal epidermis; *r* the dorsal epidermis; *s* the dorsal epidermis; *t* the dorsal epidermis; *u* the dorsal epidermis; *v* the dorsal epidermis; *w* the dorsal epidermis; *x* the dorsal epidermis; *y* the dorsal epidermis; *z* the dorsal epidermis.

FIG. 132.—Sagittal-shaped embryonic shield of a quail of about eight days, with the fore part of the primitive axis (on right) of dorsal axis. (From *Wetzel*.) *a* dorsal furrow, at the middle of the mandibular plate; *b*, *c* primitive groove (mouth); *d*, *e* dorsal (stem) axis; *f*, *g* ventral (parietal) axis. In the narrow middle part the first three primitive segments may be seen.

part of the body it is only brought about by the division of the coelom-pouches into two sections—a dorsal *epimeric* (dorsal segment or provertebra) and a ventral *hypomeric* (or ventral segment)—by a frontal constriction. In the amphioxus each of the former makes a muscular pouch, and each of the latter a sex-pouch or gonad.

These important processes of differentiation in the mesoderm, which we will consider more closely in the next chapter,

loud-yolk (Fig. 92). The chorda (*ch*) has separated from the dorsal middle line of the notoderm; to the right and left of it are the two halves of the mesoderm, or the two coelom-folds. A narrow cleft in the latter indicates the body-cavity (*coel*); this separates the two plates of the coelom-pouches, the lower (visceral) and upper (parietal). The broad dorsal furrow (*df*) formed by the mandibular plate (*m*) is still wide open, but is divided from the lateral body-plates

(A) by the parallel medullary swellings, which eventually close.

During these processes important changes are taking place in the outer germinal layer (the "skin-sense layer"). The continued rise and growth of the dorsal swellings causes their higher parts to bend together at their free borders, approach nearer and nearer (Fig. 136 w), and finally unite. Thus in the end we

descent it is a thoroughly natural process. The phylogenetic explanation of it is that the central nervous system is the organ by means of which all intercourse with the outer world, all psychic action and sense-perception, are accomplished; hence it was bound to develop originally from the outer and upper surface of the body, or from the outer skin. The medullary tube afterwards separates completely from




FIG. 133.




FIG. 134.

FIG. 133.—Human embryo at the neural-stage, cranial end of an early stage, from the end of the second week, magnified twenty-five times. (From *Grundriss*.)

FIG. 134.—Dorsal-shaped embryonic stage of a rabbit of nine days. (From *Kühn*.) (Back view from above.) *a*, skin-sense or dorsal shield (with eight pairs of primitive segments); *a1*, parietal or ventral zone; *a2*, notochord; *a3*, notochord; *a4*, heart; *a5*, pericardial cavity; *a6*, amniotic cavity; *a7*, amniotic cavity; *a8*, eye-vesicles; *a9*, fore brain; *a10*, middle brain; *a11*, hind brain; *a12*, head brain; *a13*, head brain; *a14*, head brain; *a15*, head brain; *a16*, head brain; *a17*, head brain; *a18*, head brain; *a19*, head brain; *a20*, head brain; *a21*, head brain; *a22*, head brain; *a23*, head brain; *a24*, head brain; *a25*, head brain; *a26*, head brain; *a27*, head brain; *a28*, head brain; *a29*, head brain; *a30*, head brain; *a31*, head brain; *a32*, head brain; *a33*, head brain; *a34*, head brain; *a35*, head brain; *a36*, head brain; *a37*, head brain; *a38*, head brain; *a39*, head brain; *a40*, head brain; *a41*, head brain; *a42*, head brain; *a43*, head brain; *a44*, head brain; *a45*, head brain; *a46*, head brain; *a47*, head brain; *a48*, head brain; *a49*, head brain; *a50*, head brain; *a51*, head brain; *a52*, head brain; *a53*, head brain; *a54*, head brain; *a55*, head brain; *a56*, head brain; *a57*, head brain; *a58*, head brain; *a59*, head brain; *a60*, head brain; *a61*, head brain; *a62*, head brain; *a63*, head brain; *a64*, head brain; *a65*, head brain; *a66*, head brain; *a67*, head brain; *a68*, head brain; *a69*, head brain; *a70*, head brain; *a71*, head brain; *a72*, head brain; *a73*, head brain; *a74*, head brain; *a75*, head brain; *a76*, head brain; *a77*, head brain; *a78*, head brain; *a79*, head brain; *a80*, head brain; *a81*, head brain; *a82*, head brain; *a83*, head brain; *a84*, head brain; *a85*, head brain; *a86*, head brain; *a87*, head brain; *a88*, head brain; *a89*, head brain; *a90*, head brain; *a91*, head brain; *a92*, head brain; *a93*, head brain; *a94*, head brain; *a95*, head brain; *a96*, head brain; *a97*, head brain; *a98*, head brain; *a99*, head brain; *a100*, head brain.

get from the open dorsal furrow, the upper cleft of which becomes narrower and narrower, a closed cylindrical tube (Fig. 137 w). This tube is of the utmost importance; it is the beginning of the central nervous system, the brain and spinal marrow, the *medullary tube*. This embryonic fact was formerly looked upon as very mysterious. We shall see presently that in the light of the theory of

the outer germinal layer, and is surrounded by the middle parts of the provertebra and forced inwards (Fig. 146). The remaining portion of the skin-sense layer (Fig. 93 A) is now called the horn-plate or horn-layer, because from it is developed the whole of the outer skin or epidermis, with all its horny appendages (nails, hair, etc.).

A totally different organ, the *prorenal*

(primitive kidney) duct (nephros), is found to be developed at an early stage from the ectoderm. This is originally a quite simple, tube-shaped, lengthy duct, or straight canal, which runs from front to rear at each side of the provertebrae (on the outer side, Fig. 93 *neg.*). It origi-

the first trace of it does not come from the skin-sense layer, but the skin-fibre layer.

The inner germinal layer, or the gut-fibre layer (Fig. 93 *abd.*), remains unchanged during these processes. A little later, however, it shows a quite flat, groove-like depression in the middle line




FIG. 132.—Dorsal-shaped embryonic shield of an *Ascidium* (*Pediculus*), three days old. (From *Seelische*.) [Back from front above; all heart-vent or dorsal shield (with right parts of primitive septum) at lateral or ventral sides; *ep* epibranchial area, *op* opercular area; *ab* halves of the heart, *st* fore-end & hind-end in the median line we see the chorda (*ch*) through the top-aperture medullary tube (*m*). *a* primitive segment, *pr* primitive streak (or primitive mouth).]

nates, it seems, out of the horn-plate at the side of the medullary tube, in the gap that we find between the provertebral and the lateral plates. The proventral duct is visible in this gap even at the time of the severance of the medullary tube from the horn-plate. Other observers think that

of the embryonic shield, directly under the chorda. This depression is called the gastric groove or furrow. This at once indicates the future lot of this germinal layer. As this ventral groove gradually deepens, and its lower edges bend towards each other, it is formed into a closed tube.

the alimentary canal, in the same way as the medullary groove grows into the medullary tube. The gut-fibre layer (Fig. 137*f*), which lies on the gut-gland layer (*d*), naturally follows it in its folding. Moreover, the incipient gut-wall consists from the first of two layers, internally the gut-gland layer and externally the gut-fibre layer.

The formation of the alimentary canal resembles that of the medullary tube to this extent—in both cases a straight groove or furrow arises first of all in the middle line of a flat layer. The edges of this furrow then bend towards each other, and join to form a tube (Fig. 137). But the two processes are really very different. The medullary tube closes in its whole length, and forms a cylindrical tube, whereas the alimentary canal remains open in the middle, and its cavity continues for a long time in connection with the cavity of the embryonic vesicle. The open connection between the two cavities is only closed at a very late stage, by the construction of the navel. The closing of the medullary tube is effected from both sides, the edges of the groove joining together from right and left. But the closing of the alimentary canal is not only effected from right and left, but also from front and rear, the edges of the ventral groove growing together from every side towards the navel. Throughout the three higher classes of vertebrates the whole of this process of the construction of the gut is closely connected with the formation of the navel, or with the separation of the embryo from the yolk-sac or umbilical vesicle.

In order to get a clear idea of this, we must understand carefully the relation of the embryonic shield to the germulative area and the embryonic vesicle. This is done best by a comparison of the five stages, which are shown in longitudinal section in Figs. 138–142. The embryonic shield (*e*), which at first projects very slightly over the surface of the germulative area, soon begins to rise higher above it, and to separate from the embryonic vesicle. At this point the embryonic shield, looked at from the dorsal surface, shows still the original simple sandal-shape (Figs. 133–135). We do not yet see any trace of articulation into head, neck, trunk, etc., or limbs. But the embryonic shield has increased greatly in thickness, especially in the anterior part. It now has the appearance

of a thick, oval swelling, strongly curved over the surface of the germulative area. It begins to sever completely from the embryonic vesicle, with which it is connected at the ventral surface. As this severance proceeds, the back bends more and more; in proportion as the embryo grows the embryonic vesicle decreases, and at last it merely hangs as a small vesicle from the belly of the embryo (Fig. 142*dt*). In consequence of the growth-movements which cause this severance, a groove-shaped depression is formed at the surface of the vesicle, the *limiting furrow*, which surrounds the vesicle in the shape of a pit, and a circular mouth or dam (Fig. 139*dt*) is formed at the outside of this pit by the elevation of the contiguous parts of the germinal vesicle.

In order to understand clearly this important process, we may compare the embryo to a fortress with its surrounding



FIG. 138. Transverse section of the embryonic shield at the end of the first day of incubation, transversed about (ventral) surface. The edges of the medullary plate (*m*), the medullary ridges (*m*), which separate the medullary from the horizontal (*h*), are bending towards each other. At each side of the shield (*e*) the primitive segment plates (*s*) have separated from the lateral plates (*l*), a gut-gland layer. (From Haeckel.)

rampart and trench. The ditch consists of the outer part of the germulative area, and comes to an end at the point where the area passes into the vesicle. The important fold of the middle germinal layer that belongs about the formation of the body-cavity spreads beyond the borders of the embryo over the whole germulative area. At first this middle layer reaches as far as the germulative area, the whole of the roof of the embryonic vesicle consists in the beginning only of the two original limiting layers, the outer and inner germinal layers. Hence, as far as the germulative area extends the germinal layer splits into the two plates we have already recognised in it, the outer skin-fibre layer and the inner gut-fibre layer. These two plates diverge considerably, a clear fluid gathering between them (Fig. 140*mm*). The inner plate, the gut-fibre layer, remains on the inner layer of the embryonic vesicle (on the gut-gland layer). The

outer plate, the skin-fibre layer, lies close on the outer layer of the germinal area, or the skin-sac layer, and separates together with this from the embryonic vesicle. From these two united outer plates is formed a continuous membrane. This is the circular mound that rises higher and higher round the whole embryo, and at last joins above it (Figs. 133-143 *ans*). To return to our illustration of the fortress, we must imagine the circular rampart to be extraordinarily high and towering far above the fortices. Its edges bend over like the cornice of an overhanging wall of rock that would enclose the fortress; they form a deep hollow, and at last join together above.

the original embryonic vesicle, start from the open belly of the embryo (Fig. 133 *st*). In more advanced embryos, in which the gastric wall and the ventral wall are nearly closed, it hangs out of the navel-opening in the shape of a small vesicle with a stalk (Figs. 141, 142 *st*). The more the embryo grows, the smaller becomes the vitelline (yolk) sac. At first the embryo looks like a small appendage of the large embryonic vesicle. Afterwards it is the yolk-sac, or the remainder of the embryonic vesicle, that seems a small pouch-like appendage of the embryo (Fig. 142 *st*). It ceases to have any significance in the end. The very wide opening, through which the gastric cavity



FIG. 142.—Three diagrammatic transverse sections of the embryonic disk of the higher vertebrates to show the degree of the cellular organs from the lowest greenish layers. In Fig. A the embryonic tube (*n*) and the alimentary canal (*a*) are still open grooves. In Fig. B the embryonic tube (*n*) and the dorsal wall are closed, but the alimentary canal (*a*) and the ventral wall are open, the preoral ducts (*u*) are not off from the horn-plate (*h*) and externally connected with segmental preoral canals. In Fig. C both the embryonic tube and the dorsal wall above and the alimentary canal and ventral wall below are closed. All the open grooves have become closed tubes. The growing hollows are directed upwards. The letters have the same meaning as in all three figures: *h* horn-plate layer, *n* embryonic tube, *u* preoral ducts, *z* anal pit, *z* primitive-ventrals, *v* dorsal wall, *d* ventral wall, *e* body-cavity or coelom, *f* gut-fibre layer, *g* primitive artery (aorta), *o* primitive vein (inferior vena cava), *s* gut-fibre layer, *a* alimentary canal.

In the end the fortress lies entirely within the hollow that has been formed by the growth of the edges of this large rampart.

As the two outer layers of the germinal area thus rise in a fold about the embryo, and join above it, they come at last to form a spacious sac-like membrane about it. This envelope takes the name of the germinal membrane, or water-membrane, or amnion (Fig. 142 *ans*). The embryo floats in a watery fluid, which fills the space between the embryo and the amnion, and is called the amniotic fluid (Figs. 141, 142 *st*). We will deal with this remarkable formation and with the allantois later on (Chapter XV.). In front of the allantois the yolk-sac or gut-bell vesicle (*st*), the remainder of

at first communicates with the umbilical vesicle, becomes narrower and narrower, and at last disappears altogether. The rest, the small pit-like depression that we find in the developed man in the middle of the abdominal wall, is the spot at which the remainder of the embryonic vesicle (the umbilical vesicle) originally entered into the ventral cavity, and joined on to the growing gut.

The origin of the navel coincides with the complete closing of the external ventral wall. In the amnion the ventral wall originates in the same way as the dorsal wall. Both are formed substantially from the skin-fibre layer, and externally covered with the horn-plate, the border section of the skin-sac layer. Both come into

existence by the conversion of the four flat germinal layers of the embryonic shield into a double tube by folding from opposite directions; above, at the back, we have the vertebral canal which encloses

the closing in the middle of the dorsal wall take place in the same way as the medullary tube, which is henceforth enclosed by the vertebral tube. Thus is formed the dorsal wall, and the medullary

tube takes up a position inside the body. In the same way the provertebril mass grows afterwards round the chorda, and forms the vertebral column. Below this the inner and outer edge of the provertebral plate splits on each side into two horizontal plates, of which the upper forms between the chorda and medullary tube, and the lower between the chorda and mantle tube.

As the plates meet from both sides above and below the chorda, they completely enclose it, and so form the tubular, outer, chondri-sheath, the sheath from



Fig. 243.

the medullary tube, and below, at the belly, the wall of the body-cavity which contains the alimentary canal (Eur. 112).

We will consider the formation of the dorsal wall first, and that of the ventral wall afterwards. (Figs. 143-147). In the middle of the dorsal surface of the embryo there is originally, as we already know, the medullary (*mr*) tube directly underneath the horn-plate (*h*), from the middle part of which it has been developed. Later, however, the provertebral plates (*pw*) grow over from the right and left between these originally connected parts (Figs. 146, 146b). The upper and inner edges of the two provertebral plates push between the horn-plate and



Fig. 14

[illegible]

medullary tube, force them away from each other, and finally join between them in a seam that corresponds to the middle line of the back. The coalescence of these two dorsal plates and

We find in the construction of the ventral wall precisely the same processes

as in the formation of the dorsal wall (Fig. 132 B, Fig. 144 A, Fig. 146 AA). It is formed on the flat embryonic shield of the annulotes from the upper plates of the

yell sac (Fig. 105). The external navel in the skin is the definitive point of the closing of the ventral wall; this is visible in the developed body as a small depression,




FIG. 145

parietal zone. The right and left parietal plates bend downwards towards each other, and grow round the gut in the same way as the gut itself closes. The outer part of the lateral plates forms the ventral wall or the lower wall of the body, the two lateral plates bending considerably on the inner side of the amiotic fold, and growing towards each other from right and left. While the alimentary canal is closing, the body-wall also closes on all sides. Hence the ventral wall, which encloses the whole ventral cavity below, consists of two parts, two lateral plates that bend towards each other. These approach each other all along, and at last meet at the navel. We ought, therefore, really to distinguish two navels, an inner and an outer one. The internal or intestinal navel is the definitive point of the closing of the gut wall, which puts an end to the open communication between the ventral cavity and the cavity of the




FIG. 146

With the formation of the external navel and the closing of the alimentary canal is connected the formation of two cavities, which we call the capital and the pelvic sections of the visceral cavity. As the embryonic shield lies flat on the wall of the embryonic vesicle at first, and only gradually separates from it, its fore and hind ends are independent in the beginning; on the other hand, the middle part of the ventral surface is connected with the yolk-sac by means of the vitelline or umbilical duct (Fig. 147 m). This leads to a notable curving of the dorsal surface, the head-end bends downwards towards the breast and the tail-end towards the

As a result of these processes the embryo attains a shape that may be compared to a wooden shoe, or, better still, to an overturned canoe. Imagine a canoe or boat with both ends rounded and a small covering before and behind; if this canoe is turned upside down, so that the curved keel is uppermost, we have a fair picture of the canoe-shaped embryo (Fig. 147). The upturned convex keel corresponds to the middle line of the back; the small chamber underneath the fore-deck represents the capital cavity, and the small chamber under the rear-deck the pelvic chamber of the gut (cf. Fig. 140).




FIG. 147.—Median longitudinal section of the embryo of a chick (15th day of incubation), seen from the right side (head to the right and to the left). Dorsal half dark, with convex outline. *a* gut, *m* mouth, *g* gullet, *h* heart, *b* ventricle of the heart, *c* vitelline or umbilical duct, *d* yolk-sac, *e* vitelline or umbilical duct, *f* vitelline or umbilical duct, *g* vitelline or umbilical duct, *h* vitelline or umbilical duct, *i* vitelline or umbilical duct, *j* vitelline or umbilical duct, *k* vitelline or umbilical duct, *l* vitelline or umbilical duct, *m* vitelline or umbilical duct, *n* vitelline or umbilical duct, *o* vitelline or umbilical duct, *p* vitelline or umbilical duct, *q* vitelline or umbilical duct, *r* vitelline or umbilical duct, *s* vitelline or umbilical duct, *t* vitelline or umbilical duct, *u* vitelline or umbilical duct, *v* vitelline or umbilical duct, *w* vitelline or umbilical duct, *x* vitelline or umbilical duct, *y* vitelline or umbilical duct, *z* vitelline or umbilical duct. (From Baer.)

belly. We see this very clearly in the excellent old diagrammatic illustration given by Baer (Fig. 147), a median longitudinal section of the embryo of the chick, in which the dorsal body or epinota is deeply shaded. The embryo seems to be trying to roll up, like a hedgehog protecting itself from its pursuers. This pronounced curve of the back is due to the more rapid growth of the convex dorsal surface, and is directly connected with the severance of the embryo from the yolk-sac. The further bending of the embryo leads to the formation of the "head-cavity" of the gut (Fig. 148, above *Z*) and a similar one at the tail, known as its "pelvic cavity."

The embryo now, as it were, presses into the outer surface of the embryonic vesicle with its free ends, while it moves away from it with its middle part. As a result of this change the yolk-sac becomes henceforth only a pouch-like outer appendage at the middle of the ventral wall. The ventral appendage, growing smaller and smaller, is afterwards called the umbilical (navel) vesicle. The cavity of the yolk-sac or umbilical vesicle communicates with the corresponding visceral cavity by a wide opening, which gradually contracts into a narrow and long canal, the vitelline (yolk) duct (*ductus vitellinus*, Fig. 147 m). Hence, if we were to imagine ourselves in

the cavity of the yolk-sac, we could get from it through the yolk-duct into the middle and still wide open part of the alimentary canal. If we were to go forward from there into the head-part of the embryo, we should reach the capital cavity of the gut, the fore-end of which is closed up.

The reader will ask: "Where are the mouth and the anus?" These are not at first present in the embryo. The whole of the primitive gut-cavity is completely closed, and is merely connected in the middle by the vitelline duct with the equally closed cavity of the embryonic vesicle (Fig. 140). The two later apertures of the alimentary canal—the anus and the mouth—are secondary constructions, formed from the outer skin. In the horn-plate, at the spot where the mouth is found subsequently, a pit-like depression is formed, and this grows deeper and deeper, pushing towards the blind fore-end of the capital cavity; this is the mouth-pit. In the same way, at the spot in the outer skin where the anus is afterwards situated a pit-shaped depression appears, grows deeper and deeper, and approaches the blind hind-end of the pelvic cavity, this is the anus-pit. In the end these pits touch with their deepest and innermost points the two blind ends of this primitive alimentary canal, so that they are now only separated from them by thin membranous partitions. This membrane finally disappears, and henceforth the alimentary canal opens in front at the mouth and in the rear by the anus (Figs. 142, 147). Hence at first, if we penetrate into these pits from without, we find a partition cutting them off from the cavity of the alimentary canal, which gradually disappears. The formation of mouth and anus is secondary in all the vertebrates.

During the important processes which lead to the formation of the navel, and of the intestinal wall and ventral wall, we find a number of other interesting changes taking place in the embryonic shield of the amniotes. These relate chiefly to the prorenal ducts and the first blood-vessels. The prorenal (primitive kidney) ducts, which at first lie quite flat under the horn-plate or epiderm (Fig. 93 *sup.*), soon back towards each other in consequence of special growth movements (Figs. 143-

145 *sup.*). They depart more and more from their point of origin, and approach the gut-gland layer. In the end they lie deep in the interior, on either side of the mesentery, underneath the chords (Fig. 145 *sup.*). At the same time, the two primitive aortas change their position (cf. Figs. 138-145 *op.*): they travel inwards underneath the chords, and there coalesce at last to form a single secondary aorta, which is found under




FIG. 145.—Longitudinal section of the fore half of a chick-embryo at the end of the first day of incubation (seen from the left side). *a*, blasto-epithelium on chords. Above *a* is the blind fore-end of the ventral tube (*m*). Below *a* the capital cavity of the gut *a*, gut-gland layer, *d*, gut-gland layer, *b*, horn-plate, *c*, cavity of the blasto-epithelium, *h*, heart-ventricle, *k*, heart-cavities. (From *Reichert*.)

the rudimentary vertebral column (Fig. 145 *op.*). The cardinal veins, the first ventral blood-vessels, also back towards each other, and eventually unite immediately above the rudimentary kidneys (Figs. 145 *nc.*, 152 *op.*). In the same spot, at the inner side of the fore-kidneys, we soon see the first trace of the sexual organs. The most important part of this apparatus (apart from all its appendages) is the ovary in the female and the testicle




FIG. 129.—Longitudinal section of a human embryo of the fourth week, one-fifth of an inch long, magnified fifteen times. (From Kellmann.)




FIG. 130.

FIG. 130.—Transverse section of a human embryo of fourteen days. *mr* mesoderm tube, *ch* chorda, *mp* mesoderm, *st* stomach, *sp* somites, *lf* lateral fold, *df* dorsal fold, *ab* allantois, *tu* tubule. (From Kellmann.)




FIG. 131.

FIG. 131.—Transverse section of a sheep embryo (or young mink). *mr* mesoderm tube, *ch* chorda, *st* stomach, *sp* somites, *lf* lateral fold, *df* dorsal fold, *ab* allantois, *tu* tubule. (From H. E. Hager.)

in the male. Both develop from a small part of the out-lining of the body-cavity, at the spot where the skin-fibre layer and gut-fibre layer touch. The connection of

this embryonic gland with the prorenal ducts, which lie close to it and assume most important relations to it, is only secondary.




FIG. 125.—Transverse section of a shark-embryo with twenty-four primitive segments. (From Pfeffer.) Shows a dorsal lateral part of the medullary tube (*sp 9*), the spinal ganglia (*sp 9*) just out between it and the neuro-plaxia. *ch* chorion, an double aorta, *h. y.* hyoid, layer of gut-fibre layer, with blood-vessels on surface, one muscle, *pl. m.* on the dorsal wall of the stomach (*g. m.*). Below the cardiac vein (*v. d.*) in the prorenal duct (*v. d.*) and a segmental prorenal gland (*s. d.*). The skin-fibre layer of the body-wall (*m. s.*) is indicated in the anterior (*h. y.*) part. Between the four secondary prorenal organs and the structures formed from them there is formed embryonic connective tissue with secretory cells and vascular structures (forming a "mesenchym").

CHAPTER XIV.

THE ARTICULATION OF THE BODY¹

THE VERTEBRATE stem, to which our race belongs as one of the latest and most advanced outgrowths of the natural development of life, is rightly placed at the head

of the animal kingdom. This privilege must be accorded to it, not only because man does in point of fact soar far above all other animals, and has been fitted to

¹ The term "articulation" is used in this chapter to denote both "segmentation" and "articulation" in the ordinary sense.—T. W. M.

the position of "lord of creation"; but also because the vertebrate organisms far surpasses all the other animal-stems in size, in complexity of structure, and in the advanced character of its functions. From the point of view of both anatomy and physiology, the vertebrate stem outstrips all the other, or invertebrate, animals.

There is only one among the twelve stems of the animal kingdom that can in many respects be compared with the vertebrates, and reaches an equal, if not a greater, importance in many points. This is the stem of the articulatees, composed of three classes: 1, the annelids (earth-worms, leeches, and cognate forms); 2, the crustacea (crabs, etc.); 3, the tracheata (spiders, insects, etc.). The stem of the articulatees is superior not only to the vertebrates, but to all other animal-stems, in variety of forms, number of species, elaborateness of individuals, and general importance in the economy of nature.

When we have thus declared the vertebrates and the articulatees to be the most important and most advanced of the twelve stems of the animal kingdom, the question arises whether this special position is accorded to them on the ground of a peculiarity of organization that is common to the two. The answer is that this is really the case; it is their segmental or transverse articulation, which we may briefly call *metamerism*. In all the vertebrates and articulatees the developed individual consists of a series of successive members (segments or metameres—"parts"); in the embryo these are called primitive segments or somites. In each of these segments we have a certain group of organs reproduced in the same arrangement, so that we may regard each segment as an individual unity, or a special "individual" subordinated to the entire personality.

The similarity of their segmentation, and the consequent physiological advance in the two stems of the vertebrates and articulatees, has led to the assumption of a direct affinity between them, and an attempt to derive the former directly from the latter. The annelids were supposed to be the direct ancestors, not only of the crustacea and tracheata, but also of the vertebrates. We shall see later (Chapter XX.) that this annelid theory of the vertebrates is entirely wrong, and ignores the most important differences in the organization of the two stems. The internal articulation of the vertebrates is


just as profoundly different from the external metamerism of the articulatees as are their skeletal structure, nervous system, vascular system, and so on. The articulation has been developed in a totally different way in the two stems. The unarticulated chordula (Figs. 83-86), which we have recognized as one of the chief paleogenetic embryonic forms of the vertebrate group, and from which we have inferred the existence of a corresponding ancestral form for all the vertebrates and tunicates, is quite unthinkable as the stem-form of the articulatees.

All articulated animals came originally from unarticulated ones. This phylogenetic principle is as firmly established as the ontogenetic fact that every articulated animal-form develops from an unarticulated embryo. But the organization of the embryo is totally different in the two stems. The chordula-embryo of all the vertebrates is characterized by the dorsal medullary tube, the neurenteric canal, which passes at the primitive mouth into the alimentary canal, and the axial chorda between the two. None of the articulatees, either annelids or arthropods (crustacea and tracheata), show any trace of this type of organization. Moreover, the development of the chief systems of organs proceeds in the opposite way in the two stems. Hence the segmentation must have arisen independently in each. This is not at all surprising; we find analogous cases in the stalk-articulation of the higher plants and in several groups of other animal stems.

The characteristic internal articulation of the vertebrates and its importance in the organization of the stem are best seen in the study of the skeleton. Its chief and central part, the cartilaginous or bony vertebral column, affords an obvious instance of vertebrate metamerism; it consists of a series of cartilaginous or bony pieces, which have long been known as *vertebre* (or *vertebræ*). Each vertebra is directly connected with a special section of the muscular system, the nervous system, the vascular system, etc. Thus most of the "animal organs" take part in this vertebration. But we saw, when we were considering our own vertebrate character (in Chapter XI.), that the same internal articulation is also found in the lowest primitive vertebrates, the acrania, although here the whole skeleton consists merely of the simple chorda, and is not at all articulated.

Hence the articulation does not proceed primarily from the skeleton, but from the muscular system, and is clearly determined by the more advanced swimming-movements of the primitive chordoma-uctors.

"somites" or primitive segments, to these so-called "primitive vertebrae." If the latter name is retained at all, it should only be used of the *achyrosomae*, i.e., the small part of the somites from which the later vertebrae does actually develop.



FIGS. 151-153.—Fish-shaped embryos (right of the black, in three successive stages of development, looked at from the dorsal surface, magnified about twenty times, schematic diagrams). Fig. 151 with six pairs of blastomeres. Above a simple vesicle (AV). Medullary layer still wide open (M), greatly widened at a ventral medullary plate, at lateral plates, a band of gut-cavity (G) and foregut (FG). Fig. 152 with ten pairs of somites. Blast divided into three vesicles: a fore-vesicle, an middle-vesicle, a hind-vesicle. A band of ventral medullary layer still wide open (AV), at medullary plate. Fig. 153 with sixteen pairs of somites. Blast divided into five vesicles: a fore-vesicle, a sub-mesodermic-vesicle, an middle-vesicle, a hind-vesicle, a sub-epithelial, a type vesicle, a medullary vesicle, a hind, at ventral vesicle, at medullary plate, one primitive vesicle.

It is, therefore, wrong to describe the first rudimentary segments in the vertebrate embryo as primitive vertebrae or *provertebrae*; the fact that they have been so called for some time has led to much error and misunderstanding. Hence we shall give the name of

Articulation begins in all vertebrates at a very early embryonic stage, and this indicates the considerable phylogenetic age of the process. When the chondula (Figs. 83-86) has completed its characteristic composition, often even a little earlier, we find in the arachnoid, in the

for a time in the amphioxus) remains undivided, and, by the divergence of their parietal and visceral plates, form a body-cavity that is unified from the first. In this case, again, it is clear that we must regard the features of the younger craniota as canonically modified processes that can be traced palaeogenetically to the older acrania.

We have an interesting intermediate stage between the acrania and the fishes in these and many other respects in the cyclostoma (the hag and the lamprey, cf. Chapter XXI.)

Among the fishes the selachii, or primitive fishes, yield the most important information on these and many other phylogenetic questions (Figs. 161, 162). The careful studies of Ruckert, Van Wijbe, H. E. Ziegler, and others, have given us most valuable results. The products of

series of pairs of these vesicular dorsal segments, which have been cut off on each side from the ventral side-plates, and lie to the right and left of the chorda.

The metamorphosis of the amniotes agrees in all essential points with that of the



FIG. 164.



FIG. 165.

FIG. 164.—The lateral cervical vertebra (human).

FIG. 165.—The sixth dorsal vertebra (human).

three lower classes of vertebrates we have considered; but it varies considerably in detail, in consequence of cenogenetic disturbances that are due in the first place (like the degeneration of the coelom-pouches) to the large development of the food-yells. As the pressure of this seems to force the two middle layers together from the start, and as the solid structure of the mesoderm apparently belies the original hollow character of the sacs, the two sections of the mesoderm, which are at that time divided by the lateral fold—the dorsal segment-plates and ventral side-plates—have the appearance at first of solid layers of cells (Figs. 93-97). And when the articulation of the somites begins in the sole-shaped embryonic shield, and a couple of protovertebrae are developed in succession, constantly increasing in number towards the rear,




FIG. 163.—Frontal (or horizontal-longitudinal) section of a trilobian embryo with three pairs of primitive segments. A, chorda; B, primitive segment; with their cavity, and their plate.

the middle germinal layer are partly clear in these cases at the period when the dorsal primitive segment cavities (or myotoms, A) are still connected with the ventral body-cavity (B; Fig. 161). In Fig. 162, a somewhat older embryo, these cavities are separated. The outer or lateral wall of the dorsal segment yields the cutis-plate (cp), the foundation of the connective cutis. From its inner or median wall are developed the muscle-plate (m), the rudiment of the trunk-muscles, and the skeletal plate, the formative matter of the vertebral column (sk).

In the amphibia, also, especially the water-salamander (*Triton*), we can observe very clearly the articulation of the coelom-pouches and the rise of the primitive segments from their dorsal half (cf. Fig. 91, A, B, C). A horizontal longitudinal section of the salamander-embryo (Fig. 163) shows very clearly the



FIG. 166.—The second lumbar vertebra (human).

these tube-shaped somites (formerly called protovertebrae, or primitive vertebrae) have the appearance of solid discs, made up of mesodermic cells (Fig. 93). Nevertheless, there is for a time a ventral cavity, or provertebral cavity, even in these solid

"provertebrae" (Fig. 143 *uvd*). This vesicular condition of the provertebra is of the greatest phylogenetic interest; we must, according to the common theory, regard it as an hereditary reproduction of the hollow dorsal scutites of the amphioxus (Figs. 135-140) and the lower vertebrates (Figs. 161-163). This rudimentary "provertebral cavity" has no physiological significance whatever in the acorn-like embryo; it soon disappears, being filled up with cells of the muscular plate.

The innermost median part of the

divides into two plates, which grow round the chorda, and thus form the foundation of the body of the vertebra (*uv*). The upper plate presses between the chorda and the medullary tube, the lower between the chorda and the alimentary canal (Fig. 137 *C*). As the plates of two opposite provertebral pieces unite from the right and left, a circular sheath is formed round this part of the chorda. From this develops the body of a vertebra—that is to say, the massive lower or ventral half of the bony ring, which is called the "vertebra."




FIG. 147.—Head of a starfish embryo (*Pteraster*), one-third of an inch long, magnified twenty times (From Perrier). Seen from the ventral side.

primitive segment plates, which lies immediately on the chorda (Fig. 145 *ch*) and the medullary tube (*u*), forms the vertebral column in all the higher vertebrates (it is wanting in the lowest); hence it may be called the *axial* plate. In each of the provertebrae it is called the "sclerotome" (in opposition to the outlying muscular plate, the "myotome"). From the phylogenetic point of view the myotomes are much older than the sclerotomes. The lower or ventral part of each sclerotome (the inner and lower edge of the cube-shaped provertebra)

proper and surrounds the medullary tube (Figs. 144-146). The upper or dorsal half of this bony ring, the *vertebral arch* (Fig. 145 *uv*), arises in just the same way from the upper part of the skeletal plate, and therefore from the inner and upper edge of the cube-shaped primitive vertebra. As the upper edges of two opposing somites grow together over the medullary tube from right and left, the vertebral arch becomes closed.

The whole of the secondary vertebra, which is thus formed from the union of the skeletal plates of two provertebral pieces

and encloses a part of the chorda in its body, consists at first of a rather soft mass of cells; this afterwards passes into a firmer, cartilaginous stage, and finally into a third, permanent, bony stage. These three stages can generally be distinguished in the greater part of the skeleton of the higher vertebrates; at first most parts of the skeleton are soft, tender, and membranous, they then become cartilaginous in the course of their development, and finally bony.

At the head part of the embryo in the annelotes there is not generally a cleavage of the middle germinal layer into pre-vertebral and lateral plates, but the dorsal and ventral somites are blended from the first, and form what are called the "head-plates" (Fig. 148 A). From these are



FIG. 148A.

FIG. 148B.

FIG. 148A and B.—Head of a chick embryo, of the third day. Fig. 148A from the front, Fig. 148B from the side. *a*, rudimentary fore-limbatory pit; *b*, rudimentary eye; *c*, rudimentary ear; *d*, rudimentary ear; *e*, fore-brain; *f*, eye-cup. Of the three pairs of gill-slits, the first has passed into a process of the upper jaw (*g*) and of the lower jaw (*h*). (From Kollmer.)

formed the skull, the bony case of the brain, and the muscles and cartium of the body. The skull develops in the same way as the membranous vertebral column. The right and left halves of the head curve over the cerebral vesicle, enclose the foremost part of the chorda below, and thus finally form a simple, soft, membranous capsule about the brain. This is afterwards converted into a cartilaginous primitive skull, such as we find permanently in many of the fishes. Much later this cartilaginous skull becomes the permanent bony skull with its various parts. The bony skull in man and all the other annelotes is more highly differentiated and modified than that of the lower vertebrates, the amphibia and fishes. But as the one has arisen phylogenetically from the other, we must assume

that in the former no less than the latter the skull was originally formed from the sclerotomes of a number of (at least nine) head-somites.

While the articulation of the vertebrate body is always obvious in the episoma or dorsal body, and is clearly expressed in the segmentation of the muscular plates and vertebrae, it is more latent in the hyposome or ventral body. Nevertheless, the hyposomes of the ventral half of the body are not less important than the episomes of the animal half. The segmentation in the ventral cavity affects the following principal systems of organs: 1, the gonads or sex-glands (gonosomes), 2, the nephridia or kidneys (nephro-



FIG. 149.—Head of a dog embryo, seen from the front. *a*, the two lateral halves of the foremost cerebral vesicle; *b*, rudimentary eye; *c*, middle cerebral vesicle; *d*, fore part of gill-slits; *e*, upper part of fore-limbary process; *f*, *f'*, second, third, and fourth pairs of gill-slits; *g*, *h*, heart (right, left); *i*, left auricle; *j*, left, right ventricle; *k*, lungs of the embryo with three pairs of bronchi, which go to the gill-slits. (From Strecker.)

tonies); and 3, the hind-gut with its gill-slits (branchiostomes).

The metamorphism of the hyposome is less conspicuous because in all the craniotes the cavities of the ventral segments, in the walls of which the asexual products are developed, have long since coalesced, and formed a single large body-cavity, owing to the disappearance of the partition. This cenogenetic process is so old that the cavity seems to be unsegmented from the first in all the craniotes, and the rudiment of the gonads also is almost always unsegmented. It is the more interesting to learn that, according to the important discovery of Ruckert, this asexual structure is at first segmental even in the actual schisth, and the several

gastrostomes only bleed into a simple sexual gland on either side secondarily.

Amphioxus, the sole surviving representative of the acrania, once more yields us most interesting information; in this case the sexual glands remain segmented

cavities, formed from the hypostomites of the trunk.

The gonads are the most important segmental organs of the hypostoma, in the sense that they are phylogenetically the oldest. We find sexual glands (as pouch-




FIG. 171.—Human embryo of the fourth week (twelve)–six days old, one-fourth of an inch in length magnified twenty times. (From *Atoll*.) The rudiments of the central nervous and the roots of the spinal nerves are especially marked. Underneath the four gill-arches (left side) is the heart (with auricle, I, and ventricle, A), under the aorta the liver (L).

throughout life. The sexually mature lancelet has, on the right and left of the gut, a series of metamorous sacs, which are filled with ova in the female and sperm in the male. These segmental gonads are originally nothing else than the real gastrostomes, separate body-

like appendages of the gameto-causal system) in most of the lower animals, even in the medusae, etc., which leave no kidneys. The latter appear first (as a pair of excretory tubes) in the platodes (turbellaria), and have probably been inherited from these by the articulate


(annelids) on the one hand and the unarticulated prochordons on the other, and from these passed to the articulated vertebrates. The oldest form of the kidney system in this stem are the segmental pronephridia or pronephal canals, in the same arrangement as Boveri found them in the amphioxus. They are small canals that lie in the frontal plane, on each side of the chorda, between the episoma and hyposoma (Fig. 102 n), their internal funnel-shaped opening leads into the various body-cavities, their outer opening is the lateral furrow of the epidermis. Originally they must have had a double function, the carrying away of the urine from the epinotites and the release of the sexual cells from the hypsomites.

The recent investigations of Ruckert and Van Wyhe on the mesodermic segments of the trunk and the excretory system of the selachii show that these "primitive fishes" are closely related to the amphioxus in this further respect. The transverse section of the shark-embryo in Fig. 161 shows this very clearly.

In other higher vertebrates, also, the kidneys develop (though very differently formed later on) from similar structures, which have been secondarily derived from the segmental pronephridia of the acrania. The parts of the mesoderm at which the first traces of them are found are usually called the middle or mesenteric plates. As the first traces of the gonads make their appearance in the middle plates nearer inward (i. e. middle) from the inner funnels of the nephren-canal, it is better to count this part of the mesoderm with the hyposoma.

The chief and oldest organ of the vertebrate hyposoma, the alimentary canal, is generally described as an unsegmented organ. But we could just as well say that it is the oldest of all the segmental organs of the vertebrate, the double row of the coelom-pouches grows out of the dorsal wall of the gut, on either side of the chorda. In the brief period these segmental coelom-pouches are still openly connected with the gut, they look just like a double chain of segmented visceral glands. But apart from this, we have originally in all vertebrates an important articulation of the fore-gut, that is wanting in the lower gut, the

gut, and the gill-arches that separated them, were presumably also segmental, and distributed among the various metameres of the chain, like the gonads in the after-gut and the nephridia. In the amphioxus, too, they are still segmentally formed. Probably there was a division of labour of the hypsomites in the older (and long extinct) acrania, in such wise that those of the fore-gut took over the function of breathing and those of the after-gut that of reproduction. The former developed into gill-pouches, the latter into



—Transverse section of the chordoid

side of it is a cartilaginous kidney. The gut subdivides into the a

sex-pouches. There may have been primitive kidneys in both. Though the gills have lost their function in the higher parts of them have been generally maintained in the embryo by a tenacious heredity. At a very early stage we notice in the embryo of man and the other amniotes, at each side of the head, the remarkable and important structures which we call the gill-arches and gill-clefts (Figs. 167-170 f). They are the characteristic and indispensable organs of the amniote-embryo, and are found always in the same

The gill-clefts, which originally in the older acrania pierced the wall of the fore-

spot and with the same arrangement and structure. There are formed to the right and left in the lateral wall of the fore-gut cavity, in its foremost part, first a pair and then several pairs of sac-shaped inlets, that pierce the whole thickness of the lateral wall of the head. They are thus converted into clefts, through which one can penetrate freely from without into the gullet. The wall thickens between these branchial folds, and changes into an arch-like or sickle-shaped piece—the gill, or gullet-arch. In this the muscles and skeletal parts of the branchial

higher vertebrates they afterwards disappear. The branchial arches are converted partly into the jaws, partly into the bones of the tongue and the ear. From the first gill-cleft is formed the tympanic cavity of the ear.

There are few parts of the vertebrate organism that, like the outer covering or integument of the body, are not subject to metamerism. The outer skin (*epidermis*) is unsegmented from the first, and proceeds from the continuous horny plate. Moreover, the underlying *cutis* is also not metamerous, although it develops from the segmental structure of the cutis-plates (Figs. 161, 162 *cp*). The vertebrates are strictly and profoundly different from the articulates in these respects also.

Further, most of the vertebrates still have a number of unarticulated organs, which have arisen locally, by adaptation of particular parts of the body to certain special functions. Of this character are the sense-organs in the episcium, and the lungs, the heart, the spleen, and the large visceral glands—lungs, liver, pancreas, etc.—in the hypostomium. The heart is originally only a local spindle-shaped enlargement of the large ventral blood-vessel or principal vein, at the point where the subintestinal passes into the branchial artery, at the limit of the head and trunk (Figs. 170, 171). The three higher sense-organs—nose, eye, and ear—were originally developed in the same form in all the crinoids, as three pairs of small depressions in the skin at the side of the head.

The organ of smell, the nose, has the appearance of a pair of small pits above the mouth-aperture, in front of the head (Fig. 169 *n*). The organ of sight, the eye, is found at the side of the head, also in the shape of a depression (Figs. 169 *f*, 170 *g*), to which corresponds a large outgrowth of the foremost cerebral vesicle on each side. Farther behind, at each side of the head, there is a third depression, the first trunk of the organ of hearing (Fig. 169 *g*). As yet we can see nothing of the later elaborate structure of these organs, nor of the characteristic build of the face.

When the human embryo has reached




FIG. 175.—Transverse section of the pelvic region and hind legs of a chordate embryo of the fourth day, magnified about forty times. *A*, fore-plate, or medullary tube; *v*, canal of the tube; *h*, posterior body pit; *f*, rhomboid; *g*, hind legs; *u*, allantoic canal in the ventral wall; *e*, aorta; *g*, rhomboid; *s*, gut; *d*, gut-gland tissue; *f*, gut-lining layer; *g*, embryo; *u*, episcium; *v*, dorsal muscle; *s*, body-cavity or coelom. (From H. Haeckel.)

gut separate; a blood-vessel arch rises afterwards on their inner side (Fig. 91 *do*). The number of the branchial arches and the clefts that alternate with them is four or five on each side in the higher vertebrates (Fig. 170 *d*, *f*, *f*, *f*). In some of the fishes (selachii) and in the cyclostoma we find six or seven of them permanently.

These remarkable structures had originally the function of respiratory organs—gills. In the fishes the water that serves for breathing, and is taken in at the mouth, will always pass out by the branchial clefts at the sides of the gullet. In the

... stage of development. It can still scarcely be distinguished from any other higher vertebrate. All the chief parts of the body are now laid out: the head with its skull, the rudiments of the three higher sense-organs and the five cerebral vesicles, and the gill-arches and clasts, the trunk

significance. From it we can gather the most important phylogenetic conclusions.

There is still no trace of the limbs. Although head and trunk are separated and all the principal internal organs are laid down, there is no indication whatever of the "extremities" at this stage; they are formed later on. Here again we



FIG. 134.—Development of the lizard's legs (*Lacerta agilis* L.) with special reference to their blood-vessels. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100. (From *J. Huxley*.)

with the spinal cord, the rudiment of the vertebral column, the chain of metameres, the heart and chief blood-vessels, and the kidneys. At this stage man is a higher vertebrate, but shows no essential morphological difference from the embryo of the mammals, the birds, the reptiles, etc. This is an ontogenetic fact of the utmost

importance. It is a fact of the utmost interest. It is that the older vertebrates had no limbs as we find to be the case in the lowest vertebrates (amphioxus and the tunicates). The descendants of these limbless vertebrates only acquired limbs—two fore-legs and two hind-legs—at a much later stage of development.

at first all alike, though afterwards vary considerably in structure—becoming fins (of breast and belly) in the fishes, wings and legs in the birds, fore and hind legs in the creeping

which represent at first simple roundish knobs or plates. Gradually each of these plates becomes a large projection, in which we can distinguish a small inner part and a broader outer part. The latter



FIG. 172.—*HUMAN EMBRYO*, five weeks old, half an inch long, seen from the right, magnified ten times (From *Human Embryos and Human Larvae*). In the undisturbed foetus we see the eye, mouth, and ear. In the trunk the skin and part of the muscles have been removed, so that the cartilaginous vertebral column is free; the dorsal root of a spinal nerve goes out from each vertebra (towards the skin of the back). In the middle of the lower half of the figure part of the ribs and intercostal muscles are visible. The skin and muscles have also been removed from the right limb; the external rudiments of the five fingers of the hand, and five toes of the foot, are clearly seen without the foetal pad, and also the strong network of nerves that goes from the spinal cord to the extremities. The tail projects under the foot, and to the right of it is the first part of the umbilical

animals, arms and legs in the apes and man. All these parts develop from the same simple original structure, which forms secondarily from the trunk-wall (Figs. 172, 173). They have always the appearance of two pairs of small buds,

is the rudiment of the foot or hand, the former that of the leg or arm. The similarity of the original rudiment of the limbs in different groups of vertebrates is very striking.

How the five fingers or toes with their

If there is an intimate causal connection between the processes of embryology and stem-history, as we must assume in virtue of the laws of heredity, several important phylogenetic conclusions follow at once from these ontogenetic facts. The profound and remarkable similarity in the embryonic development of man and the

other vertebrates can only be explained when we admit their descent from a common ancestor. As a fact, this common descent is now accepted by all competent scientists, they have substituted the natural evolution for the supernatural creation of organisms.

CHAPTER XV.

FOETAL MEMBRANES AND CIRCULATION

Among the many interesting phenomena that we have encountered in the course of human embryology, there is an especial

the other viviparous mammals. As a fact, all the embryonic peculiarities that distinguish the mammals from other



FIG. 175.—Human embryos from the second to the fifteenth week, natural size, seen from (on left, the dorsal back turned towards the right) II. of fourteen days. III. of three weeks. IV. of four weeks. V. of five weeks. VI. of six weeks. VII. of seven weeks. VIII. of eight weeks. XII. of twelve weeks. XV. of fifteen weeks.

importance in the fact that the development of the human body follows from the beginning just the same lines as that of

animals are found also in man; even the ovum with its distinctive membrane (*ovum pellucidum*, Fig. 14) shows the same typical

structure in all mammals (apart from the older oviparous monotremes). It has long since been deduced from the structure of the developed man that his natural place in the animal kingdom is among the mammals. Linné (1735) placed him

with the apes, in one and the same order (*primates*), in his *Systema Naturæ*. This position is fully confirmed by comparative embryology. We see that man entirely resembles the higher mammals, and most of all the apes, in embryonic development as well as in anatomic structure. And if we seek to understand this ontogenetic agreement in the light of the biogenetic law, we find that it proves clearly and necessarily the descent of man from a series of other mammals, and proximately from the primates. The common origin of man and the other mammals from a single ancient stem-form can no longer be questioned, nor can the immediate blood-relationship of man and the ape.

The essential agreement in the whole bodily form and inner structure is still visible in the embryo of man and the other mammals at the late stage of development at which the mammal-body can be recognized as such. But at a somewhat earlier stage, in which the limbs, gill-arches, sense-organs, etc., are already outlined, we cannot yet recognize the mammal-embryo as such, or distinguish them from those

micellulars). Finally, we see that the cells which make up the primary germinal layers owe their origin in every case to the repeated cleavage of a single simple cell, the stem-cell or fertilized ovum.

It is impossible to lay too much stress on this remarkable agreement in the chief embryonic features in man and the other animals. We shall make use of it later on for our monophyletic theory of descent—the hypothesis of a common descent of man and all the metazoa from the gastræa. The first rudiments of the principal parts




FIG. 46.—Very young human embryo of the fourth week, one-fourth of an inch long (taken from the womb of a female eight hours after death). (Front limb.) a, dorsal pole; b, eye; c, lower jaw; d, e, gill-arches; f, g, hind and fourth gill-arch; h, heart; i, primitive notochord; j, fore-limb (arm); k, hind-limb (leg); l, between the two the ventral pole.

we consider still earlier stages of development, we were unable to discover any essential difference in bodily structure between the embryos of these higher vertebrates and those of the lower, the amphibia and fishes. If, in fine, we go back to the construction of the body out of the four germinal layers, we are astonished to perceive that these four layers are the same in all vertebrates, and everywhere take a similar part in the building-up of the fundamental organs of the body. If we go back to the origin of these four secondary layers, we learn that they always arise in the same way from the two primary layers; and the latter have the same significance in all the metazoa (i.e., all animals except the

of the body, especially the oldest organ, the alimentary canal, are the same everywhere, they have always the same extremely simple form. All the peculiarities that distinguish the various groups of animals from each other only appear gradually in the course of embryonic development; and the closer the relation of the various groups, the later they are found. We may formulate this as a law, which may be regarded as an appendix to our biogenetic law. This is the law of the ontogenetic connection of related animal forms. It runs: The closer the

relation of two fully-developed animals in respect of their whole bodily structure, and the nearer they are connected in the classification of the animal kingdom, the longer do their embryonic forms retain their identity, and the longer is it impossible (or only possible on the ground of subordinate features) to distinguish between their embryos. This law applies to all animals whose embryonic development is, in the main, an hereditary summary of their ancestral history, or in which the original form of development has been faithfully preserved by heredity. When, on the other hand, it has been altered by cenogenesis, or disturbance

characteristic formation of its membrane (*zona pellucida*), which clearly distinguishes it from the ovum of all other animals. When the human fetus has attained the age of fourteen days, it forms a round vesicle (or "embryonic vesicle") about a quarter of an inch in diameter. A thicker part of its border forms a simple saucer-shaped embryonic shield one-twelfth of an inch long (Fig. 133). On its dorsal side we find in the middle line the straight medullary furrow, bordered by the two parallel dorsal or medullary swellings. Behind, it passes by the neurorhence canal into the primitive gut or primitive groove. From this the folding of the two coelenteropouches proceeds in the same way as in the other mammals (cf. Figs. 98, 99). In the middle of the saucer-shaped embryonic shield the first primitive segments immediately begin to make their appearance. At this age the human embryo cannot be distinguished from that of other mammals, such as the hare or dog.

Later (for after the twenty-first day) the human embryo has doubled its length; it is $\frac{1}{5}$ th of an inch



FIG. 134.—Human embryo at the middle of the fifth week, one-third of an inch long. (From Ruedi's Lecture as in Fig. 133, except at curve of skull, at upper jaw, at neck-indentation.)

of development, we find a limitation of the law, which increases in proportion to the introduction of new features by adaptation (cf. Chapter I., pp. 4-6). Thus the apparent exceptions to the law can always be traced to cenogenesis.


When we apply to man this law of the ontogenetic connection of related forms, and run rapidly over the earliest stages of human development with an eye to it, we notice first of all the structural identity of the ovum in man and the other mammals at the very beginning (Figs. 1, 14). The human ovum possesses all the distinctive features of the ovum of the viviparous mammals, especially the

the three higher sense-organs, and the rudiments of the gill-clefts, which pierce the sides of the neck (Fig. 179, 181). The allantois has grown out of the gut behind. The embryo is already entirely enclosed in the amnion, and is only connected in the middle of the belly by the vitelline duct with the embryonic vesicle, which changes into the yolk-sac. There are no extrusions or limbs at this stage, no trace of arms or legs. The head-end has been strongly differentiated from the tail-end, and the first outlines of the cerebral vesicles in fr below, under this form more or less clearly see no real face. Moreover at this stage a special character that distinguishes the human embryo from that of other mammals.

A week later (after the fourth week, or the twenty-eighth to thirtieth day development) the human embryo is

reached a length of about one-third of an inch (Fig. 179, IV). We can now clearly distinguish the head with its various parts, inside it the five primitive cerebral vesicles (fore-brain, middle-brain, intermediate-brain, hind-brain, and after-brain); under the head the gill-arches, which divide the gill-clefts; at the sides of the head the rudiments of the eyes, a couple of pits in the outer skin, with a

head bend over the trunk, almost at a right angle. The latter is still connected in the middle of its ventral side with the embryonic vesicle; but the embryo has still further severed itself from it, so that it already hangs out as the yolk-sac. The hind part of the body is also very much curved, so that the pointed tail-end is directed towards the head. The head and face-part are sunk entirely on the



— — — Median longitudinal section of the tail of a human embryo, two-thirds of an *Anguilla* long. (From *Atlas Generalis Anatomiae*.) Med, median longitudinal section; Cafil, caudal filament; Ca, caudal artery; Sp, spinal vein; An, anus; Sag, sagittal sinus; Vag, vaginal sinus.

pair of corresponding simple vesicles growing out of the lateral wall of the fore-brain (Figs. 180, 181 a). Far behind the eyes, over the last gill-arches, we see the vesicular rudiment of the muscular organ. The rudimentary limbs are now clearly outlined—four simple buds of the shape of round plates, a pair of fore (*fg*) and a pair of hind legs (*hg*), the former a little larger than the latter. The large

still open breast. The bend soon increases so much that the tail almost touches the forehead (Fig. 179, V.; Fig. 181). We may then distinguish three or four special curves on the round dorsal surface—namely, a skull-curve in the region of the second cerebral vesicle, a neck-curve at the beginning of the spinal cord, and a tail-curve at the fore-end. This pronounced curve is only shared by man and

the higher classes of vertebrates (the annulides); it is much slighter, or not found at all, in the lower vertebrates. At this age (four weeks) man has a considerable tail, twice as long as his legs. A vertical longitudinal section through the middle plane of this tail (Fig. 182) shows that the hinder end of the spinal marrow extends to the point of the tail,

or *arteria sacralis media*, &c.), and the principal vein (*vena cava inferior* or *sacralis media*). Underneath is the opening of the anus (*an*) and the urogenital slits (*5 ug*). From this anatomic structure of the human tail it is perfectly clear that



FIG. 182.—Human embryo, four weeks old, opened on the ventral side. Ventral and dorsal walls are cut away, so as to show the contents of the pectoral and abdominal cavities. All the appendages are also removed (intestine, yellowish) and the middle part of the gut. *n* eye, *g* nose, *a* upper jaw, *g* lower jaw, *g* second, *g* third gill-arch, *ov* heart (*g* right, *g* left auricle, *g* right, *g* left ventricle), *g* origin of the aorta, *g* liver (in umbilical vein), *g* gut (with vitelline artery, cut off at *g*), *g* vitelline vein, *m* primitive kidneys, *g* rudimentary nasal glands, *g* terminal gut (cut off at the secondary *g*), *u* umbilical artery, *u* umbilical vein, *g* fore-leg, *g* hind-leg. (From Cuvier.)

as also does the underlying chorda (*ch*), the terminal continuation of the vertebral column. Of the latter, the rudiments of cervical (or lowest) vertebrae are visible—thirty-two indicates the third and thirty-six the seventh of these. Under the vertebral column we see the hindmost ends of the two large blood-vessels of the tail, the principal artery (*aorta caudalis*

FIG. 183.—Human embryo, five weeks old, opened from the ventral side (as in Fig. 182). Broad and hollowed and liver are removed. *g* outer nasal process, *g* upper jaw, *g* lower jaw, *g* tongue, *g* right, *g* left ventricle of heart, *g* left auricle, *g* origin of aorta, *g*, *g*, *g* first, second, and third aortic-arches, *g*, *g*, *g* venae cavae, *g* lungs (*g* pulmonary artery), *g* kidneys, *m* primitive kidneys (*g* left vitelline vein, *g* vitelline vein, *g* right vitelline artery, *u* umbilical artery, *u* umbilical vein), *g* vitelline duct, *g* rectum, *g* hind, *g* fore-leg, *g* hind-leg. (From Cuvier.)

it is the rudiment of an upo-tail, the last hereditary relic of a long bi-handled down to primate ancestors to the present day.

It sometimes happens that we find even external relics of this tail growing. According to the illustrated works of

Surgeon-General Bernhard Ornstein, of Greece, these tailed men are not uncommon; it is not impossible that they gave rise to the ancient fables of the satyrs. A great number of such cases are given by Max Bartels in his essay on "Tailed Man" (1884, in the *Archiv für Anthropologie*, Band XV), and critically examined. These atavistic human tails are often mobile; sometimes they contain only muscles and fat, sometimes

tion is hereditary in certain isolated tribes (especially in south-eastern Asia and the



1 John Pasternak



FIG. 184.—Human ovum of twelve to thirteen days (?). (From Allen Thomson.) a Not opened, natural size. b Opened and magnified. Within the latter chorion the tiny curved ligatus loop on the large embryonic vessels, to the left above.

rudiments of caudal vertebrae. They attain a length of eight to ten inches and more. Granville Harrison has very carefully studied one of these cases of "pig-tail," which he removed by operation from a six months' old child in 1901. The tail moved bristly when the child cried or was excited, and was drawn up when at rest.

In the opinion of some travellers and anthropologists, the atavistic tail-forma-



FIG. 185

FIG. 186

FIG. 185.—Human ovum of ten days. (From Allen Thomson.) Natural size, opened, the small foramen in the right half, above.

FIG. 186.—Human ovum of ten days, taken from the preceding ovum, magnified ten times. a yolk-sac; b neck (the medullary groove already closed); c head



FIG. 186a.—Human ovum of twenty to twenty-two days. (From Allen Thomson.) Natural size, opened, the chorion foramen a spacious vesicle, to the right wall of which the small foramen (to the right above) is attached by a short umbilical cord.



FIG. 186b.—Human ovum of twenty to twenty-two days, taken from the preceding ovum, magnified a hundred times. a lower jaw process of the first gill-arch, b upper jaw process of same, c second gill-arch (two smaller ones behind). Three gill-slits are clearly seen. d rudimentary forelimb, e rudimentary vesicle, f eye, g heart.

archipelago), so that we might speak of a special race or "species" of tailed man.

(*Homo canis*) Bartels has "no large, and almost fills the whole of the (Fig. 183, *ov*). Behind it small rudimentary lungs. The primitive kidneys (*an*) are very large, they fill the greater part of the abdominal cavity, and extend from the liver (*f*) to

physical and ethnographical knowledge of the kinds in question" (*Lectures for Anthropologic*, Band XV., p. 129)



FIG. 183.—Human embryo of sixteen to eighteen days. (From Cuvier.) Magnified. The embryo is surrounded by the amnion (*a*) and has been with this in the upturned embryonic vesicle. The belly is drawn up by the large gut-sack (*g*), and fastened to the inner wall of the embryonic membrane by the short and thick pedicle (*h*). Hence the normal convex curve of the back (Fig. 184) is here changed into an abnormal concave surface. *b* heart, *c* parallel discoidium. The spaces on the outer wall of the serotum are the roots of the branching chorionic villi, which are free at the border.

mouth (Fig. 183), we find the alimentary canal formed in the body-cavity, and for the most part cut off from the embryonic vesicle. There are both mouth and anus apertures. But the mouth-cavity is not yet separated from the nasal cavity, and the face not yet shaped. The heart shows all its four sections, it is very

first month all the chief organs are already outlined. But there are at this stage no features by which the human embryo materially differs from that of the dog, the hare, the ox, or the horse—in a word, of any other higher mammal. All these embryos have the same, or at least a very similar, form; they can at the most be

distinguished from the human embryo by the total size of the body or some other insignificant difference in size. Thus, for instance, in man the head is larger in proportion to the trunk than in the ox.

The features, by means of which we distinguish between them are not clear until later on. Even at a much more advanced stage of development, when we can distinguish the human foetus from that of



FIG. 164.—Human embryo of the fourth week, one-third of an inch long, lying in the dissected chorion.

he tail is rather longer in the dog than in man. These are all negligible differences. On the other hand, the whole internal organisation and the form and arrangement of the various organs are essentially the same in the human embryo of four weeks as in the embryos of the other mammals at corresponding stages.

It is otherwise in the second month of human development. Fig. 179 represents a human embryo of six weeks (VI), one of seven weeks (VII), and one of eight weeks (VIII), at natural size. The differences which mark off the human embryo from that of the dog and the lower mammals now begin to be more pronounced. We can see important differences at the sixth, and still more at the eighth, week, especially in the formation of the head. The size of the various sections of the brain is greater in man, and the tail is shorter. Other differences between man and the lower mammals are found in the relative size of the internal



of the nearest related mammals—the dog, especially the anthropomorphic ape. he angulates at a glance, it still closely resembles the last we got the distinctive features, and

we can distinguish the human embryo confidently at the first glance from that of all other mammals during the last four months of fetal life—from the sixth to the ninth month of pregnancy. Then we begin to find also the differences between the various races of men, especially in regard to the formation of the skull and the face. (Of Chapter XXIII.)

The striking resemblance that persists so long between the embryo of man and of the higher apes disappears much earlier in the lower apes. It naturally remains

famous Miss Julia Pastrana, Fig. 185), it will be admitted to represent a higher stage of development. There are still people among us who look especially to the face for the "image of God in man." The long-nosed ape would have more claim to this than some of the stumpy-nosed human individuals one meets.

This progressive divergence of the human from the animal form, which is based on the law of the ontogenetic connection between related forms, is found in the structure of the internal organs as well




FIG. 184.—Human embryo with its membranes, six weeks old. The outer cavity of the womb even at the chorion, thickly covered with its branching villi, a product of the syncytial membrane. The embryo is retracted in its dorsal position. The yolk-sac is reduced to a small pear-shaped umbilical vesicle, its thin pedicle, the long vitelline duct, is enclosed in the embryonic cord. In the latter, behind the vitelline duct is the much shorter pedicle of the allantois, the inner lamina of which (the peritoneal layer) forms a long allantoic tube. The outer lamina is attached to the inner wall of the outer embryonic coat, and forms the placenta there. (Holt diagrammatic.)

longest in the large anthropomorphic apes (gorilla, chimpanzee, orang, and gibbon). The physiognomic similarity of these animals, which we find so great in their earlier years, lessens with the increase of age. On the other hand, it remains throughout life in the remarkable long-nosed ape of Borneo (*Nasalis larvatus*). Its finely-shaped nose would be regarded with envy by many a man who has too little of that organ. If we compare the face of the long-nosed ape with that of abnormally ape-like human beings (such as the

as in external form. It is also expressed in the construction of the envelopes and appendages that we find surrounding the fetus externally, and that we will now consider more closely. Two of these appendages—the amnion and the allantois—are only found in the three higher classes of vertebrates, while the third, the yolk-sac, is found in most of the vertebrates. This is a circumstance of great importance, and it gives us valuable data for constructing man's genealogical tree.

As regards the external membrane that encloses the ovum in the mammal womb,

we find it just the same in man as in the higher mammals. The ovum is, the reader will remember, first surrounded by the transparent structureless *coelomus* or *zona pellucida* (Figs. 1, 14). But very soon, even in the first week of development, this is replaced by the permanent chorion. This is formed from the external layer of the amnion, the *scrolemma*, or "serous membrane," the formation of which we shall consider presently; it surrounds the foetus and its appendages as a broad, completely-closed sac; the space between the two, filled with clear watery fluid, is the *serotocœlum*, or *interamniotic cavity*,

to one-third of an inch in diameter (Figs. 186-188). As a large quantity of fluid gathers inside it, the chorion expands more and more, so that the embryo only occupies a small part of the space within the vesicle. The villi of the chorion grow larger and more numerous. They branch out more and more. At first the villi cover the whole surface, but they afterwards disappear from the greater part of it, they then develop with proportionately greater vigour at a spot where the placenta is formed from the allantois.

When we open the chorion of a human embryo of three weeks, we find on the




FIG. 191.—Diagram of the embryonic organs of the mammal (fetal membranes and appendages). (From Turner). *A*, *U*, *H* outer, middle, and inner germ layers of the embryonic shell, which is figured in median longitudinal section. *ALC*, allantois.

("extra-embryonic body-cavity"). — The smooth surface of the sac is quickly covered with numbers of tiny tufts, which are like the villi of the placenta (Figs. 192, 198 *chr*).

ponding depressions that are formed by the tubular glands of the mucous membrane of the maternal womb. Thus, the ovum secures its permanent seat (Figs. 186-194).

In human ova of eight to twelve days this external membrane, the chorion, is already covered with small tufts or villi, and forms a ball or spheroid of one-fourth

antral side of the foetus a large sac, filled with fluid. This is the *umbilical vesicle*, the origin of which we have considered previously. The larger the embryo becomes the smaller we find the yolk-sac. In the end we find the remainder of it in the shape of a small pear-shaped vesicle, fastened to a long thin stalk (or pedicle), and hanging from the open belly of the foetus (Fig. 194). This pedicle is the *vitellic duct*, and is separated from the body at the closing of the navel.

Behind the yolk-sac a second appendage,

In this case there is at an early stage an intimate blending of the chorion of the embryo and the part of the mucous lining of the womb to which it attaches. The villi of the chorion with the blood-vessels they contain grow so completely into the tissue of the uterus, which is rich in blood, that it becomes impossible to separate them, and they form together a sort of cake. This comes away as the "afterbirth" at parturition, at the same time, the part of the mucous lining of the womb that has united inseparably with the chorion is torn away, hence it is called the *decidua* ("falling-away mem-

brana")—namely, that part of the mucous lining of the womb which unites intimately with the chorion-villi of the foetal placenta. The internal or false decidua (*interna* or *reflexa*, Fig. 196 *dr*, Fig. 199 *f*) is that part of the mucous lining of the womb which encloses the remaining surface of the ovum, the smooth chorion (*chorion laeve*), in the shape of a special thin membrane. The origin of these three different deciduous membranes, in regard to which quite erroneous views (still retained in their names) formerly prevailed, is now quite clear. The external decidua vera is the specially modified and subsequently




FIG. 199.—Male embryo of the flowering-gibbon (*Macaca leonina*) of Sumatra, very much natural size. In the left the dissected uterus, of which only the dorsal half is given. The embryo has been taken out, and the limbs folded together, it is still connected by the umbilical cord with the center of the decidua placenta, which is attached to the inside of the uterus. This embryo takes the head-position in the womb, and thus is normal in that also.

brane"), and also the "decidua-membrana," because it is perforated like a sieve. We find a decidua of this kind in most of the higher placentals, but it is only in man and the anthropoid apes that it divides into three parts—the outer, inner, and placental decidua. The external or true decidua (Fig. 196 *dr*, Fig. 199 *f*) is the part of the mucous lining of the womb that clothes the inner surface of the uterine cavity wherever it is not connected with the placenta. The placental or spongy decidua (*placentalis* or *serotina*, Fig. 196 *dr*, Fig. 199 *d*) is really the placenta itself, or the maternal part of it (*placenta*

detachable superficial stratum of the original mucous lining of the womb. The placental decidua *serotina* is that part of the preceding which is completely transformed by the ingrowth of the chorion-villi, and is used for constructing the placenta. The inner decidua *reflexa* is formed by the rise of a circular fold of the mucous lining (at the border of the decidua vera and serotina), which grows over the foetus (like the amnion) to the

The peculiar anatomic features that characterise the human foetal membrane are found in just the same way in the higher

FETAL MEMBRANES AND CIRCULATION

apes. Until recently it was thought that the human embryo was distinguished by its peculiar construction of a solid allantois and a special ventral pedicle, and that the umbilical cord developed from this in a different way than in the other mammals. The opponents of the unwelcome "ape-theory" laid great stress on this, and thought they had at last discovered an important indication that separated man from all the other placental. But the

described the allantois has no blood-vessels at any moment of its existence. But the other two vesicles, the yolk-sac and the allantois, are equipped with large blood-vessels, and these effect the nourishment of the embryonic body. We may take the opportunity to make a few general observations on the first circulation in the embryo and its central organ, the heart. The first blood-vessels, the heart, and the first blood itself, are formed from the




FIG. 104.—Frontal section of the pregnant human womb. (If I hurry in the middle of the amniotic cavity by the ventral pedicle of the allantois.)


remarkable discoveries published by the distinguished zoologist Seale in 1890 proved that man shares these peculiarities of placentation with the anthropoid apes, though they are not found in the other apes. Thus the very feature which was advanced by our critics as a disproof became a most important piece of evidence in favour of our pithecolid origin.

Of the three vesicular appendages of the amniotic embryo which we have now

gut-fibre layer. Hence it was called by earlier embryologists the "vascular layer." In a sense the term is quite correct. But it must not be understood as if all the blood-vessels in the body came from this layer, or as if the whole of this layer were taken up only with the formation of blood-vessels. Neither of these suppositions is true. Blood-vessels may be formed independently in other parts, especially in the various products of the skin-fibre layer.



FIG. 149.—Human fetus, twelve weeks old, with its membranes. Natural size. The umbilical cord is pulled from its canal to the placenta, & amnion & chorion, & placenta & folds of tube on smooth surface decidual. (From R. du Roidar.)



—Mature human fetus (at the end of pregnancy, in its natural position, taken out of the uterus the lower surface of the uterus (on the left) at the placenta, which is connected by the umbilical cord with the child's navel. (From Bernhard Schmidt.)

The first blood-vessels of the mammal embryo have been considered by us previously, and we shall study the development of the heart in the second volume.

In every vertebrate it lies at first in the ventral wall of the fore-gut, or in the ventral (or cardiac) mesentery, by which it is connected for a time with the wall of the body. But it soon severs itself from the place of its origin, and lies freely in a cavity—the cardiac

They rise in the wall of the fore-gut, which they enclose in a sense, and then unite above, in the upper wall of the fore-gut-cavity, to form a large single artery, that runs backward immediately under the chorda, and is called the aorta (Fig. 202 *As*). The first pair of aorta-arches rise on the inner wall of the first pair of gill-arches, and so lie between the first gill-arch (*h*) and the fore-gut (*d*), just as we find them throughout life in the




FIG. 202.—Vitalis vessels in the germinal area of a chick-embryo, at the close of the third day of incubation. (From *Reichert*.) The detached germinal area is seen from the ventral side. The notochord is dark, the nerve light. H, heart. A, aorta. A, aorta. R, Q, right and left cephalic vessels. A, aorta. S, C, ventral branches. L, D, and P, Q, right and left cephalic-vertebral vessels. S, C, ventral branches. A, C, dorsal branches. S, C, and P, Q, fore and hind cerebral vessels.

cavity. For a short time it is still connected with the former by the thin plate of the mesocardium. Afterwards it lies quite free in the cardiac cavity, and is only directly connected with the gut-wall by the vessels which issue from it.

The fore-end of the spindle-shaped tube, which soon bends into an S-shape (Fig. 203), divides into a right and left branch. These tubes are bent upwards arch-wise, and represent the first arches of the aorta.

fishes. The single aorta, which results from the conjunction of these two first vascular arches, divides again immediately into two parallel branches, which run backwards on either side of the chorda. These are the primitive aortae which we have already mentioned; they are also called the posterior vertebral arteries. These two arteries now give off at each side, behind, at right angles, four or five branches, and these pass from the embryonic body to the germinal area; they

are called *omphalo-mesenteric* or *vitelline* arteries. They represent the first beginning of a *fœtal* circulation. Thus, the first blood-vessels pass over the embryonic body and reach as far as the edge of the germinal area. At first they are inclined to the dark or "vascular" area. But they

of the embryonic vesicle. In this end, the whole of the yolk-sac is covered with a vascular net-work. These vessels rather food for

the yolk-sac and convey it to the embryonic body. This is done by the veins, which pass first from the germinal area, and afterwards from the yolk-sac, to the farther end of the heart. They are called *vitelline*, or, frequently, *omphalo-mesenteric*, veins.

These vessels naturally atrophy with the degeneration of the umbilical vesicle, and the vitelline circulation is replaced by a second, that of the allantois. Large blood-vessels are developed in the wall of the urinary sac or the allantois, as before, from the gut-fibre layer. These vessels grow larger and larger, and are very closely connected with the vessels that develop in the body of the embryo itself. Thus, the secondary, allantoic circulation gradually takes the place of the original vitelline circulation. When the allantois has attached itself to the inner wall of the chorion and been converted into the placenta, its blood-vessels alone effect the nourishment of the embryo. They are called *umbilical* vessels, and are originally double—a pair of umbilical arteries and a pair of umbilical veins. The two umbilical veins (Fig. 183 a), which convey blood from the placenta to the heart, open at first into the united vitelline veins. The latter then disappear, and the right umbilical vein goes with them, so that henceforth a single large vein, the left umbilical vein, conducts all the blood from the placenta to the heart of the embryo. The two arteries of the allantois, or the umbilical arteries (Figs. 183 a, 184 a), are merely the ultimate terminations of the primitive aorta, which are strongly developed afterwards. This umbilical circulation is retained until the nine months of embryonic life are over, and the human embryo enters into the world as an independent individual. The umbilical cord (Fig. 196 af), in which these large blood-vessels pass from the embryo to the placenta, comes away, together with the latter, in the after-birth, and

with the use of the lungs begins an entirely new form of circulation, which is confined to the body of the infant.

There is a great phylogenetic significance in the perfect agreement which we find between man and the anthropoid apes in these important features of . . . bryonic circulation, and the special construction of the placenta and the umbilical cord. We must infer from it a close blood-relationship of man and the anthropomorphic apes—a common descent of them from one and the same extinct



FIG. 183. — Spot-shaped embryo of the dog, from the ventral side, magnified about ten times. Is first under the forehead we then see the first pair of arteries; underneath is the S-shaped heart, at the sides of which are the secondary veins. The lower part of the embryo is the yolk-sac, which is still attached.

See page 171. The two pairs of vitelline arteries are given off. (From Huxley's "Lessons in Comparative Zoology.")

group of lower apes. Huxley's "pithoco-metra-principle" applies to these ontogenetic features as much as to any other morphological relations: "The differences in construction of any part of the body are less between man and the anthropoid apes than between the latter and the lower apes."

This important Huxleyian law, the chief consequence of which is "the descent of man from the ape," has lately been confirmed in an interesting and unexpected way from the side of the experimental

physiology of the blood. The experiments of Hans Friedenthal at Berlin have shown that human blood, mixed with the blood of lower apes, has a poisonous

As we know that the blood is only possible without injury in the case of two closely related animals of



FIG. 105.—LAR or white-headed gibbon (*Hylobates lar* or *allmanni*), from the Indian mainland (From Becken)

effect on the latter, the serum of the one destroys the blood-cells of the other. But this does not happen when human blood is mixed with that of the anthropoid ape.

In the same family, we have another proof of the close blood-relationship, in the literal sense of the word, of man and the anthropoid ape.

The existing anthropoid apes are only a small remnant of a large family of eastern apes (or *Catarrhinae*), from which man was evolved about the end of the Tertiary period. They fall into two groups

eight to twelve species of it in the East Indies. I made observations of four of them during my voyage in the East Indies (1901), and had a specimen of the ash-grey gibbon (*Hylodates leuciscus*) living



FIG. 204.—YOUNG ORANG (*Setyrus orang*), sitting.

graphical groups—the Asiatic and the African anthropoids. In each group we can distinguish two genera. The subject of these four genera is the gibbon (*Hylodates*, FIG. 203); there are from

for several months in the garden of my house in Java. I have described the interesting habits of this ape (regarded by the Malays as the wild descendant of men who had lost their way) in my *Malayische*

Ressetrisjen (chap xi). Psychologically, he showed a good deal of resemblance to the children of my Malay hosts, with whom he played and formed a very close friendship.

The second, larger and stronger, genus of Asiatic anthropoid ape is the orang (*Satyrus*), he is now found only in the islands of Borneo and Sumatra. Selenka,

liar and salient cheek-pads in the elderly male; these are wanting in the other group, the ordinary orang-outang (*Esau/orus*).

Several species have lately been distinguished in the two genera of the black African anthropoid apes (chimpanzee and gorilla). In the genus *Anthropithecus* (or *Anthropopithecus*, formerly) *Troglodytes*,



FIG. 205.—WILD ORANG (*Dasyrctes sumatra*). (From R. Fock and Latschmann.)

who has published a very thorough *Study of the Development and Cranial Structure of the Anthropoid Apes* (1899), distinguishes ten races of the orang, which may, however, also be regarded as "local varieties or species." They fall into two sub-genera or genera. One group, *Dasyrctes* (orang-batang, Fig. 205), is distinguished for the strength of its limbs, and the formation of very pecu-

liar and salient cheek-pads in the elderly male; these are wanting in the other group, the ordinary orang-outang (*Esau/orus*). Several species have lately been distinguished in the two genera of the black African anthropoid apes (chimpanzee and gorilla). In the genus *Anthropithecus* (or *Anthropopithecus*, formerly) *Troglodytes*, the bald-headed chimpanzee, *A. calvus* (Fig. 206), and the gorilla-like *A. majus* differ very strikingly from the ordinary *Anthropithecus aeyer* (Fig. 207), not only in the size and proportion of many parts of the body, but also in the peculiar shape of the head, especially the ears and lips, and in the hair and colour. The controversy that still continues as to whether these different forms of



FIG. 104. — The bald-headed shimpansze (*Anthrophormus catenax*). From a series of specimens described by Frank Redford in 1897 as *Trogodytes cavens*, differs considerably from the ordinary *A. niger* (Fig. 107) in the structure of the head, the colouring, and the absence of hair in parts.

chimpanzees and orang are "merely local varieties" or "true species" as an idle one, as in all such disputes of classifiers there is an utter absence of clear ideas as to what a species really is.

Of the largest and most famous of all the anthropoid apes, the gorilla, Paschen has lately discovered a giant-form in the interior of the Cameroons, which seems to differ from the ordinary species (*Gorilla*

to that of man, but it is substantially the same. "The same 200 bones, arranged in the same way, form our internal skeleton; the same 300 muscles effect our movements, the same hair covers our skin, the same groups of ganglionic cells compose the ingenious mechanism of our brain; the same four-chambered heart is the central pump of our circulation." The really existing differences in the



FIG. 207.—Female chimpanzee (*Anthropus niger*). (From Rehn.)

gila Fig. 208), not only by its unusual size and strength, but also by a special formation of the skull. This giant gorilla (*Gorilla gila*, Fig. 209) is six feet eight inches long; the span of its great arms is about nine feet; its powerful chest is twice as broad as that of a strong man.

The whole structure of this huge anthropoid ape is not merely very similar

shape and size of the various parts are explained by differences in their growth, due to adaptation to different habits of life and unequal use of the various organs. This of itself proves morphologically the descent of man from the ape. We will return to the point in the twenty-third chapter. But I wanted to point already to this important solution of "the question of questions," because that agreement

FETAL MEMBRANES AND CIRCULATION

in the formation of the embryonic membranes and in fetal circulation which I have described affords a particularly weighty proof of it. It is the more instructive as even ontogenetic structures

may in certain circumstances have a high phylogenetic value. In conjunction with the other facts, it affords a striking confirmation of our biogenetic law.



Fig. 106.—Female gorilla. (From Hensley.)



Fig. 10. Male giant gorilla (*Gorilla silver*), from Yavondo, in the interior of the Caprivi. Killed by
Kauwara, killed by O. K. K.

THE EVOLUTION OF MAN

A POPULAR SCIENTIFIC STUDY

ERNST HAECKEL

VOL. II.

HUMAN STEM-HISTORY, OR PHYLOGENY

AND BY JOSEPH McCABE

WATTS & Co.,
17, JOHNSON'S COURT, FLEET STREET, LONDON, E.C.

1923

*Printed (or made) in Great Britain
by Watts & Co., Johnson's Court, Fleet Street,
London, E C 4*

CONTENTS OF VOL. II.

	PAGE
LIST OF ILLUSTRATIONS	v
INDEX	359
<hr/>	
CHAP	
XVI. STRUCTURE OF THE LANCELET AND THE SEA-SQUID	179
XVII. EMBRYOLOGY OF THE LANCELET AND THE SEA-SQUID	191
XVIII. DURATION OF THE HISTORY OF OUR SPECIES	199
XIX. OUR PROTIST ANCESTORS	207
XX. OUR WORM-LIKE ANCESTORS	218
XXI. OUR FISH-LIKE ANCESTORS	229
XXII. OUR FIVE-TOED ANCESTORS	239
XXIII. OUR APE ANCESTORS	253
XXIV. EVOLUTION OF THE NERVOUS SYSTEM	266
XXV. EVOLUTION OF THE SENSE-ORGANS	280
XXVI. EVOLUTION OF THE ORGANS OF MOVEMENT	292
XXVII. EVOLUTION OF THE ALIMENTARY SYSTEM	308
XXVIII. EVOLUTION OF THE VASCULAR SYSTEM	318
XXIX. EVOLUTION OF THE SEXUAL ORGANS	330
XXX. RESULTS OF ANTHROPOGENY	348

LIST OF ILLUSTRATIONS

210. The lancelet	181	260. Fossil amphibian	240
211. Section of the head of the lancelet	181	261. Larva of the spotted salamander	241
212. Section of an amphioxus-larva	184	262. Larva of common frog	241
213. Diagram of preceding	184	263. Fossil mailed amphibian	243
214. Section of a young amphioxus	185	264. The New Zealand lizard	243
215. Diagram of preceding	185	265. <i>Homocephalus pulchellus</i>	246
216. Transverse section of lancelet	186	266. Skull of a Permian lizard	247
217. Section through the middle of the lancelet	187	267. Skull of a therapsid	248
218. Section of a primitive-fish embryo	187	268. Lower jaw of a primitive mammal	249
219. Section of the head of the lancelet	188	269-270. The ornithomimus	250
220-221. Organisation of an ascidian	189	271. Lower jaw of a promammal	251
222-224. Sections of young amphioxus-larva	190	272. The crab-eating opossum	252
225. An appendiculus	196	273. Fetal membranes of the human embryo	254
226. <i>Chroococcus minor</i>	197	274. Skull of a fossil lemur	255
227. <i>Applanocapsa perimorialis</i>	209	275. The slender loris	256
228. <i>Protospongia</i>	210	276. The white-nosed ape	257
229. Original ovum-cleavage	211	277. The drill-baboon	258
230. Morula	212	278-281. Skeletons of man and the anthropoid apes	260
231-232. <i>Magnosphera planula</i>	214	282. Skull of the Java ape-man	263
233. Modern gastrula	216	284. Section of the human skin	267
234-235. <i>Prophyria perimorialis</i>	217	285. Epidermic cells	268
236-237. Aculae of <i>Gastrophyscus</i>	217	286. Rudimentary lachrymal glands	269
238. <i>Olynthus</i>	218	287. The female breast	270
239. <i>Aphanostomus</i> Langu	222	288. Mammary gland of a new-born infant	271
240-241. A turbellarian	223	289. Embryo of a bear	272
242-243. <i>Chaetomus</i>	224	290. Human embryo	273
244. A nemertine worm	225	291. Central marrow of a human embryo	273
245. An enteropneust	225	292-293. The human brain	274-275
246. Section of the branchial gut	226	294-296. Central marrow of human embryo	276
247. The marine lamprey	230	297. Head of a chick embryo	277
248. Fossil primitive fish	231	298. Brain of three craniole embryos	277
249. Embryo of a shark	232	299. Brain of a shark	277
250. Man-eating shark	233	300. Brain and spinal cord of a frog	278
251. Fossil angel-shark	234	301. Brain of an ox-embryo	278
252. Tooth of a gigantic shark	235	302-303. Brain of a human embryo	278-279
253-255. <i>Crossopterygii</i>	236	304. Brain of the rabbit	279
256. Fossil dipneust	237		
257. The Australian dipneust	237		
258-259. Young <i>ceratodus</i>	238		

	PAGE		PAGE
309. Head of a shark - - -	281	353-356. Sections of head of lamprey -	314
309-310. Heads of chick-embryos -	282	357. Viscera of a human embryo -	315
311. Section of mouth of human embryo - - - -	283	358. Red blood-cells - - - -	319
312. Diagram of mouth-nose cavity -	284	359. Vascular tissue - - - -	319
313-314. Heads of human embryos -	285	360. Section of trunk of a chick-embryo - - - -	320
315-316. Face of human embryo -	286	361. Microcytes - - - -	321
317. The human eye - - - -	287	362. Vascular system of an animal -	322
318. Eye of the chick embryo - -	288	363. Head of a fish-embryo - -	322
319. Section of eye of a human embryo -	288	364-370. The five arterial arches -	323-324
320. The human ear - - - -	289	371-372. Heart of a rabbit-embryo -	325
321. The body labyrinth - - - -	289	373-374. Heart of a dog-embryo -	325
322. Development of the labyrinth -	290	375-377. Heart of a human embryo -	326
323. Primitive skull of human embryo -	291	378. Heart of adult man - - -	326
324. Rudimentary muscles of the ear -	291	379. Section of head of a chick-embryo - - - -	327
325-326. The human skeleton - -	293	380. Section of a human embryo -	328
327. The human vertebral column -	294	381-382. Sections of a chick-embryo -	329
328. Piece of the dorsal cord - -	294	383. Embryos of sagitta - - -	333
329-330. Dorsal vertebrae - - -	295	384. Kidneys of bledianoma - -	334
331. Intervertebral disk - - -	296	385. Section of embryonic shield -	335
332. Human skull - - - -	296	386-387. Primitive kidneys - -	336
333. Skull of new-born child - -	297	388. Pig-embryo - - - -	337
334. Head-skeleton of a primitive fish - - - -	297	389. Human embryo - - - -	337
335. Skulls of nine primates - -	298	390-392. Rudimentary kidneys and sexual organs - - - -	338
335-338. Evolution of the fin - -	299	393-394. Urinary and sexual organs of salamander - - -	338
339. Skeleton of the fore-leg of an amphibian - - - -	300	395. Primitive kidneys of human embryo - - - -	339
340. Skeleton of gorilla's hand - -	300	396-398. Urinary organs of ox-embryos - - - -	340
341. Skeleton of human hand - -	300	399. Sexual organs of water-mole -	341
342. Skeleton of hand of six mammals -	301	400-401. Original position of sexual glands - - - -	342
343-345. Arm and hand of three anthropoids - - - -	302	402. Urogenital system of human embryo - - - -	342
346. Section of fish's tail - - -	303	403. Section of ovary - - - -	343
347. Human skeleton - - - -	303	404-406. Graafian follicles - - -	343
348. Skeleton of the giant gorilla -	303	407. A ripe Graafian follicle - -	344
349. The human stomach - - -	303	408. The human ovum - - - -	345
350. Section of the head of a rabbit-embryo - - - -	310		
351. Shark's teeth - - - -	311		
352. Gut of a human embryo - -	312		
353-354. Gut of a dog embryo - -	313		

CHAPTER XVI.

STRUCTURE OF THE LANCELET AND THE SEA-SQUIRT

In turning from the embryology to the phylogeny of man—from the development of the individual to that of the species—

and the direct causal link that exists between these two man

evolution. This important finds its simplest expression in "the fundamental law of organic development," the content and purport of which we have fully considered in the first chapter. According to this biogenetic law, ontogeny is a brief and condensed recapitulation of phylogeny. If this compendious reproduction were complete in all cases, it would be very easy to construct the whole story of evolution on an embryonic basis. When we wanted to know the ancestors of any higher organism, and, therefore, of man—to know from what forms the race as a whole has been evolved—we should merely have to follow the series of forms in the development of the individual from the ovum; we could then regard each of the successive forms as the representative of an extinct ancestral form. However, this direct application of ontogenetic facts to phylogenetic ideas is possible, without limitations, only in a very small section of the animal kingdom. There are, it is true, still a number of lower invertebrates (for instance, some of the Zoophyta and Vermalis) in which we are justified in recognizing at once each embryonic form as the historical reproduction, or *effigie*, as it were, of an extinct ancestor. But in the great majority of the animals, and in the case of man, this is impossible, because the embryonic forms themselves have been modified through the change of the conditions of existence, and have lost their original character to some extent. During the immeasurable course of organic history, the many millions of years during which life was developing on our planet, sec-

dary changes of the embryonic forms have taken place in most animals. The young of animals (not only let larvae, but also the embryos enclosed in the womb) may be modified influence of the e

tion to the conditions of life; even species are altered during the embryonic development. Moreover, it is an advantage for all higher organisms (and the advantage is greater the more advanced they are) to curtail and simplify the original course of development, and thus to obliterate the traces of their ancestors. The higher the individual organism is in the animal kingdom, the less completely does it reproduce in its embryonic development the series of its ancestors, for reasons that are as yet only partly known to us. The fact is easily proved by comparing the different developments of higher and lower animals in any single stem.

In order to appreciate this important feature, we have distributed the embryological phenomena in two groups, *palinogenetic* and *anagenetic*. Under *palinogenesis* we count those facts of embryology that we can directly regard as a faithful synopsis of the corresponding stem-history. By *anagenesis* we understand those embryonic processes which we cannot directly correlate with corresponding evolutionary processes, but must regard as modifications or falsifications of them. With this careful discrimination between *palinogenetic* and *anagenetic* phenomena, our biogenetic law assumes the following more precise shape: rapid and brief development of the individual (ontogeny) is a condensed synopsis of the long and slow history of the stem (phylogeny): this synopsis is the faithful and complete in proportion as the original features have been preserved by heredity, and modifications have not been introduced by adaptation.

In order to distinguish correctly between palingenetic and cenogenetic phenomena in embryology, and deduce sound conclusions in connection with stem-history, we must especially make a comparative study of the former. In doing this it is best to employ the methods that have long been used by geologists for the purpose of establishing the succession of the sedimentary rocks in the crust of the earth. Thus solid crust, which encloses the glowing central mass like a thin shell, is composed of different kinds of rocks: there are, firstly, the volcanic rocks which were formed directly by the cooling at the surface of the molten mass of the earth; secondly, there are the sedimentary rocks, that have been made out of the former by the action of water, and have been laid in successive strata at the bottom of the sea. Each of these sedimentary strata was at first a soft layer of mud, but in the course of thousands of years it condensed into a solid, hard mass of stone (sandstone, limestone, marl, etc.), and at the same time permanently preserved the solid and imperishable bodies that had chanced to fall into the soft mud. Among these bodies, which were either fossilized or left characteristic impressions of their forms in the soft slime, we have especially the more solid parts of the animals and plants that lived and died during the deposit of the silty strata.

Hence each of the sedimentary strata has its characteristic fossils, the remains of the animals and plants that lived during that particular period of the earth's history. When we make a comparative study of these strata, we can survey the whole series of such periods. All geologists are now agreed that we can demonstrate a definite historical succession in the strata, and that the lowest of them were deposited in very remote, and the uppermost in comparatively recent, times. However, there is no part of the earth where we find the series of strata in its entirety, or even approximately complete. The succession of strata and of corresponding historical periods generally given in geology is an ideal construction, formed by piecing together the various partial discoveries of the succession of strata that have been made at different points of the earth's surface (cf. Chapter XVIII.).

We must act in this way in constructing the phylogeny of man. We must try to piece together a fairly complete picture of

the series of our ancestors from the various phylogenetic fragments that we find in the different groups of the animal kingdom. We shall see that we are really in a position to form an approximate picture of the evolution of man and the mammals by a proper comparison of the embryology of very different animals—a picture that we could never have framed from the ontogeny of the mammals alone. As a result of the above-mentioned cenogenetic processes—those of disturbed and curtailed heredity—whole series of lower stages have dropped out in the embryonic development of man and the other mammals, especially from the earliest periods, or been falsified by modification. But we find these lower stages in their original purity in the lower vertebrates and their invertebrate ancestors. Especially in the lowest of all the vertebrates, the lancelet or *Amphioxus*, we have the oldest stem-forms completely preserved in the embryonic development. We also find important evidence in the fishes, which stand between the lower and higher vertebrates, and throw further light on the course of evolution in certain periods. Next to the fishes come the amphibians, from the embryology of which we can also draw instructive conclusions. They represent the transition to the higher vertebrates, in which the middle and older stages of ancestral development have been either distorted or curtailed, but in which we find the more recent stages of the phylogenetic process well preserved in ontogeny. We are thus in a position to form a fairly complete idea of the past development of man's ancestors within the vertebrate stem by putting together and comparing the embryological developments of the various groups of vertebrates. And when we go below the lowest vertebrates and compare their embryology with that of their invertebrate relatives, we can follow the genealogical tree of our animal ancestors much farther, down to the very lowest groups of animals.

In entering the obscure paths of this phylogenetic labyrinth, clinging to the Ariadne-thread of the biogenetic law and guided by the light of comparative anatomy, we will first, in accordance with the methods we have adopted, discover and arrange those fragments from the manifold embryonic developments of very different animals from which the stem-history of man can be composed. I would call attention particularly to the fact that

employ this method with the same confidence and right as the geologist. No geologist has ever had ocular proof that the vast rocks that compose our Carboniferous or Jurassic or Cretaceous strata were really deposited in water. Yet no one doubts the fact. Further, no geologist has ever learned by direct observa-

tion these various sedimentary formations deposited in a certain order: yet all are agreed as to this order. This

cannot be rationally understood that they

are. These hypotheses are and indispen-

geologic

the same value, for the same reasons. In formulating them we are acting on the same inductive and deductive methods, and with almost equal confidence, as the geologist. We hold them to be correct, and claim the status of "biological theories" for them, because we cannot understand the nature and origin of man and the other organisms without them,

knowledge of c
ological hypo

ineteenth century are now universally admitted, so our phylogenetic hypotheses, which are still regarded as fantastic in certain quarters, will sooner or later be generally received. It is true that, as all soon appear, our task is not so simple as that of the geologist. It is just as much more difficult and complex as man's organisation is more elaborate than the structure of the rocks.

When we approach this task, we find a auxiliary of the utmost importance in the comparative anatomy and embryology of two lower animal-forms. One of these animals is the lancelet (*Amphioxus*), the other the sea-squirt (*Ascidia*). Both of these animals are very instructive. Both are at the border between the two chief divisions of the animal kingdom—

vertebrates. They

the already mentioned
the Amphiox

(actinias, lampreys, fishes, dipnoans, amphibia, reptiles, birds, and mammals). Following the example of Lamarck, it is usual to put all the other animals together under the head of invertebrates. But, as I have often mentioned already, the group

is composed of a number of very different stems. Of these we have no interest just now

thinoderms, molluscs, and articulates, as they are independent branches of the animal-tree, and have nothing to do with the vertebrates. On the other hand, we are greatly concerned with a very interesting group that has only recently been carefully studied, and that has

the vertebrates.

the stem of the Tunicates. One member of this group, the sea-squirt, very closely approaches the lowest vertebrate, the Amphioxus, in its essential internal structural embryonic development. Until one had any idea of the close con-

cent

ness, it was a very fortunate accident that the embryology of these related forms was discovered just at the time when the question of the descent of the vertebrates from the invertebrates came to the front. In order to understand it properly, we must first consider these remarkable animals in their fully-developed forms and compare their anatomy.

It lives on the flat, sandy parts of the Mediterranean coast, partly buried in the sand, and is apparently found in a number of seas.* It has been found in the North Sea (on the British and Scandinavian coasts and in Heligoland) and at various places on the Mediterranean (for instance, at Nice, Naples, and Messina). It is also found on the coast of Brazil and in the most distant parts of the Pacific Ocean (the coast of Peru, Bonnev, China, Australia, etc.). Recently eight to ten species of the *amphioxus* have been determined, distributed in two or three genera.

Johannes Müller classed the lancelet with the fishes, although he pointed out that the differences between this simple vertebrate and the lowest fishes are much greater than between the fishes and the amphibia. But this was far from expressing the real significance of the

... suddenly lay down the following principle: The *Amphioxus* differs more from the fishes than the fishes do from

* See the simple monograph by Arthur Willey, *Amphioxus and the Ancestry of the Vertebrates*, 1891.

(*cranium*). Of these the only living representatives are the *Amphioxus* and *Pneumphoxus*, though there must have been a number of different species at an early period of the earth's history.

Opposed to the Acrania is the second division of the vertebrates, which comprises all the other members of the stem, from the fishes up to man. All these vertebrates have a head quite distinct from the trunk, with a skull (*cranium*) and brain; all have a centralised heart, fully-formed kidneys, etc. Hence they are called the *Craniota*. These Craniotes are, however, without a skull in their earlier period. As we already know from embryology, even man, like every other mammal, passes in the earlier course of his development through the important stage which we call the *chordate*; at this lower stage the animal has neither vertebrae nor skull nor limbs (Figs 83-86). And even after the formation of the primitive vertebrae has begun, the segmented *fistulus* of the annelotes still has for a long time the simple form of a lyre-shaped disk or a vandel, without limbs or extremities. When we compare this embryonic condition, the vandel-shaped *fistulus*, with the developed lancelet, we may say that the *amphioxus* is, in a certain sense, a permanent sandal-embryo, or a permanent embryonic form of the Acrania; it never rises above a low grade of development which we have long since passed.

The fully-developed lancelet (Fig. 220) is about two inches long, is colourless or of a light red tint, and has the shape of a narrow lancelet-formed leaf. The body is pointed at both ends, but much compressed at the sides. There is no trace of limbs. The outer skin is very thin and delicate, naked, transparent, and composed of two different layers, a simple external stratum of cells, the epidermis, and a thin underlying cutis-layer. Along the middle line of the back runs a narrow fin-fringe which expands behind into an oval tail-fin, and is continued below in a short anus-fin. The fin-fringe is supported by a number of square elastic fin-plates.

In the middle of the body we find a thin string of cartilage, which goes the whole length of the body from front to back, and is pointed at both ends (Fig. 220 e). This straight, cylindrical rod (somewhat compressed for a time) is the axial rod or the *chorda dorsalis*; in the lancelet this is the only trace of a vertebral column. The *chorda* develops no further, but retains

its original simplicity throughout life. It is enclosed by a firm membrane, the *chorda-sheath* or *perichorda*. The real features of this and of its dependent formations are best seen in the transverse section of the *Amphioxus* (Fig. 221). The *perichorda* forms a cylindrical tube immediately over the *chorda*, and the central nervous system, the medullary tube, is enclosed in it. This important psychic organ also remains in its simplest shape throughout life, as a cylindrical tube, terminating with almost equal plainness at either end, and enclosing a narrow canal in its thick wall. However, the fore end is a little rounder, and contains a small, almost imperceptible bulbous swelling of the canal. This must be regarded as the beginning of a rudimentary brain. At the foremost end of it there is a small black pigment-spot, a rudimentary eye; and a narrow canal leads to a superficial sense-organ. In the vicinity of this optic spot we find at the left side a small ciliated depression, the single olfactory organ. There is no organ of hearing. This defective development of the higher sense-organs is probably, in the main, not an original feature, but a result of degeneration.

Underneath the axial rod or *chorda* runs a very simple alimentary canal, a tube that opens on the ventral side of the animal by a mouth in front and anus behind. The oral mouth is surrounded by a ring of cartilage, on which there are twenty to thirty cartilaginous threads (organs of touch, Fig. 220 a). The alimentary canal divides into sections of about equal length by a constriction in the middle. The fore section, or head-gut, serves for respiration; the hind section, or trunk-gut, for digestion. The limit of the two alimentary regions is also the limit of the two parts of the body, the head and the trunk. The head-gut or branchial gut forms a broad gill-crate, the grilled wall of which is pierced by numbers of gill-clefts (Fig. 220 d). The fine bars of the gill-crate between the clefts are strengthened with firm parallel rods, and these are connected in pairs by cross-rods. The water that enters the mouth of the *Amphioxus* passes through these clefts into the large surrounding branchial cavity or *atrium*, and then pours out behind through a hole in it, the respiratory pore (*porus branchialis*, Fig. 220 e). Below, on the ventral side of the gill-crate, there is in the middle

line a ciliated groove with a glandular wall (the hypobranchial groove), which is also found in the Ascidia and the larva of the Cyclostoma. It is interesting because the thyroid gland in the larynx of the higher vertebrates (underneath the "Adam's apple") has been developed from it.

Behind the respiratory part of the gut or liver (hepatic) gut. The small particles that the Amphioxus takes in with the water—Infusoria, diatoms, particles of decomposed plants and animals, etc.—pass from the gill-crate into the digestive part of the canal, and are used up as food.

From a somewhat enlarged portion, that corresponds to the stomach (Fig. 210 r), a long, pouch-like blind sac proceeds straight forward (f). It lies beneath

pulsating in their whole length, and thus driving the colourless blood through the entire body. On the under-side of the gill-crate, in the middle line, there is the trunk of a large vessel that corresponds to the heart of the vertebrates and the trunk of the big artery that proceeds from it; the

number of small vascular arches on each side from this branchial artery, and form little heart-shaped swellings or *bulbula* (m) at their points of departure; they advance along the branchial arches, between the gill-clefts and the fore-gut, and unite, as branchial veins, above the gill-crate in a large trunk blood-vessel that runs under the chorda dorsalis. This is the principal artery or primitive





FIG. 210.

FIG. 210.—Transverse section of an Amphioxus larva, with five gill-clefts, through the middle of the body. FIG. 211.—Diagram of the preceding. (From Haeckel.) A epibranchium, B medullary tube, C chorda, D outer choriodermis, E second epibranchium, F subintestinal vein, G tube, 2 muscle-plate (myotome), 3 skeletal plate (leptotome), 4 notochord, 5 parietal space between dorsal and ventral notochord, 6 slender layer, 7 gut-tube (gut), 8 ventral body-cavity, 9 11 splanchnic vein (ventral body-cavity).



on the left side of the gill-crate, and ends blindly about the middle of it. This is the liver of the Amphioxus, the simplest kind of liver that we meet in any vertebrate. In man also the liver develops, as we shall see, in the shape of a pouch-like blind sac, that forms out of the alimentary canal behind the stomach.

The formation of the circulatory system in this animal is no less interesting. All the other vertebrates have a compressed, thick, pouch-shaped heart, which develops from the wall of the gut at the throat, and from which the blood-vessels proceed; in the Amphioxus there is no special centralised heart, driving the blood by its pulsations. This movement is effected, as in the annelids, by the thin blood-vessels themselves, which discharge the function of the heart, contracting and

expanding in their whole length, and thus driving the colourless blood through the entire body. The branches which it gives off to all parts of the body unite again in a larger venous vessel at the under-side of the gut, called the subintestinal vein (Figs. 210 d, 211 k). This single main vessel of the Amphioxus goes like a closed circular water-conduit along the alimentary canal through the whole body, and pulsates in its whole length above and below. When the upper tube contracts the lower one is filled with blood, and vice versa. In the upper tube the blood flows from front to rear, then back from rear to front in the lower vessel. The whole of the long tube that runs along the ventral side of the alimentary canal and contains venous blood may be called the "principal vein," and may be compared to the ventral vessel in the worms. On the other hand, the long

gut, and are ejected through the mouth.

Above the sexual glands, at the dorsal angle of the atrium, we find the kidneys. These important excretory organs could not be found in the *Amphioxus* for a long

other vertebrates (Fig. 218 *B*). Their internal aperture (Fig. 217 *B*) opens into the body-cavity; their outer aperture into the atrium (*C*). The pronephal canals lie in the middle of the line of the head, outwards




FIG. 216.—Transverse section of the lancelet, in the fore half. (From Swinhö.) The entire covering is the simple cut-layer of the epidermis (*B*). Under this is the thin perium, the subcutaneous tissue of which is indicated. A small connective-tissue partition between the atrium (*M*) and to the chondrodermis. *N* notochord. *L* body-cavity. *A* atrium. *G* upper wall of same. *H* inner wall. *P* pronephal canal. *K* kidney. *O* outer wall of same. *S* sexual gland.

time, on account of their remote position and their smallness; they were discovered in 1890 by Theodor Boveri (Fig. 217 *S*). They are short segmented canals, corresponding to the primitive kidneys of the

from the uppermost section of the gill-arches, and have important relations to the branchial vessels (*H*). For this reason, and in their whole arrangement, the primitive kidneys of the *Amphioxus*

show clearly that they are equivalent to the preoral canals of the Craniotes (Fig. 218 *B*). The preoral duct of the latter (Fig. 218 *C*) corresponds to the branchial cavity or atrium of the former (Fig. 217 *C*).

If we sum up the results of our anatomic study of the *Amphioxus*, an-

both animals in the same division of the animal kingdom. Nevertheless, this classification is indisputably just. Man is only a more advanced stage of the vertebral type that we find unmistakably in the *Amphioxus* in its characteristic features. We need only recall the features of the ideal Primitive Vertebrate




FIG. 217.—Transverse section through the middle of the *Amphioxus*. (From Sherrin). On the left a gill-rod has been struck, and on the right is gill-tissue, consequently on the left we see the whole of a preoral canal (*A*), on the right only the section of its border. *A* genital chamber (ventral section of the gonocoel), *B* preoperculum, *C* the oesophagus, *D* the body cavity, *E* the ventral cavity, *F* the subintestinal vein, *G* the aorta (the left branch connected by a branchial vessel with the subintestinal vein), *H* the renal vessel.

compare them with the familiar organization of man, we shall find an immense distance between the two. As a fact, the highest summit of the vertebrate organization which man represents is in every respect so far above the lowest stage, at which the lancelet remains, that one would at first scarcely believe it possible to class





FIG. 218.—Transverse section through the middle of the primitive *Ichthyops*. (A) the left preoperculum, (B) the right preoperculum, (C) the oesophagus, (D) the body cavity, (E) the ventral cavity, (F) the subintestinal vein, (G) the aorta, (H) the renal vessel.

given in a former chapter, and compare it with the lower stages of human embryonic development, to convince ourselves of our close relationship to the lancelet. (Cf. Chapter XI.)

It is true that the *Amphioxus* is far below all other living vertebrates. It is true that it has no separate head, no developed brain or skull, the characteristic feature of the other vertebrates.

It is (probably as a result of degeneration) without the auscultory organ and the centralised heart that all the others have; and it has no fully-formed kidneys. Every single organ in it is simpler and less advanced than in any of the others. Yet the characteristic connection and arrangement of all the organs is just the same as in the other vertebrates. All these, moreover, pass, during their embryonic development, through a stage in which their whole organisation is no higher than that of the Amphioxus, but is substantially identical with it.



double; *b* atrium *c* cranial *cd* cranial (body) ducts *f* foregut *g* gill-pouches *kb* kidney buds *i* intestine *e* epibranchial groove *a* aortic arches *sr* spiracles *b* branchial gill *c* central canal *d* dorsal canal *e* epibranchial groove *f* foregut *g* gill-pouches

In order to see this quite clearly, it is particularly useful to compare the Amphioxus with the youthful forms of those vertebrates that are classified next to it. This is the class of the Cyclostoma. There are to-day only a few species of

be distributed in two groups. One group comprises the hag-fishes or Myxinoidei. The other group are the Petrolampreys, which are a


familiar delicacy in their marine form. These Cyclostoma are usually classified with the fishes. But they are far below the true fishes, and form a very interesting connecting-group between them and the lancelet. One can see how closely they approach the latter by comparing a young lamprey with the Amphioxus. The chorda is of the same simple character in both, also the medullary tube, that lies above the chorda, and the canal below it. However, in the lamprey the spinal cord lies in front into a simple pear-shaped cerebral vesicle, and at each side of it there are a very simple eye and a rudimentary auditory vesicle. The nose is a single pit, as in the Amphioxus. The two sections of the gut are also just the same and very rudimentary in the lamprey. On the other hand, we see a great advance in the structure of the heart, which is found underneath the gills in the shape of a centralised muscular tube, and is divided into an auricle and a ventricle. Later on the lamprey advances still further, and gets a skull, five cerebral vesicles, a series of independent gill-pouches, etc. This makes all the more interesting the striking resemblance of its immature form to the developed and asexually mature Amphioxus.

While the Amphioxus is thus connected through the Cyclostoma with the fishes, and so with the series of the higher vertebrates, it is, on the other hand, very closely related to a lowly invertebrate marine animal, from which it seems to be entirely remote at first glance. This remarkable animal is the sea-squirt or Ascidia, which was formerly thought to be closely related to the mussel, and so classed in the molluscs. But since the remarkable embryology of these animals was discovered in 1860, there can be no question that they have nothing to do with the molluscs. To the great astonishment of zoologists, they were found, in their whole individual development, to be closely related to the vertebrates. When fully developed the Ascidia are shapeless lumps that would not, at first sight, be taken for animals at all. The oval body, frequently studded with knobs or uneven and lumpy, in which we can discover no special external organs, is

or the floor of the sea. Many species look like potatoes, others like melon-cacti, others like prunes. Many of the Ascidia form transparent crusts or

deposits on stones and marine plants. Some of the larger species are eaten like oysters. Fishermen, who know them very well, think they are not animals, but plants. They are sold in the fish markets of many of the Italian coast-towns with other lower marine animals under the name of "sea-fruit" (*frutti di mare*). There is nothing about them to show that they are animals. When they are taken out of the water with the net the most one can perceive is a slight contraction of the body that causes water to spout out in two places. The bulk of the *Ascidia* are very small, at the most a few inches long. A few species are a foot or more in length. There are many species of them, and they are found in every sea. As in the case of the *Acrania*, we have no fossilised remains of the class, because they have no hard and fossilisable parts. However, they must be of great antiquity, and must go back to the primordial epoch. The name of "Tunicates" is given to the whole class to which the *Ascidia* belong, because the body is enclosed in a thick

occupied by the large grated branchial sac (*br*). This is so like the gill-craie of the *Amphioxus* in its whole arrangement that the resemblance was pointed out by the English naturalist Goodair, years ago, before anything was known of the relationship of the two animals. As a fact, even in the *Ascidia* the mouth (*a*) opens first into this wide branchial



The asc.—Organisation of an *Ascidia* (left view), the dorsal side is turned to the right and the ventral to the left. The mouth (*a*) above, the cloaca (*c*) below, the respiratory pore (*r*) on the side, the siphon (*s*) at the top, the tail (*t*) at the bottom. The branchial sac is formed by a number of diaphragms, over which the food is passed. The rectum opens into the cloaca (*c*) from which the excrement is expelled with the respiratory water. The mantle-hole or siphon (*s*), in the *Gegartia*.

sac. The respiratory water passes through the lattice-work of the branchial sac into the branchial cavity, and is ejected from this by the respiratory pore (*r*). Along the ventral side of the branchial sac runs a ciliated groove—the hypobranchial groove which we have previously found at the same spot in the *Amphioxus*. The food of the *Ascidia* also

has a number of remarkable of

vegetal substance
slurps of the plant
the wood. The

that have a real cellulose or woody coat. Sometimes the cellulose mantle is brightly coloured, at other times colourless. Not infrequently it is set with needles or hairs,

The hind end, which corresponds to the tail of the *Amphioxus*, is usually attached, often by means of regular roots. The dorsal and ventral sides differ a good deal internally, but frequently cannot be distinguished externally. If we open the thick tunic or mantle in order to examine the internal organisation, we first find a spacious cavity filled with water—the mantle-cavity or respiratory cavity (Fig. 220 *cf*). It is also called the branchial cavity and the cloaca, because it receives the excrements and sexual products as well as the respiratory water. The greater part of the respiratory cavity is

consists of tiny organisms, infusoria, diatoms, parts of decomposed marine plants and animals, etc. These pass with the water into the gill-crate and the digestive part of the gut at the end of it, at first into an enlargement of it that represents the stomach. The adjoining small intestine usually forms a loop, bends forward, and opens by an anus (Fig. 220 *e*), not directly outwards, but first into the mantle cavity; from this the excrements are ejected by a common outlet (*d'*) together with the used-up



= —Organization of an Ascidia (as in Fig.

also usually confounded with the whole atrium, or peribranchial cavity, full of water.

There is Ascidia of

axial skeleton. It is the more interesting that the young animal that emerges from the ovum has a chorda, and that there is a rudimentary medullary tube above it. The latter is wholly atrophied in the developed Ascidia, and looks like a small nerve-ganglion in front above the gill-crate. It corresponds to the upper "gullet-ganglion" or "primitive brain" in other vertebrates. Special sense-organs are either wanting altogether or are only found in a very rudimentary form, as simple optic spots and touch-capsules or tentacles that surround the mouth. The muscular system is very slightly and irregularly developed. Immediately under the thin cuticle, and closely connected with it, we find a thin muscle

c, as in the worms. On the other hand, the Ascidia has a centralized heart. In this respect it seems to be more advanced than the Amphioxus. On the ventral side of the gut, some distance behind the gill-crate, there is a spindle-shaped heart. It retains permanently the simple tubular form that we find temporarily as the first structure of the heart in the vertebrates. This simple heart of the Ascidia has, however, a remarkable peculiarity. It contracts in alternate directions. In all other animals the beat of the heart is always in the same direction (generally from rear to front); it changes in the Ascidia to the reverse direction. The heart contracts first from the rear to the front, stands still for a minute, and then begins to beat the

water through the openings of the mantle. (From *Meier-Schwarz*.)

water and the sexual products. The outlet is sometimes called the branchial pore, and sometimes the cloaca or ejection-aperture. In many of the Ascidia a glandular mass opens into the gut, and this represents the liver. In some there is another gland besides the liver, and this is taken to represent the kidneys. The body-cavity proper, or coelom, which is filled with blood and encloses the hepatic gut, is very narrow in the Ascidia, as in the Amphioxus, and is here

alternately as arteries and veins. This feature is found in the Tunicates alone.

Of the other chief organs we have still to mention the sexual glands, which lie right behind in the body-cavity. All the Ascidia are hermaphrodites. Each individual has a male and a female gland, and so is able to fertilize itself. The ripe ova (Fig. 221 *o*) fall directly from the ovary (*o*) into the mantle-cavity. The male sperm is conducted

the testicle (*t*) by a special duct (*nd*). Fertilization is accomplished here, and in many of the Ascidia developed embryos are found. These are then

ejected with the breathing-water through the cloaca (*q*), and so "born alive."

If we now glance at the entire structure of the simple Ascidia (especially *Phallusia*, *Cynthia*, etc.) and compare it with that of the Amphioxus, we shall find that the two have few points of contact. It is true that the fully-developed Ascidia resembles the Amphioxus in several important features of its internal structure, and especially in the peculiar character of the gill-grate and gut. But

in most other features of organization it is so far removed from it, and is so unlike it in external appearance, that the really close relationship of the two was not discovered until their embryology was studied. We will now compare the embryonic development of the two animals, and find to our great astonishment that the same embryonic form develops from the ovum of the Amphioxus as from that of the Ascidia—a typical *chaetula*.

CHAPTER XVII.

EMBRYOLOGY OF THE LANCELET AND THE SEA-SQUIRT

so prominent that there was the greatest difficulty in the earlier stages of classification in determining the affinity of these two great groups. When scientists began to speak of the affinity of the various animal groups in more than a figurative—in a genealogical—sense, this question came at once to the front, and seemed to constitute one of the chief obstacles to the carrying-out of the evolutionary theory. Even earlier, when they had studied the relations of the chief groups, without any idea of real genealogical connection, they believed they had found here and there among the invertebrates points of contact with the vertebrates.

One of the worms, especially, seemed to approach the vertebrates in structure, such as the marine arrow-worm (*Sagitta*). But on closer study the analogies proved untenable. When Darwin gave an impulse to the construction of a real stem-history of the animal kingdom by his reform of the theory of evolution, the solution of this problem was found to be particularly difficult. When I made the first attempt in my *General Morphology* (1866) to work out the theory and apply it to classification, I found no problem of phylogeny that gave me so much trouble

But just at this time the true link was discovered, and at a point where it was least expected. Towards the end of 1866 two works of the Russian zoologist, Kowalevsky, who had lived for some time at Naples, and studied the embryology of the lower animals, were issued in the publications of the St. Petersburg Academy. A fortunate accident had directed the attention of this able observer almost simultaneously to the embryology of the lowest vertebrate, the Amphioxus, and that of an invertebrate, the close affinity of which to the Amphioxus had been least suspected, the Ascidia. To the extreme astonishment of all zoologists who were interested in this important question, there turned out to be the utmost resemblance in structure from the commencement of development between these two very different animals—the lowest vertebrate and the misshaped, sessile invertebrate. With this undeniable identity of ontogenesis, which can be demonstrated to an astounding extent, we had, in virtue of the biogenetic law, discovered the long-sought genealogical link, and definitely identified the invertebrate group that represents the nearest blood-relatives of the vertebrates.

The discovery was confirmed by other zoologists, and there can no longer be any doubt that of all the classes of invertebrates that of the Tunicates is most closely related to the vertebrates, and of the Tunicates the nearest are the Ascidians. We cannot say that the vertebrates are descended from the Ascidians—and still less the reverse—but we can say that of all the invertebrates it is the Tunicates, and, within this group, the Ascidians, that are the nearest blood-relatives of the ancient stem-form of the vertebrates. We must assume as the common ancestral group of both stems an extinct family of the extensive *vermula*-stem, the *Prochordones* or *Prochordata* ("primitive chorda-animals").

In order to appreciate fully this remarkable fact, and especially to secure the sound basis we seek for the genealogical tree of the vertebrates, it is necessary to study thoroughly the embryology of both these animals, and compare the individual development of the *Amphioxus* step by step with that of the *Ascidia*. We begin with the ontogeny of the *Amphioxus*.

From the concordant observations of Kovalevsky at Naples and Hatschek at Messina, it follows, firstly, that the ovum-segmentation and gastrulation of the *Amphioxus* are of the simplest character. They take place in the same way as we find them in many of the lower animals of different invertebrate stems, which we have already described as original or primordial; the development of the *Ascidia* is of the same type. Sexually-mature specimens of the *Amphioxus*, which are found in great quantities at Messina from April or May onwards, begin as a rule to eject their sexual products in the evening; if you catch them about the middle of a warm night and put them in a glass vessel with seawater, they immediately eject through the mouth their accumulated sexual products, in consequence of the disturbance. The males give out masses of sperm, and the females discharge ova in such quantity that many of them stick to the filaments about their mouths. Both kinds of cells pass first into the mantle-cavity after the opening of the gonads, proceed through the gill-clefts into the branchial gut, and are discharged from this through the mouth.

The ova are simply round cells. They are only $\frac{1}{16}$ of an inch in diameter, and thus are only half the size of the mammal

ova, and have no distinctive features. The clear protoplasm of the mature ovum is made so turbid by the numbers of dark granules of food-yolk or deutoplasm scattered in it that it is difficult to follow the process of fecundation and the behaviour of the two nuclei during it (p. 51). The active elements of the male sperm, the cone-shaped spermatozoa, are similar to those of most other animals (cf. Fig. 20). Fecundation takes place when these lively ciliated cells of the sperm approach the ovum, and seek to penetrate into the yolk-matter or the cellular substance of the ovum with their head-part—the thicker part of the cell that encloses the nucleus. Only one spermatozoon can bore its way into the yolk at one pole of the ovum-ovus, its head or nucleus coalesces with the female nucleus, which remains after the extrusion of the directive bodies from the germinal vesicle. Thus is formed the "stem-nucleus," or the nucleus of the "stem-cell" (cytula, Fig. 2). This now undergoes total segmentation, dividing into two, four, eight, sixteen, thirty-two cells, and so on. In this way we get the spherical, mulberry-shaped body, which we call the *morula*.

The segmentation of the *Amphioxus* is not entirely regular, as was supposed after the first observations of Kovalevsky (1866). It is not completely equal, but a little unequal. As Hatschek afterwards found (1879), the segmentation-cells only remain equal up to the *morula*-stage, the spherical body of which consists of thirty-two cells. Then, as always happens in unequal segmentation, the more sluggish vegetal cells are outstripped in the cleavage. At the lower or vegetal pole of the ovum a crowd of eight large endodermic cells remains for a long time unchanged, while the other cells divide, owing to the formation of a series of horizontal circles, into an increasing number of crowns of sixteen cells each. Afterwards the segmentation-cells get more or less irregularly displaced, while the segmentation-cavity enlarges in the centre of the *morula*; in the end the former all lie on the surface of the latter, so that the fetus attains the familiar blastula shape and form: the wall of which consists of a single stratum of cells (Fig. 38 A-C). This layer is the blastoderm, the simple epithelium from the cells of which all the tissues of the body proceed.

These important early embryonic processes take place so quickly in the *Amphioxus* that four or five hours after fecundation, or about midnight, the spherical blastula is completed. A pit-like depression is then formed at the vegetal pole of it, and in consequence of this the hollow sphere doubles on itself (Fig. 38 D). This pit becomes deeper and deeper (Fig. 38 E, F); at last the invagination (or doubling) is complete, and the inner or folded part of the blastula-wall lies on the inside of the outer wall. We thus get a hollow hemisphere, the thin wall of which is made up of two layers of cells (Fig. 38 G). From hemispherical the body soon becomes almost spherical once more, and then oval, the internal cavity enlarging considerably and its mouth growing narrower (Fig. 213). The form which the *Amphioxus*-embryo has thus reached is a real "cup-larva" or *gastrula*, of the original simple type that we have previously described as the "bell-gastrula" or *archegastrula* (Figs. 20 35).

As in all the other animals that form an *archegastrula*, the whole body is nothing but a simple gastric sac or stomach, its internal cavity is the primitive gut (*pro-gut* or *archenteron*, Fig. 38 G, 35 d), and its aperture the primitive mouth (*protoma* or *blastopore*, e). The wall is at once gut-wall and body-wall. It is composed of two simple cell-layers, the familiar primary germinal layers. The inner layer or the invaginated part of the blastoderm, which immediately encloses the gut-cavity is the endoderm, the inner or vegetal germ-layer, from which develop the wall of the alimentary canal and all its appendages, the coelom-pouches, etc. (Figs. 35, 36 f). The outer stratum of cells, or the non-invaginated part of the blastoderm, is the ectoderm, the outer or animal germ-layer, which provides the outer skin (epidermis) and the nervous system (e). The cells of the endoderm are much larger, darker, and more fatty than those of the ectoderm, which are clearer and less rich in fatty particles. Hence before and during invagination there is an increasing differentiation of the inner from the outer layer. The animal cells of the outer layer soon develop vibratory hairs; the vegetal cells of the inner layer do so much later. A thread-like process grows out of each cell, and effects continuous vibratory movements. By the vibrations of these slender hairs the gastrula of the *Amphi-*

oxus swims about in the sea, when it has pierced the thin ovolemma, like the gastrula of many other animals (Fig. 36). As in many other lower animals, the cells have only one whip-like hair each, and so are called *flagellate* (whip) cells (in contrast with the *ciliated* cells, which have a number of short lashes or cilia).

In the further course of its rapid development the roundish bell-gastrula becomes elongated, and begins to flatten on one side, parallel to the long axis. The flattened side is the subsequent dorsal side; the opposite or ventral side remains curved. The latter grows more quickly than the former, with the result that the primitive mouth is forced to the dorsal side (Fig. 39). In the middle of the dorsal surface a shallow longitudinal groove or furrow is formed (Fig. 79), and the edges of the body rise up on each side of this groove in the shape of two parallel swellings. This groove is, of course, the dorsal furrow, and the swellings are the dorsal or medullary swellings; they form the first structure of the central nervous system, the medullary tube. The medullary swellings now rise higher, the groove between them becomes deeper and deeper. The edges of the parallel swellings curve towards each other, and at last unite, and the medullary tube is formed (Figs. 83 m, 83 n). Hence the formation of a medullary tube out of the outer skin takes place in the naked dorsal surface of the free-swimming larva of the *Amphioxus* in just the same way as we have found in the embryo of man and the higher animals within the foetal membranes.

Simultaneously with the construction of the medullary tube we have in the *Amphioxus*-embryo the formation of the chorda, the coelom-pouches, and the mesoderm proceeding from their wall. These processes also take place with characteristic simplicity and clearness, so that they are very instructive to compare with the *vermalia* on the one hand and with the higher vertebrates on the other. While the medullary groove is sinking in the middle line of the flat dorsal side of the oval embryo, and its parallel edges unite to form the ectodermic neural tube, the single chorda is formed directly underneath them, and on each side of this a parallel longitudinal fold, from the dorsal wall of the primitive gut. These longitudinal folds of the endoderm proceed from the primitive mouth, or from its lower

and hinder edge. Here we see at an early stage a couple of large entodermic cells, which are distinguished from all the others by their great size, round form, and fine-grained protoplasm; they are the two promesoblasts, or polar cells of the mesoderm (Fig. 83 p). They indicate the ~~medial~~ starting-point of the two colorers, which grow from this spot and outer germinal layers, sever themselves from the primitive gut, and provide the cellular material for the middle layer.

Immediately after their formation two colom-pouches of the Amphioxus divide into several parts by longitudinal and transverse folds. Each of the pouches is divided into an upper dorsal and a lower ventral section by a couple of lateral longitudinal folds (Fig. 82). But these are again divided by several parallel transverse folds into a number of successive sacs, the primitive segments or somites (formerly called by the unsuitable name of "primitive vertebrae"). They have a different future above and below. The upper or dorsal segments, the *episomites*, lose their cavity later on, and form with their cells the muscular plates of the trunk. The lower or ventral segments, the *hypsomites*, corresponding to the lateral plates of the cranio-embryo, fuse together in the upper part owing to the disappearance of their lateral walls, and thus form the later body-cavity (metacoel); in the lower part they remain separate, and afterwards form the segmental gonads.

In the middle, between the two lateral in-folds of the primitive gut, a single central organ develops at an early stage in the middle line of its dorsal wall. This is the dorsal chorda (Fig. 83, 84 ch). This axial rod, which is the foundation of the later vertebral column in all the vertebrates, and is the only representative of it in the *Amphioxus*, originates from the entoderm.

In consequence of these important folding-processes in the primitive gut, the simple entodermic tube divides into four different sections:—I., underneath, ventral side, the permanent alimentary canal or permanent gut; II., above, at the dorsal side, the axial rod or chorda; and III., the two colom-sacs, which immediately sub-divide into two structures—IIIa., above, on the dorsal side, the *episomites*, the double row of primitive or muscular segments; and IIIb., below, on each side of the gut, the *hypsomites*,

the two lateral plates that give rise to the sex-glands, and the cavities of which partly unite to form the body-cavity. At the same time, the neural or medullary tube is formed above the chorda, on the dorsal surface, by the closing of the parallel medullary swellings. All these processes, which outline the typical structure of the vertebrate, take place with astonishing rapidity in the embryo of the *Amphioxus*; in the afternoon of the first day, or twenty-four hours after fertilisation, the young vertebrate, the typical ~~vertebrate~~ ^{Amphioxus} is to eight times.

The chief occurrence on the second day of development is the construction of the

the alimentary tube is found to be entirely closed, after the closing of the primitive mouth; it only communicates behind by the neurenteric canal with the medullary tube. The permanent mouth is a secondary formation, at the opposite end. Here, at the end of the second day, we find a pit-like depression in the outer skin, which penetrates inwards into the closed gut. The anus is formed behind in the same way a few hours later (in the vicinity of the additional gastrula-mouth). In man and the higher verte-

brates, as we have seen, as flat pits in the outer skin, they then penetrate inwards, gradually becoming connected with the blind ends of the closed gut-tube. During the second day the *Amphioxus*-embryo undergoes few other changes. The number of primitive segments increases, and generally amounts to fourteen, some forty-eight to fifty hours after impregnation.

Almost simultaneously with the formation of the mouth the first gill-cleft breaks through in the fore section of the *Amphioxus*-embryo (generally forty hours after the commencement of development). It is ~~at first~~ ^{at first} independent, as the food material stored is completely used up. The further development of the free larva takes place very slowly, and

The body becomes much longer, and is compressed at the sides, the head-end being broadened in a sort of triangle. Two rudimentary sense-organs are developed in it. Inside we find the first blood-vessels, an upper or dorsal vessel, corresponding to the aorta, between the gut and the dorsal cord, and a lower or ventral

Now, the gills or respiratory organs also are formed at the fore-end of the alimentary canal. The whole of the anterior or respiratory section of the gut is converted into a gill-plate, which is pierced trellis-like by numbers of branchial-holes, as in the scudica. This is done by the foremost part of the gut-wall joining star-wise with the outer skin, and the formation of clefts

we have previous-
lative Vertebrate" (Figs 98-102). But the body afterwards undergoes various modifications, especially in the fore-part. These modifications do not concern us, as they depend on special adaptations, and do not affect the hereditary vertebrate type. When the free-swimming Amphib-




FIG. 98.




FIG. 99.

at the point of connection, piercing the wall and leading into the gut from without. At first there are very few of these branchial clefts; but there are soon a number of them—first in one, then in two.

The foremost gill-cleft is the oldest

In the early stages of embryonic development of fine gill-clefts, supported on a number of stiff branchial rods; these are connected in pairs by transverse rods.

At an early stage of embryonic develop-




FIG. 100. Sections of young lancelets, from the side (C).

FIG. 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631, 632, 633, 634, 635, 636, 637, 638, 639, 640, 641, 642, 643, 644, 645, 646, 647, 648, 649, 650, 651, 652, 653, 654, 655, 656, 657, 658, 659, 660, 661, 662, 663, 664, 665, 666, 667, 668, 669, 670, 671, 672, 673, 674, 675, 676, 677, 678, 679, 680, 681, 682, 683, 684, 685, 686, 687, 688, 689, 690, 691, 692, 693, 694, 695, 696, 697, 698, 699, 700, 701, 702, 703, 704, 705, 706, 707, 708, 709, 710, 711, 712, 713, 714, 715, 716, 717, 718, 719, 720, 721, 722, 723, 724, 725, 726, 727, 728, 729, 730, 731, 732, 733, 734, 735, 736, 737, 738, 739, 740, 741, 742, 743, 744, 745, 746, 747, 748, 749, 750, 751, 752, 753, 754, 755, 756, 757, 758, 759, 760, 761, 762, 763, 764, 765, 766, 767, 768, 769, 770, 771, 772, 773, 774, 775, 776, 777, 778, 779, 780, 781, 782, 783, 784, 785, 786, 787, 788, 789, 790, 791, 792, 793, 794, 795, 796, 797, 798, 799, 800, 801, 802, 803, 804, 805, 806, 807, 808, 809, 810, 811, 812, 813, 814, 815, 816, 817, 818, 819, 820, 821, 822, 823, 824, 825, 826, 827, 828, 829, 830, 831, 832, 833, 834, 835, 836, 837, 838, 839, 840, 841, 842, 843, 844, 845, 846, 847, 848, 849, 850, 851, 852, 853, 854, 855, 856, 857, 858, 859, 860, 861, 862, 863, 864, 865, 866, 867, 868, 869, 870, 871, 872, 873, 874, 875, 876, 877, 878, 879, 880, 881, 882, 883, 884, 885, 886, 887, 888, 889, 890, 891, 892, 893, 894, 895, 896, 897, 898, 899, 900, 901, 902, 903, 904, 905, 906, 907, 908, 909, 910, 911, 912, 913, 914, 915, 916, 917, 918, 919, 920, 921, 922, 923, 924, 925, 926, 927, 928, 929, 930, 931, 932, 933, 934, 935, 936, 937, 938, 939, 940, 941, 942, 943, 944, 945, 946, 947, 948, 949, 950, 951, 952, 953, 954, 955, 956, 957, 958, 959, 960, 961, 962, 963, 964, 965, 966, 967, 968, 969, 970, 971, 972, 973, 974, 975, 976, 977, 978, 979, 980, 981, 982, 983, 984, 985, 986, 987, 988, 989, 990, 991, 992, 993, 994, 995, 996, 997, 998, 999, 1000.

FIG. 100. Sections of young lancelets, from the side (C). FIG. 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631, 632, 633, 634, 635, 636, 637, 638, 639, 640, 641, 642, 643, 644, 645, 646, 647, 648, 649, 650, 651, 652, 653, 654, 655, 656, 657, 658, 659, 660, 661, 662, 663, 664, 665, 666, 667, 668, 669, 670, 671, 672, 673, 674, 675, 676, 677, 678, 679, 680, 681, 682, 683, 684, 685, 686, 687, 688, 689, 690, 691, 692, 693, 694, 695, 696, 697, 698, 699, 700, 701, 702, 703, 704, 705, 706, 707, 708, 709, 710, 711, 712, 713, 714, 715, 716, 717, 718, 719, 720, 721, 722, 723, 724, 725, 726, 727, 728, 729, 730, 731, 732, 733, 734, 735, 736, 737, 738, 739, 740, 741, 742, 743, 744, 745, 746, 747, 748, 749, 750, 751, 752, 753, 754, 755, 756, 757, 758, 759, 760, 761, 762, 763, 764, 765, 766, 767, 768, 769, 770, 771, 772, 773, 774, 775, 776, 777, 778, 779, 780, 781, 782, 783, 784, 785, 786, 787, 788, 789, 790, 791, 792, 793, 794, 795, 796, 797, 798, 799, 800, 801, 802, 803, 804, 805, 806, 807, 808, 809, 810, 811, 812, 813, 814, 815, 816, 817, 818, 819, 820, 821, 822, 823, 824, 825, 826, 827, 828, 829, 830, 831, 832, 833, 834, 835, 836, 837, 838, 839, 840, 841, 842, 843, 844, 845, 846, 847, 848, 849, 850, 851, 852, 853, 854, 855, 856, 857, 858, 859, 860, 861, 862, 863, 864, 865, 866, 867, 868, 869, 870, 871, 872, 873, 874, 875, 876, 877, 878, 879, 880, 881, 882, 883, 884, 885, 886, 887, 888, 889, 890, 891, 892, 893, 894, 895, 896, 897, 898, 899, 900, 901, 902, 903, 904, 905, 906, 907, 908, 909, 910, 911, 912, 913, 914, 915, 916, 917, 918, 919, 920, 921, 922, 923, 924, 925, 926, 927, 928, 929, 930, 931, 932, 933, 934, 935, 936, 937, 938, 939, 940, 941, 942, 943, 944, 945, 946, 947, 948, 949, 950, 951, 952, 953, 954, 955, 956, 957, 958, 959, 960, 961, 962, 963, 964, 965, 966, 967, 968, 969, 970, 971, 972, 973, 974, 975, 976, 977, 978, 979, 980, 981, 982, 983, 984, 985, 986, 987, 988, 989, 990, 991, 992, 993, 994, 995, 996, 997, 998, 999, 1000.

ous-larva is three months old, it abandons its pelagic habits and changes into the young animal that lives in the sand. In spite of its smallness (one-eighth of an inch), it has substantially the

same adult. As regards the remaining organs of the Amphioxus, we need only mention that the gonads or sexual glands are developed very late, immediately out of the inner cell-layer of the

body-cavity. Although we can find afterwards no continuation of the body-cavity (Fig. 216 *E'*) in the lateral walls of the mantle-cavity, in the gill-covers or mantle-folds (Fig. 224 *E'*), there is one present in the beginning (Fig. 223 *LA*). The sexual cells are formed below, at the bottom of this continuation (Fig. 224 *S*). For the rest, the subsequent development into the adult Amphioxus of the larva we have followed is so simple that we need not go further into it here.

We may now turn to the embryology of the Ascidia, an animal that seems to stand so much lower and to be so much more simply organised, remaining for the greater part of its life attached to the bottom of the sea like a shapeless lump. It was a fortunate accident that Kowalevsky first examined just those larger specimens of the Ascidiae that show most clearly the relationship of the vertebrates to the invertebrates, and the larvae of which behave exactly like those of the Amphioxus in the first stages of development. This resemblance is so close in the main features that we have only to repeat what we have already said of the ontogenesis of the Amphioxus.

The ovum of the larger Ascidia (*Phallusia*, *Cynthia*, etc.) is a simple round cell of $\frac{1}{16}$ to $\frac{1}{8}$ of an inch in diameter. In the thick fine-grained yolk we find a clear round germinal vesicle of about $\frac{1}{16}$ of an inch in diameter, and this encloses a small embryonic spot or nucleolus. Inside the membrane that surrounds the ovum, the stem-cell of the Ascidia, after fecundation, passes through just the same metamorphoses as the stem-cell of the Amphioxus. It undergoes total segmentation; it divides into two, four, eight, sixteen, thirty-two cells, and so on. By continued total cleavage the morula, or mulberry-shaped cluster of cells, is formed. Fluid gathers inside it, and thus we get once more a globular vesicle (the blastula); the wall of this is a single stratum of cells, the blastoderm. A real gastrula (a simple bell-gastrula) is formed from the blastula by invagination, in the same way as in the amphioxus.

Up to this there is no definite ground in the embryology of the Ascidiae for bringing them into close relationship with the Vertebrates; the same gastrula is formed in the same way in many other animals of different stems. But we now find an embryonic process that is peculiar to the Vertebrates, and that proves irrefragably the affinity of the Ascidiae to the

Vertebrates. From the epidermis of the gastrula a medullary tube is formed on the dorsal side, and, between this and the primitive gut, a chorda, these are the organs that are otherwise only found in Vertebrates. The formation of these very important organs takes place in the Ascidia-gastrula in precisely the same way as in that of the Amphioxus. In the Ascidia (as in the other case) the oval gastrula is first flattened on one side—the subsequent dorsal side. A groove or furrow (the medullary groove) is sunk in the middle line of the flat surface, and two parallel longitudinal swellings arise on either side from the skin layer. These medullary swellings join together over the furrow, and form a tube, in this case, again, the neural or medullary tube is at first open in front, and connected with the primitive gut behind by the neurontic canal. Further, in the Ascidia-larva also the two permanent apertures of the alimentary canal only appear later, as independent and new formations. The permanent mouth does not develop from the primitive mouth of the gastrula; this primitive mouth closes up, and the later anus is formed near it by invagination from without, on the hinder end of the body, opposite to the aperture of the medullary tube.

During these important processes, that take place in just the same way in the Amphioxus, a tail-like projection grows out of the posterior end of the larva-body, and the larva folds itself up within the round exocoelom in such a way that the dorsal side is curved and the tail is forced on to the ventral side. In this tail is developed—starting from the primitive gut—a cylindrical string of cells, the fore end of which pushes into the body of the larva, between the alimentary canal and the neural canal, and is no other than the chorda dorsalis. This important organ had hitherto been found only in the Vertebrates, not a single trace of it being discoverable in the Invertebrates. At first the chorda only consists of a single row of large endodermic cells. It is afterwards composed of several rows of cells. In the Ascidia-larva, also, the chorda develops from the dorsal middle part of the primitive gut, while the two coelom-pouches detach themselves from it on both sides. The simple body-cavity is formed by the coalescence of the two.

When the Ascidia-larva has attained

this stage of development it begins to move about in the ovotermus. This causes the membrane to burst. The larva emerges from it, and swims about on its ear-like tail.

These five have been known for a long time. They were first observed by Darwin during his voyage round the world in 1833. They resemble tadpoles in outward appearance, and use their tails as oars, as the tadpoles do. However, this lively and highly-developed condition does not last long. At first there is a progressive development, the foremost part of the medullary tube enlarges into a brain, and inside this two single sense-organs are developed, a dorsal auditory vesicle and a ventral eye. Then a heart is formed on the ventral side of the animal, or the lower wall of the gut, in the same simple form and at the same spot at which the heart is developed in man, and

solid, spindle-shaped

backward, as is the case with the adult Ascidia. In this way the sanguineous fluid accumulated in the hollow muscular tube is driven in alternate directions into the blood-vessels, which develop at both ends of the medullary tube. One principal vessel runs along the dorsal side of the gut, another along its ventral side. The former corresponds to the aorta and the dorsal vessel in the worms. The other corresponds to the subintestinal vein and the ventral vessel of the worms.

With the formation of these organs the progressive development of the Ascidia comes to an end, and degeneration sets in. The free-swimming larva sinks to the floor of the sea, abandons its locomotive habits, and attaches itself to stones, marine plants, mussel-shells, corals, and other objects; this is done with the part of the body that was foremost in movement. The attachment is effected by a number of out-growths, usually three, which can be seen even in the free-swimming larva. The tail is lost, as there is no further use for it. It undergoes a fatty degeneration, and disappears with the chorda dorsalis. The tailless body changes into an unshapely tube, and, by the atrophy of some parts and the modification of others, gradually

assumes the appearance we have already described.

Among the living Tunicates there is a very interesting group of small animals that remain throughout life at the stage of development of the tailed, free Ascidia-larva, and swim about briskly in the sea by means of their broad ear-tail. These are the remarkable *Copepata* (*Appendi-*




FIG. 225.—An Appendicularia (*Copepata*), seen from the left. *a*, mouth; *b*, branchial gut; *c*, gullet; *d*, stomach; *e*, brain; *f*, brain (ganglion above the gullet); *g*, auditory vesicles; *h*, dorsal groove under the gills; *i*, heart; *j*, notochord; *k*, urinary; *l*, chorda; *m*, tail.

caris and *Tenillerns*, Fig. 225). They are the only living Vertebrates that have throughout life a chorda dorsalis and a neural string above it, the latter must be regarded as the prolongation of the cerebral ganglion and the equivalent of the medullary tube. Their branchial gut also opens directly outwards by a pair of

branchial clefts. These instructive Cope-lata, comparable to permanent Ascidia-larvæ, come next to the extinct Prochordonia, those ancient worms which we must regard as the common ancestors of the Tunicates and Vertebrates. The chorda of the Appendicularia is a long, cylindrical string (Fig. 225 c), and serves as an attachment for the muscles that work the flat out-tail.

Among the various modifications which the Ascidia-larva undergoes after its establishment at the sea-floor, the most interesting (after the loss of the axial rod) is the atrophy of one of its chief organs, the medullary tube. In the Amphioxus the spinal marrow continues to develop, but in the Ascidia the tube soon shrinks into a small and insignificant nervous ganglion that lies above the mouth and the gill-crate, and is in accord with the extremely slight mental power of the animal. This insignificant relic of the medullary tube seems to be quite beyond comparison with the nervous centre of the vertebrate, yet it started from the same structure as the spinal cord of the Amphioxus. The sense-organs that had been developed in the fore part of the neural tube are also lost; no trace of them can be found in the adult Ascidia. On the other hand, the alimentary canal becomes a most extensive organ. It divides presently into two sections—a wide fore or branchial gut that serves for respiration, and a narrower hind or hepatic gut that accomplishes digestion. The branchial or head-gut of the Ascidia is small at first, and opens directly outwards only by a couple of lateral ducts or gill-clefts—a permanent arrangement in the Cope-lata. The gill-clefts are developed in the same way as in the Amphioxus.

Under greatly increased
a large gill-crate, pierced like lat
work. In the middle line of its ven-
side we find the hypobranchial groove.
The mantle or cloaca-cavity (the atrium)
that surrounds the gill-crate is also

formed in the same way in the Ascidia as in the Amphioxus. The ejection-opening of this peribranchial cavity corresponds to the branchial pore of the Amphioxus. In the adult Ascidia the branchial gut and the heart on its ventral side are almost the only organs that recall the original affinity with the vertebrate.

The further development of the Ascidia in detail has no particular interest for us, and we will not go into it. The chief result that we obtain from its embryology is the complete agreement with that of the Amphioxus in the earliest and most important embryonic stages. They do not begin to diverge until after the medullary tube and alimentary canal, and the axial rod with the muscles between the two, have been formed. The Amphioxus continues to advance, and resembles the embryonic forms of the higher vertebrates; the Ascidia degenerates more and more, and at last, in its adult condition, has the appearance of a very imperfect invertebrate.

If we now look back on all the remarkable features we have encountered in the structure and the embryonic development of the Amphioxus and the Ascidia, and compare them with the features of man's embryonic development which we have previously studied, it will be clear that I have not exaggerated the importance of these very interesting animals. It is evident that the Amphioxus from the vertebrate side and the Ascidia from the invertebrate form the bridge by which we can span the deep gulf that separates the two great divisions of the animal kingdom. The radical agreement of the lancelet and the sea-squirt in the first and most important stages of development shows something more than their affinity and their proximity in classification, it shows a common blood-relationship and their origin and the same. In this throws considerable light on the oldest roots of man's genealogical tree.

CHAPTER XVIII.

DURATION OF THE HISTORY OF OUR STEM

Our comparative investigation of the anatomy and ontogeny of the *Amphioxus* and *Ascidia* has given us invaluable assistance. We have, in the first place, bridged the wide gulf that has existed up to the present between the Vertebrates and Invertebrates; and, in the second place, we have discovered in the embryology of the *Amphioxus* a number of ancient evolutionary stages that have long since disappeared from human embryology, and have been lost, in virtue of the law of curtailed heredity. The chief of these stages are the spherical *blastula* (in its simplest primary form) and the succeeding *archigastrea*, the pure, original form of the *gastrula* which the *Amphioxus* has preserved to this day, and which we find in the same form in a number of Invertebrates of various classes. Not less important are the later embryonic forms of the *colomula*, the *chordula*, etc.

Thus the embryology of the *Amphioxus* and the *Ascidia* has so much increased our knowledge of man's stem-history that, although our empirical information is still very incomplete, there is now no defect of any great consequence in it. We may now, therefore, approach our proper task, and reconstruct the phylogeny of man in its chief lines with the aid of this evidence of comparative anatomy and ontogeny. In this the reader will soon see the immense importance of the direct application of the biogenetic law. But before we enter upon the work it will be useful to make a few general observations that are necessary to understand the processes afloat.

We must say a few words with regard to the period in which the human race was evolved from the animal kingdom. The first thought that occurs to one in

between the duration of man's ontogeny and phylogeny. The individual needs only nine months for his complete development, from the fecundation of the ovum to the moment when he leaves the maternal womb. The human embryo runs its whole course in the brief space of

forty weeks (as a rule, 280 days). In many other mammals the time of the embryonic development is much the same as in man—for instance, in the cow. In the horse and ass it takes a little longer, forty-three to forty-five weeks, in the camel, thirteen months. In the largest mammals, the embryo needs a much longer period for its development in the womb—a year and a half in the rhinoceros, and ninety weeks in the elephant. In these cases pregnancy lasts twice as long as in the case of man, or one and three-quarter years. In the smaller mammals the embryonic period is much shorter. The smallest mammals, the dwarf-mice, develop in three weeks; hares in four weeks, rats and marmots in five weeks, the dog in nine, the pig in seventeen, the sheep in twenty-one and the goat in thirty-six. Birds develop still more quickly. The chick only needs, in normal circumstances, three weeks for its full development. The duck needs twenty-five days, the turkey twenty-seven, the peacock thirty-one, the swan forty-two, and the cassowary sixty-five. The smallest bird, the humming-bird, leaves the egg after twelve days. Hence the duration of individual development within the fetal membranes is, in the mammals and birds, clearly related to the absolute size of the body of the animal in question. But this is not the only determining feature. There are a number of other circumstances that have an influence on the period of embryonic development. In the *Amphioxus* the earliest and most important embryonic processes take place so rapidly that the *blastula* is formed in four hours, the *gastrula* in six, and the typical vertebrate form in twenty-four.

In every case the duration of ontogeny links into the enormous period that has been necessary for phylogeny, or the gradual development of the ancestral series. This period is not measured by years or centuries, but by thousands and millions of years. Many millions of years had to pass before the most advanced

vertebrate, man, was evolved, step by step, from his ancient unicellular ancestors. The opponents of evolution, who declare that this gradual development of the human form from lower animal forms, and ultimately from a unicellular organism, is an incredible miracle, forget that the same miracle takes place within the space of nine months in the embryonic development of every human being. Each of us has, in the forty weeks—properly speaking, in the first four weeks—of his development in the womb, passed through the same series of transformations that our animal ancestors underwent in the course of millions of years.

It is impossible to determine even approximately, in hundreds or even thousands of years, the real and absolute duration of the phylogenetic period. But for some time now we have, through the research of geologists, been in a position to assign the relative length of the various sections of the organic history of the earth. The immediate data for determining this relative length of the geological periods are found in the thickness of the sedimentary strata—the strata that have been formed at the bottom of the sea or in fresh water from the mud or slime deposited there. These successive layers of limestone, sandstone, slate, marl, etc., which make up the greater part of the rocks, and are often several thousand feet thick, give us a standard for computing the relative length of the various periods.

To make the point quite clear, I must say a word about the evolution of the earth in general, and point out briefly the chief features of the story. In the first place, we encounter the principle that on our planet organic life began to exist at a definite period. That statement is no longer disputed by any competent geologist or biologist. The organic history of the earth could not commence until it was possible for water to settle on our planet in fluid condition. Every organism, without exception, needs fluid water as a condition of existence, and contains a considerable quantity of it. Our own body, when fully formed, contains sixty to seventy per cent. of water in its tissues, and only thirty to forty per cent. of solid matter. There is even more water in the body of the child, and still more in the embryo. In the earlier stages of development the human foetus contains more than ninety per cent. of water, and not ten per cent. of solids. In the lower marine animals,

especially certain medusae, the body consists to the extent of more than ninety-nine per cent. of sea-water, and has not one per cent. of solid matter. No organism can exist or discharge its functions without water. No water, no life!

But fluid water, on which the existence of life primarily depends, could not exist on our planet until the temperature of the surface of the incandescent sphere had sunk to a certain point. Up to that time it remained in the form of steam. But as soon as the first fluid water could be condensed from the envelope of steam, it began its geological action, and has continued down to the present day to modify the solid crust of the earth. The final outcome of this incessant action of the water—wearing down and dissolving the rocks in the form of rain, hail, snow, and ice, as running stream or boiling surge—is the formation of mud. As Huxley says in his admirable *Lectures on the Causes of Phenomena in Organic Nature*, the chief document as to the past history of our earth is mud; the question of the history of past ages resolves itself into a question about the formation of mud.

As I have said, it is possible to form an approximate idea of the relative age of the various strata by comparing them at different parts of the earth's surface. Geologists have long been agreed that there is a definite historical succession of the different strata. The various superimposed layers correspond to successive periods in the organic history of the earth, in which they were deposited in the form of mud at the bottom of the sea. The mud was gradually converted into stone. This was lifted out of the water owing to variations in the earth's surface, and formed the mountains. As a rule, four or five great divisions are distinguished in the organic history of the earth, corresponding to the larger and smaller groups of the sedimentary strata. The larger periods are then sub-divided into a series of smaller ones, which usually number from twelve to fifteen. The comparative thickness of the groups of strata enables us to make an approximate calculation of the relative length of these various periods of time. We cannot say, it is true, "In a century a stratum of a certain thickness (about two feet) is formed on the average; therefore, a layer 1,000 feet thick must be 500,000 years old." Different strata of the same thickness may need very different periods for their formation. But from

DURATION OF THE HISTORY OF OUR STEM

the thickness or size of the stratum we may draw some conclusion as to the relative length of the period.

The first and oldest of the four or five chief divisions of the organic history of the earth is called the primordial, archaic, or archeozoic period. If we compute the total average thickness of the sedimentary strata at about 120,000 feet, this first

(7.6) indicate—possibly 9.6. Of late years the thickness of the archaic rocks has been put at 90,000 feet.

The primordial period falls into three subordinate sections—the Laurentian, Huronian, and Cambrian, corresponding to the three chief groups of rocks which comprise the archaic formation. The immense period during which these

**SYNOPSIS OF THE PALEONTOLOGICAL FORMATIONS,
OR THE FOSSILIFEROUS STRATA OF
THE CRUST**

[illegible]

period comprises 70,000 feet, or the greater part of the whole. For this and other reasons we may at once conclude that the corresponding primordial or archæolithic period must have been in itself much longer than the whole of the remaining periods together, from its close to the present day. It was probably much longer than the figures I have quoted

were forming in the primitive ocean probably comprises more than 50,000,000 years. At the commencement of it the oldest and simplest organisms were formed by spontaneous generation—the Monera, with which the history of life on our planet opened. From these were first developed unicellular organisms of the simplest character, the Protozoa.

and Protozoa (pseudotomes, amoebae, rhizopods, infusoria, and other Protists). During this period the whole of the invertebrate ancestors of the human race were evolved from the unicellular organisms. We can deduce this from the fact that we already find remains of fossilized fishes (Selachii and Ganoids) towards the close of the following Silurian period. These are much more advanced and much younger than the lowest vertebrates, the Amphioxus, and the numerous skull-less vertebrates, related to the Amphioxus, that must have lived at that time. The whole of the invertebrate ancestors of the human race must have preceded these.

The primordial age is followed by a much shorter division, the *paleozoic* or *Primary* age. It is divided into four long periods, the Silurian, Devonian, Carboniferous, and Permian. The Silurian strata are particularly interesting because they contain the first fossil traces of vertebrates—teeth and scales of Selachii (*Palaodus*) in the lower, and Ganoids (*Pteraspis*) in the upper Silurian. During the Devonian period the "old red sandstone" was formed; during the Carboniferous period were deposited the vast coal-measures that yield us our chief combustible material, in the Permian (or the *Dyas*). In fine, the new red sandstone, the Zechstein (magnesian limestone), and the Kupferschiefer (marl-slate) were formed. The collective depth of these strata is put at 30,000 to 45,000 feet. In any case, the paleozoic age, taken as a whole, was much shorter than the preceding and much longer than the subsequent periods. The strata that were deposited during this primary epoch contain a large number of fossils; besides the invertebrate species there are a good many vertebrates, and the fishes preponderate. There were so many fishes, especially primitive fishes (of the shark type) and plated fishes, during the Devonian, and also during the Carboniferous and Permian periods, that we may describe the whole paleozoic period as "the age of fishes." Among the paleozoic plated fishes or Ganoids the *Crossopterygii* and the *Ctenodipterina* (dipnoists) are of great importance.

During this period some of the fishes began to adapt themselves to living on land, and so gave rise to the class of the amphibians. We find in the Carboniferous period fossilized remains of five-toed amphibians, the oldest terrestrial, air-breathing vertebrates. These amphibians

increase in variety in the Permian epoch. Towards the close of it we find the first Amniotes, the ancestors of the three higher classes of Vertebrates. These are lizard-like animals; the first to be discovered was the *Proteropsaurus*, from the marl at Eisenach. The rise of the earliest Amniotes, among which must have been the common ancestor of the reptiles, birds, and mammals, is put back towards the close of the paleozoic age by the discovery of these reptile remains. The ancestors of our race during this period were at first represented by true fishes, then by dipnoists and amphibians, and finally by the earliest Amniotes, or the *Protamniotes*.

The third chief section of the organic history of the earth is the *Mesozoic* or *Secondary* period. This again is subdivided into three divisions: Triassic, Jurassic, and Cretaceous. The thickness of the strata that were deposited in this period, from the beginning of the Triassic to the end of the Cretaceous period, is altogether about 15,000 feet, or not half as much as the paleozoic deposits. During this period there was a very brisk and manifold development in all branches of the animal kingdom. There were especially a number of new and interesting forms evolved in the vertebrate stem. Bony fishes (*Tetrapods*) make their first appearance. Reptiles are found in extraordinary variety and number; the extinct giant-serpents (*diapsauria*), the sea-serpents (*thalisauria*), and the flying lizards (*pterosauria*) are the most remarkable and best known of these. On account of this predominance of the reptile-class, the period is called "the age of reptiles." But the bird-class was also evolved during this period; they certainly originated from some division of the lizard-like reptiles. This is proved by the embryological identity of the birds and reptiles and their comparative anatomy, and, among other features, from the circumstance that in this period there were birds with teeth in their jaws and with tails like lizards (*Archaeopteryx*, *Odontornis*).

Finally, the most advanced and (for us) the most important class of the vertebrates, the mammals, made their appearance during the mesozoic period. The earliest fossil remains of them were found in the latest Triassic strata—lower jaws of small ungulates and mammals. More numerous remains are found a little later

in the Jurassic, and some in the Cretaceous. All the mammal remains that we have from this section belong to the lower plesiomammals and marsupials; among these were most certainly the ancestors of the human race. On the other hand, we have not found a single indisputable fossil of any higher mammal (a placental) in the whole of this period. This division of the mammals, which includes man, was not developed until later, towards the close of this or in the following period.

The fourth section of the organic history of the earth, the Tertiary or Cenozoic age, was much shorter than the preceding. The strata that were deposited during this period have a collective thickness of only about 3,000 feet. It is subdivided into four sections—the Eocene, Oligocene, Miocene, and Pliocene. During these periods there was a very varied development of higher plant and animal forms, the fauna and flora of our planet approached nearer and nearer to the character that they bear to-day. In particular, the most advanced class, the mammals, began to preponderate. Hence the Tertiary period may be called "the age of mammals." The highest section of this class, the placentals, now made their appearance, to this group the human race belongs. The first appearance of man, or, to be more precise, the development of man from some closely-related group of apes, probably falls in either the miocene or the pliocene period, the middle or the last section of the Tertiary period. Others believe that man properly so-called—man endowed with speech—was not evolved from the non-speaking ape-man (*Pithecanthropus*) until the following, the anthropozoic, age.

In this fifth and last section of the organic history of the earth we have the full development and dispersion of the various races of men, and so it is called the *Anthropozoic* as well as the *Quaternary* period. In the imperfect condition of paleontological and ethnographical science we cannot as yet give a confident answer to the question whether the evolution of the human race from some extinct ape or lemur took place at the beginning of this or towards the middle or the end of the Tertiary period. However, this much is certain: The development of civilisation falls in the anthropozoic age, and this is merely an insignificant fraction of the vast period of the whole history of life. When we

remember this, it seems ridiculous to restrict the word "history" to the civilised period. If we divide into a hundred equal parts the whole period of the history of life, from the spontaneous generation of the first bacteria to the present day, and if we then represent the relative duration of the five chief sections or ages, as calculated from the average thickness of the strata they contain, as percentages of this, we get something like the following relation:—

I. Archeozoic or archæozoic (primordial) age	53	6
II. Paleozoic or paleozoic (primary) age	32	1
III. Mesozoic or mesozoic (secondary) age	11	5
IV. Cenozoic or cenozoic (tertiary) age	2	3
V. Anthropozoic or anthropozoic (quaternary) age	0	5

100 0

In any case, the "historical period" is an insignificant quantity compared with the vast length of the preceding ages, in which there was no question of human existence on our planet. Even the important Cenozoic or Tertiary period, in which the first placentals or higher mammals appear, probably amounts to little over two per cent. of the whole organic age.

Before we approach our proper task, and, with the aid of our ontogenetic acquirements and the biogenetic law, follow step by step the paleontological development of our animal ancestors, let us glance for a moment at another, and apparently quite remote, branch of science, a general consideration of which will help us in the solving of a difficult problem. I mean the science of comparative philology. Since Darwin gave new life to biology by his theory of selection, and raised the question of evolution on all sides, it has often been pointed out that there is a remarkable analogy between the development of languages and the evolution of species. The comparison is perfectly just and very instructive. We could hardly find a better analogy when we are dealing with some of the difficult and obscure features of the evolution of species. In both cases we find the action of the same natural laws.

All philologists of any competence in their science now agree that all human languages have been gradually evolved from very rudimentary beginnings. The

idea that speech is a gift of the gods—an idea held by distinguished authorities only fifty years ago—is now generally abandoned, and only supported by theologians and others who admit no natural development whatever. Speech has been developed simultaneously with its organs, the larynx and tongue, and with the functions of the brain. Hence it will be quite natural to find in the evolution and classification of languages the same features as in the evolution and classification of organic species. The various groups of languages that are distinguished in philology as primitive, fundamental, parent, and daughter languages, dialects, etc., correspond entirely in their development to the different categories which we classify in zoology and botany as stems, classes, orders, families, genera, species, and varieties. The relation of these groups, partly co-ordinate and partly subordinate, in the general scheme is just the same in both cases, and the evolution follows the same lines in both.

When, with the assistance of this tree, we follow the formation of the various languages that have been developed from the common root of the ancient Indo-Germanic tongue, we get a very clear idea of their phylogeny. We shall see at the same time how analogous this is to the development of the various groups of vertebrates that have arisen from the common stem-form of the primitive vertebrate. The ancient Indo-Germanic root-language divided first into two principal stems—the Slavo-Germanic and the Aryo-Romanic. The Slavo-Germanic stem then branches into the ancient Germanic and the ancient Slavo-Lettic tongues; the Aryo-Romanic into the ancient Aryan and the ancient Greco-Roman. If we still follow the genealogical tree of these four Indo-Germanic tongues, we find that the ancient Germanic divides into three branches—the Scandinavian, the Gothic, and the German. From the ancient German came the High German and Low German; to the latter belong the Frisian, Saxon, and modern Low-German dialects. The ancient Slavo-Lettic divided first into a Baltic and a Slav language. The Baltic gave rise to the Lett, Lithuanian, and old-Prussian varieties; the Slav to the Russian and South-Slav in the south-east, and to the Polish and Czech in the west.

We find an equally prolific branching of its two chief stems when we turn to

the other division of the Indo-Germanic languages. The Greco-Roman divided into the Thracian (Albano-Greek) and the Italo-Celtic. From the latter came the divergent branches of the Italic (Roman and Latin) in the south, and the Celtic in the north. From the latter have been developed all the British (ancient British, ancient Scotch, and Irish) and Gallic varieties. The ancient Aryan gave rise to the numerous Iranian and Indian languages.

This "comparative anatomy" and evolution of languages admirably illustrates the phylogeny of species. It is clear that in structure and development the primitive languages, mother and daughter languages, and varieties, correspond exactly to the classes, orders, genera, and species of the animal world. In both cases the "natural" system is phylogenetic. As we have been convinced from comparative anatomy and ontogeny, and from paleontology, that all past and living vertebrates descend from a common ancestor, so the comparative study of dead and living Indo-Germanic tongues proves beyond question that they are all modifications of one primitive language. This view of their origin is now accepted by all the chief philologists who have worked in this branch and are unprejudiced.

But the point to which I desire particularly to draw the reader's attention in this comparison of the Indo-Germanic languages with the branches of the vertebrate stem is, that one must never confuse direct descendants with collateral branches, nor extinct forms with living. This confusion is very common, and our opponents often make use of the erroneous ideas it gives rise to for the purpose of attacking evolution generally. When, for instance, we say that man descends from the ape, this is from the lemur, and the lemur from the marsupial, many people imagine that we are speaking of the living species of those orders of mammals that they find stuffed in our museums. Our opponents then foist this idea on us, and say, with more astuteness than intelligence, that it is quite impossible; or they ask us, by way of physiological experiment, to turn a kangaroo into a lemur, a lemur into a gorilla, and a gorilla into a man! The demand is childish, and the idea it rests on erroneous. All these living forms have diverged more or less from the ancestral form; none of them could engender the

same posterity that the stem-form really produced thousands of years ago.

It is certain that man has descended from some extinct mammal; and we should just as certainly class this in the order of apes if we had it before us. It is equally certain that this primitive ape descended in turn from an unknown lemur, and this from an extinct marsupial. But it is just as clear that all these extinct ancestral forms can only be claimed as belonging to the living order of mammals in virtue of their essential internal structure and their resemblance in the decisive anatomic characteristics of each order. In external appearance, in the characteristics of the *genus* or *species*, they would differ more or less, perhaps very considerably, from all living representatives of those orders. It is a universal and natural procedure in phylogenetic development that the stem-forms themselves, with their specific peculiarities, have been extinct for some time. The forms that approach nearest to them among the living species are more or less — perhaps very substantially — different from them. Hence in our phylogenetic inquiry and in the comparative study of the living, divergent descendants, there can only be a question of determining the greater or less remoteness of the latter from the ancestral form. Not a single one of the older stem-forms has continued unchanged down to our time.

We find just the same thing in comparing the various dead and living languages that have developed from a common primitive tongue. If we examine our genealogical tree of the Indo-Germanic languages in this light, we see at once that all the older or parent tongues, of which we regard the living varieties of the stem as divergent daughter or grand-daughter languages, have been extinct for some time. The Aryan-Romantic and the Slavo-Germanic tongues have completely disappeared; so also the Aryan, the Græco-Roman, the Slavo-Lettic, and the ancient Germanic. Even their daughters and grand-daughters have been lost; all the living Indo-Germanic languages are only related in the sense that they are divergent descendants of common stem-forms. Some forms have diverged more, and some less, from the original stem-form.

This easily demonstrable fact illustrates very well the analogous case of the origin of the vertebrate species. Phylogenetic

comparative philology here yields a strong support to phylogenetic comparative zoology. But the one can adduce more direct evidence than the other, as the paleontological material of philology — the old monuments of the extinct tongue — have been preserved much better than the paleontological material of zoology, the fossilised bones and imprints of vertebrates.

We may, however, trace man's genealogical tree not only as far as the lower mammals, but much further — to the amphibia, to the shark-like primitive fishes, and, in fine, to the skull-less vertebrates that closely resembled the *Amphioxus*. But this must not be understood in the sense that the existing *Amphioxus*, or the sharks or amphibia of to-day, can give us any idea of the external appearance of these remote stem-forms. Still less must it be thought that the *Amphioxus* or any actual shark, or any living species of amphibia, is a real ancestral form of the higher vertebrates and man. The statement can only rationally mean that the living forms I have referred to are *collateral lines* that are much more closely related to the extinct stem-forms, and have retained the resemblance much better, than any other animals we know. They are still so like them in regard to their distinctive internal structure that we should put them in the same class with the extinct forms if we had those before us. But no direct descendants of these earlier forms have remained unchanged. Hence we must entirely abandon the idea of finding direct ancestors of the human race in their characteristic *external form* among the living species of animals. The essential and distinctive features that still connect living forms more or less closely with the extinct common stem-forms lie in the internal structure, not the external appearance. The latter has been much modified by adaptation. The former has been more or less preserved by heredity.

Comparative anatomy and ontogeny prove beyond question that man is a true vertebrate, and, therefore, man's special genealogical tree must be connected with that of the other Vertebrates, which spring from a common root with him. But we have also many important grounds in comparative anatomy and ontogeny for assuming a common origin for all the Vertebrates. If the general theory of

evolution is correct, all the Vertebrates, including man, come from a single common ancestor, a long-extinct "Primitive Vertebrate." Hence the genealogical tree of the Vertebrates is at the same time that of the human race.

Our task, therefore, of constructing man's genealogy becomes the larger aim of discovering the genealogy of the entire vertebrate stem. As we now know from the comparative anatomy and ontogeny of the Amphioxus and the Ascidia, this is in turn connected with the genealogical tree of the Lowervertebrates (directly with that of the Vermalia), but has no direct connection with the independent stems of the Articulates, Molluscs, and Echinoderms. If we do thus follow our ancestral tree through various stages down to the lowest worms, we come inevitably to the *Gastrea*, that most instructive form that gives the clearest possible picture of an animal with two germinal layers. The *Gastrea* itself has originated from the simple multicellular vesicle, the *Blasodea*, and this in turn must have been evolved from the lowest circle of unicellular animals, to which we give the name of Protozoa. We have already considered the most important primitive type of these, the unicellular *Amœba*, which is extremely instructive when compared with the human ovum. With this we reach the lowest of the solid data to which we are to apply our biogenetic law, and by which we may deduce the extinct ancestor from the embryonic form. The amœboid nature of the young ovum and the unicellular condition in which (as stem-cell or cytula) every human being begins its existence justify us in affirming that the earliest ancestors of the human race were simple amœboid cells.

But the further question now arises: "Whence came these first amœbæ with which the history of life began at the commencement of the Laurentian epoch?" There is only one answer to this. The earliest unicellular organisms can only have been evolved from the simplest organisms we know, the *Micæra*. These are the simplest living things that we can conceive. Their whole body is nothing but a particle of plasma, a granule of living albuminous matter, discharging of itself all the essential vital functions that form the material basis of life. Thus we come to the last, or, if you prefer, the first, question in connection with evolution—the question of the origin of the *Micæra*.

This is the real question of the origin of life, or of spontaneous generation.

We have neither space nor occasion to go further in this Chapter into the question of spontaneous generation. For this I must refer the reader to the fifteenth chapter of the *History of Creation*, and especially to the second book of the *General Morphology*, or to the essay on "The *Micæra* and Spontaneous Generation" in my *Studies of the Micæra and other Protists*.¹ I have given there fully my own view of this important question. The famous botanist Nagell afterwards (1884) developed the same ideas. I will only say a few words here about this obscure question of the origin of life, in so far as our main subject, organic evolution in general, is affected by it. Spontaneous generation, in the definite and restricted sense in which I maintain it, and claim that it is a necessary hypothesis in explaining the origin of life, refers solely to the evolution of the *Micæra* from inorganic carbon-compounds. When living things made their first appearance on our planet, the very complex nitrogenous compound of carbon that we call *plasma*, which is the earliest material embodiment of vital action, must have been formed in a purely chemical way from inorganic carbon-compounds. The first *Micæra* were formed in the sea by spontaneous generation, as crystals are formed in the mother-water. Our demand for a knowledge of causes compels us to assume this. If we believe that the whole inorganic history of the earth has proceeded on mechanical principles without any intervention of a Creator, and that the history of life also has been determined by the same mechanical laws, if we see that there is no need to admit creative action to explain the origin of the various groups of organisms; if it is utterly irrational to assume such creative action in dealing with the first appearance of organic life on the earth.

This much-disputed question of "spontaneous generation" seems so obscure, because people have associated with the term a mass of very different, and often very absurd, ideas, and have attempted to solve the difficulty by the crudest experiments. The real doctrine of the spontaneous generation of life cannot possibly be refuted by experiments.

¹ The English reader will find a business and up-to-date chapter on this subject in Huxley's recently-written and translated *Wonders of Life*.—T. J. W.

Every experiment that has a negative result only proves that no organism has been formed out of inorganic matter in the conditions—highly artificial conditions—we have established. On the other hand, it would be exceedingly difficult to prove the theory by way of experiment; and even if Monera were still formed daily by spontaneous generation (which is quite possible), it would be very difficult, if not impossible, to find a solid proof of it. Those who will not admit the spontaneous generation of the first living things in our sense must have recourse to a supernatural miracle; and this is, as a matter of fact, the desperate resource to which our "exact" scientists resort.

A famous English physicist, Lord Kelvin (then Sir W. Thomson), attempted to dispense with the hypothesis of spontaneous generation by assuming that the organic inhabitants of the earth were developed from germs that came from the inhabitants of other planets, and that chance to fall on our planet on fragments of their original home, or meteorites. This hypothesis found many supporters, among others the distinguished German physicist, Helmholtz. However, it was refuted in 1872 by the able physicist, Friedrich Zöllner, of Leipzig, in his work,

On the Nature of Comets. He showed clearly how unscientific this hypothesis is; firstly in point of logic, and secondly in point of scientific content. At the same time he pointed out that our hypothesis of spontaneous generation is "a necessary condition for understanding nature according to the law of causality."

I repeat that we must call in the aid of the hypothesis only as regards the Monera, the structureless "organisms without organs." Every complex organism must have been evolved from some lower organism. We must not assume the spontaneous generation of even the simplest cell, for this itself consists of at least two parts—the internal, firm nuclear substance, and the external, softer cellular substance or the protoplasm of the cell-body. These two parts must have been formed by differentiation from the indifferent plasma of a moneron, or a cytode. For this reason the natural history of the Monera is of great interest; here alone can we find the means to overcome the chief difficulties of the problem of spontaneous generation. The actual living Monera are specimens of such organless or structureless organisms, as they must have been formed by spontaneous generation at the commencement of the history of life.

CHAPTER XIX.

OUR PROTIST ANCESTORS

UNDER the guidance of the biogenetic law—the basis of the evidence we

are the ...
Phylogeny

as a whole is an inductive process—the totality of the biological processes in the life of plants, animals, and man we have gathered a confident inductive idea that the whole organic population of our planet has been moulded on a harmonious law of evolution. All the interesting

phenomena that we meet in ontogeny and paleontology, comparative anatomy and cytology, the distribution and habits of organisms—all the important general laws that we abstract from the phenomena of these sciences, and combine in harmonious unity—are the broad bases of our great biological induction.

But when we come to the application of this law, and seek to determine with its aid the origin of the various species of organisms, we are compelled to frame

hypotheses that have essentially a *deductive* character, and are inferences from the general law to particular cases. But these special deductions are just as much justified and necessitated by the rigorous laws of logic as the inductive conclusions on which the whole theory of evolution is built. The doctrine of the animal ancestry of the human race is a special deduction of this kind, and follows with logical necessity from the general inductive law of evolution.

I must point out at once, however, that the certainty of these evolutionary hypotheses, which rest on clear special deductions, is not always equally strong. Some of these inferences are now beyond question; in the case of others it depends on the knowledge and the competence of the inquirer what degree of certainty he attributes to them. In any case, we must distinguish between the *absolute* certainty of the general (inductive) theory of descent and the *relative* certainty of special (deductive) evolutionary hypotheses. We can never determine the whole ancestral series of an organism with the same confidence with which we hold the general theory of evolution as the sole scientific explanation of organic modifications. The special indication of stem-forms in detail will always be more or less incomplete and hypothetical. This is quite natural. The evidence on which we build is imperfect, and always will be imperfect; just as in comparative philology.

The first of these documents, paleontology, is exceedingly incomplete. We know that all the fossils yet discovered are only an insignificant fraction of the plants and animals that have lived on our planet. For every single species that has been preserved for us in the rocks there are probably hundreds, perhaps thousands, of extinct species that have left no trace behind them. This extreme and very unfortunate incompleteness of the paleontological evidence, which cannot be pointed out too often, is easily explained. It is absolutely inevitable in the circumstances of the fossilization of organisms. It is also due in part to the incompleteness of our knowledge in this branch. It must be borne in mind that the great majority of the stratified rocks that compose the crust of the earth have not yet been opened. We have only a few specimens of the innumerable shells that are buried in the vast mountain ranges of Asia and Africa.

Only a part of Europe and North America has been investigated carefully. The whole of the fossils known to us certainly do not amount to a hundredth part of the remains that are really buried in the crust of the earth. We may, therefore, look forward to a rich harvest in the future as regards this science. However, our paleontological evidence will (for reasons that I have fully explained in the sixteenth chapter of the *History of Creation*) always be defective.

The second chief source of evidence, ontogeny, is not less incomplete. It is the most important source of all for special phylogeny; but it has great defects, and often fails us. We must, above all, clearly distinguish between polygenetic and conogenetic phenomena. We must never forget that the laws of curtailed and disturbed heredity often make the original course of development almost unrecognizable. The recapitulation of phylogeny by ontogeny is only fairly complete in a few cases, and is never whole complete. As a rule, it is precisely the earliest and most important embryonic stages that suffer most from alteration and condensation. The earlier embryonic forms have had to adapt themselves to new circumstances, and so have been modified. The struggle for existence has had just as profound an influence on the freely moving and still immature young forms as on the adult forms. Hence in the embryology of the higher animals, especially, polygenosis is much restricted by conogenosis; it is to-day, as a rule, only a faded and much altered picture of the original evolution of the animal's ancestors. We can only draw conclusions from the embryonic forms to the stem-history with the greatest caution and discrimination. Moreover, the embryonic development itself has only been fully studied in a few species.

Finally, the third and most valuable source of evidence, comparative anatomy, is also, unfortunately, very imperfect; for the simple reason that the whole of the living species of animals are a mere fraction of the vast population that has dwelt on our planet since the beginning of life. We may confidently put the total number of these at more than a million species. The number of animals whose organisation has been studied up to the present in comparative anatomy is proportionately very small. Here, again, future research will yield incalculable treasures.

But, for the present, in view of this patent incompleteness of our chief sources of evidence, we must naturally be careful not to lay too much stress in human phylogeny on the particular animals we have studied, or regard all the various stages of development with equal α -form.

In my first efforts to construct the series of man's ancestors I drew up a list of, at first ten, afterwards twenty to thirty, forms that may be regarded more or less certainly as animal ancestors of the human race, or as stages that in a sense mark off the chief sections in the long story of evolution from the unicellular organism to man. Of these twenty to thirty stages, ten to twelve belong to the older group of the Invertebrates and eighteen to twenty to the younger division of the Vertebrates.

In approaching, now, the difficult task of establishing the evolutionary succession of these thirty ancestors of humanity since the beginning of life, and in venturing to lift the veil that covers the earliest secrets of the earth's history, we must undoubtedly look for the first living things among the wonderful organisms that we call the Monera; they are the simplest organisms known to us—in fact, the simplest we can conceive. Their whole body consists merely of a simple particle or globule of structureless plasma or plasmon. The

distinguish between the plasmon of the cytodas and the protoplasm of the cells. This distinction is of the utmost importance for the purposes of evolution. As I have often said, we must recognize two different stages of development in these "elementary organisms," or plastids ("builders"), that represent the ultimate units of organic individuality. The earlier and lower stage are the unnucleated cytodas, the body of which consists of only one kind of albuminous matter—the homogeneous plasmon or "formative matter." The later and higher stage are the nucleated cells, in which we find a differentiation of the original plasmon into two different formative substances—the caryoplasm of the nucleus and the cytoplasm of the body of the cell (cf. pp. 37 and 42).

The Monera are permanent: Their whole body consists of soft structureless plasmon. However carefully we examine it with our finest chemical reagents and most powerful microscopes,



FIG. 26.—*Chromococcus ruber* (Nagel), magnified 1,000 times. A phytomonadon, the globular plastid of which secretes a gelatinous structureless mesoplasm. The unnucleated globule of plasma (blue-green color) increases in simple division (and).

discoveries of the last four decades have led us to believe with increasing certainty that wherever a natural body exhibits the vital processes of nutrition, reproduction, voluntary movement, and sensation, we have the action of a nitrogenous carbon-compound of the chemical group of the albuminoids, this plasma (or protoplasm) is the material basis of all vital functions. Whether we regarded the function, in the monistic sense, as the direct action of the material substratum, or whether we take matter and force to be distinct things in the dualistic sense, it is certain that we have not as yet found any living organism in which the exercise of the vital functions is not inseparably bound up with plasma.

The soft slimy plasmon of the body of the moneron is generally called "protoplasm," and identified with the cellular matter of the ordinary plant and animal cells. But we must, to be accurate,

we can find no definite parts or no anatomic structure to it. Hence, the Monera are literally organisms without organs; in fact, from the philosophic point of view they are not organisms at all, since they have no organs. They can only be called organisms in the sense that they are capable of the vital functions of nutrition, reproduction, sensation, and movement. If we were to try to imagine the simplest possible organism, we should frame something like the moneron.


The Monera that we find to-day in various forms fall into two groups according to the nature of their nutrition—the *Phytomonera* and the *Zoomonera*; from the physiological point of view, the former are the simplest specimens of the plant (*phytes*) kingdom, and the latter of the animal (*zoes*) world. The *Phytomonera*, especially in their simplest form, the *Chromococcus* (*Phycocchromococcus* or *Cyanophyceae*), are the most primitive and the

oldest of living organisms. The typical genus *Chloococcus* (Fig. 226) is represented by several fresh-water species, and often forms a very delicate bluish-green deposit

..... and wood in ponds and ditches, of round, light green particles, from of an inch in diameter.

The whole life of these homogeneous globules of plasma consists of simple growth and reproduction by cleavage.

When the tiny particle has reached certain size by the continuous of inorganic matter, it divides equal halves, by a constriction middle. The two daughter formed immediately begin



of cytoblast membrane

similar vital process. It is the same with the brown *Prorocentrum* (formerly called the *Prorococcus marinus*); it forms large masses of floating matter in the arctic seas. The tiny plasma-globules of species are of a greenish-brown colour,

..... if an able

the simplest *Chloococcales*, but we find one in other members of the same family; in *Aphanizomenon* (Fig. 227) the enveloping of the plastids combine; in *Chloococcus* they are retained through several generations, so that the little plasma-globules are enfolded in many layers of membrane.

Next to the *Chloococcales* come the Bacteria, which have been evolved from them by the remarkable change in nutrition which gives us the simple explanation of the difference of plant

the protist kingdom. The *Chloococcus* build up their plasma directly from organic matter; the Bacteria feed on organic matter. Hence, if we logically divide the protist kingdom into plasma-forming Protophyta and plasma-consuming Protococcus, we must class the Bacteria with the latter; it is quite illogical to describe them—as is still often done—as *Schizomycetes*, and class them with the true fungi. The Bacteria, like the *Chloococcales*, have no nucleus. As is well known, they play an important part in modern biology as the causes of fermentation and putrefaction, and of tuberculosis, typhus, cholera, and other infectious diseases, and as parasites, etc. But we cannot linger now to deal with those very interesting features; the Bacteria have no relation to man's genealogical tree.

We may now turn to consider the remarkable *Protamoeba*, or un-nucleated Amoeba. I have, in the first volume, pointed out the great importance of the ordinary Amoeba in connection with several weighty questions of general biology. The tiny *Protamoeba*, which are found both in fresh and salt water, have the same unshapely form and irregular movements of their simple naked body as the real Amoeba, but they differ from them very materially in having no nucleus in their cell-body. The short, blunt, finger-like processes that are thrust out at the surface of the creeping *Protamoeba* serve for getting food as well as for locomotion. They multiply by simple cleavage (Fig. 228).

The next stage to the simple cytoblasts of the Monera in the genealogy of mankind (and all other animals) is the simple cell, or the most rudimentary form of the cell which we find living independently to-day as the Amoeba. The earliest process of inorganic differentiation in the structureless body of the Monera led to into two different substances—the karyoplasm and the cytoplasm. The karyoplasm is the inner and firmer part of the cell, the substance of the nucleus. The cytoplasm is the outer and softer part, the substance of the body of the cell. By this important differentiation of the plasma into nucleus and cell-body, the

organized cell was evolved from the structureless cytozoa, the nucleated from the un-nucleated plasmod. That the first cells to appear on the earth were formed from the Monera by such a differentiation seems to us the only possible view in the present condition of science. We have a direct instance of this earliest process of differentiation to-day in the ontogeny of many of the lower Protists (such as the Gregarinas).

The unicellular form that we have in the ovum has already been described as the reproduction of a corresponding unicellular stem-form, and to this we have ascribed the organisation of an Amoeba (cf. Chapter VI.). The irregular-shaped Amoeba, which we find living independently to-day in our fresh and salt water, is the least definite and the most primitive of all (Fig. 16).

the unripe ova (the protists that we find in the ovaries of animals) cannot be distinguished from the common Amoeba, we must regard the Amoeba as the primitive form that is reproduced in the embryonic stage of the amoeboid ovum to-day, in accordance with the biogenetic law. I have already pointed out, in proof of the striking resemblance of the two cells, that the ova of many of the sponges were formerly regarded as parasitic Amoebae (Fig. 18). Large unicellular organisms like the Amoeba were found creeping about inside the body of the sponge, and were thought to be parasites. It was afterwards discovered that they were really the ova of the sponge from which the embryos were developed. As a matter of fact, these sponge-ova are so much like many of the Amoebae in size, shape, the character of their nucleus, and movement of the pseudopodia, that it is impossible to distinguish them without knowing their subsequent development.

relation of the
of it to some
ant amoeboid ancestral form, supply
the answer to the old problem: "Which
was first, the egg or the chick?" We
can now give a very plain answer to this
riddle, with which our opponents have
often tried to drive us into a corner. The
egg came a long time before the chick.
We do not mean, of course, that the egg
existed from the first as a bird's egg, but
as an indifferent amoeboid cell of the
simplest character. The egg lived for
thousands of years as an independent

unicellular organism, the Amoeba. The
egg, in the modern physiological sense of
the word, did not make its appearance
until the descendants of the unicellular
Protistoon had developed into multicellular
animals, and these had undergone sexual
differentiation. Even then the egg was
first a gastraea-egg, then a platoda-egg,
then a termalia-egg, and chordonia-egg;
later still a crura-egg, then fish-egg,
amphibia-egg, reptile-egg, and finally
bird's egg. The bird's egg we have ex-
perience of daily is a highly complicated

product, the result of countless
hereditary processes that have taken place
of years

The simple Protophyta, and from these our
protist ancestors were developed after-
wards the morphological point of
view both the vegetal and the animal
Protists were simple organisms, indi-




FIG. 16.—A mossopron (*Frustramochia*) in the act of reproduction. A The whole mossopron, showing how an ordinary amoeba by throwing out chromatids produces. B It divides into two halves by a constriction in the middle. C The two halves separate, and each becomes an independent individual. (Huxley suggested.)

individuality of the first order, or plasmids.
The ancestors are complex organ-
isms, individuals of a higher order
—social aggregations of a plurality of
cells. The earliest of these, the *Metazoa*,
which represent the third stage in our
genealogy, are very simple associations of
homogeneous, indifferent cells — un-
differentiated colonies of social Amoebae or
Infusoria. To understand the nature and
origin of these problems

they follow step by step the first embryonic
products of the stem-cell. In all the
Metazoa the first embryonic process is
the repeated cleavage of the stem-cell, or
first segmentation-cell (Fig. 229). We
have already fully considered this process,
and found that all the different forms of
it may be reduced to one type, the original
equal or primordial segmentation (cf.
Chapter VIII.). In the genealogical tree

of the Vertebrates this pallogenic form of segmentation has been preserved in the *Amphioxus* alone, all the other Vertebrates having ontogenetically modified forms of cleavage. In any case, the latter were developed from the former, and so the segmentation of the ovum in the *Amphioxus* has a great interest for us (cf. Fig. 38). The outcome of this repeated cleavage is the formation of a round

small communities of *Amoebae* arise by the side of these ectoblastic Protocysts, the water-cells produced by cleavage remaining joined together. The advantages in the struggle for life which these communities had over the isolated cells favoured their formation and their further development. We find plenty of these cell-colonies or communities to-day in both fresh and salt water. They belong to various groups

Fig. 22.—Original or pallogenic ovum-cleavage. The stem-cell or cytula, formed by fission of the ovum, divides by repeated regular cleavage first into two (A), then four (B), then eight (C), and finally a large number of daughter-cells (D).

cluster of cells, composed of homogenous indifferent cells of the simplest character (Fig. 230). This is called the *morula* (= mulberry-embryo) on account of its resemblance to a mulberry or blackberry.

It is clear that this *morula* reproduces for us to-day the simple structure of the multicellular animal that succeeded the unicellular amoeboid form in the early Laurentian period. In accordance with the biogenetic law, the *morula* recalls the ancestral form of the *Amoeba*, or simple colony of Protocysts. The first cell-com-

of our race that succeeded phylogener to the *Morula*, we have only to follow the further embryonic development of the *morula*. We then see that the social cells of the round cluster secrete a sort of jelly or a watery fluid inside their globular body, and they themselves rise to the surface of it (Fig. 29 F, G). In this way the solid mulberry-embryo becomes a hollow sphere, the wall of which is composed of a single layer of cells. We call this layer the *blastoderm*, and the sphere itself the *blastula*, or embryonic vesicle.

This intervening *blastula* is very important. The conversion of the *morula* into a hollow ball proceeds on the same lines originally in the most diverse stems—as, for instance, in many of the zoophytes and worms, the ascidians, many of the echinoderms and molluscs, and in the amphioxus. Moreover, in the animals in which we do not find a real pallogenic *blastula* the defect is clearly due to ontogenetic causes, such as the formation of food-yolk and other embryonic adaptations.

We may, therefore, the ontogenetic *blastula* is the reproduct of a very early phylogenetic stem-form, all the stems

stem-form, which was in the main constructed like the *blastula*. In many of the lower animals the *blastula* is not developed



FIG. 230.—*Morula*, or mulberry-shaped embryo.

unities to be formed, which laid the early foundation of the higher multicellular body, must have consisted of homogeneous and simple amoeboid cells. The oldest *Amoebae* lived isolated lives, and even the amoeboid cells that were formed by the segmentation of these unicellular organisms must have continued to live independently for a long time. But gradually

within the foetal membranes, but in the open water. In those cases each blastodermic cell begins at an early stage to thrust out one or more mobile hair-like processes; the body swims about by the vibratory movement of these lashes or whips (Fig. 29 F).

We still find, both in the fresh water, various kinds of primitive multicellular organisms that substantially the blastula in it may be in

hollow vesicles or gelatinous balls, with a wall composed of a single layer of ciliated homogeneous cells.

There is one kind even among the Protozoa—the familiar *Volvocina*, formerly classed with the *Volvox*. *Globose* is found in the ponds in the spring—a small, green, gelatinous globule, swimming about by means of the strokes of its lashes, which rise in pairs from the

similar *Halosphaera viridis* also, which we find marine plankton (floating matter). number of green cells form a simple ball at the surface of the gelatinous ball; but in this case there are no cilia.

Some of the infusoria of the flagellata-class (*Stigma*, *Algaephora*, etc.) are similar in structure to these vegetal clusters, but differ in their animal nutrition; they form the special group of the *Catantaria*. In September, 1889, I studied the development of one of these graceful animals on the island of Gux-Oe, off the coast of Norway (*Algaephora planula*, Figs. 231 and 232). The fully-formed body is a gelatinous ball, with its wall composed of thin

After reaching maturity the community is dissolved. Each cell then lives independently. changes into a creeping amoeba afterwards contracts, and cloths with a structureless membrane. then looks just like an ordinary

When it has been in this condition for some time the cell divides into two, four, eight, sixteen, thirty-two, and sixty-four cells. These arrange themselves in a round vesicle, thrust out vibratory lashes, burst the capsule, and swim about in the sea

blastula, with similar construction, of many of the lower animals, we can confidently deduce from them that there was a very early and long-extinct common stem-form of substantially the same structure as the blastula. We may call it the *Blastae*. Its body consisted, when fully formed, of a simple hollow ball, filled with fluid or structureless jelly, with a wall composed of a single stratum of ciliated cells. There were probably many genera of these

Laurentian period, forming a special class of marine protists.

plant kingdom also the simple hollow sphere is found to be an elementary form of the multicellular organism. At the surface and below the surface (down to a depth of 2,000 yards) of the sea these green globules swimming about, with a wall composed of a single layer of chlorophyll-bearing cells. The botanist Schenck gave them the name of *Halosphaera viridis* in 1879.

The next stage to the *Blastula*, and the sixth in our genealogical tree, is the *Gastrula* that is developed from it. As we have already seen, this ancestral form is particularly important. That it once existed is proved with certainty by the *gastrula*, which we find temporarily in the ontogenesis of all the Metazoa (Fig. 29 J, A). As we saw, the original, palingenetic form of the *gastrula* is a round or oval uni-axial body, the simple cavity of which (the primitive gut) has an aperture at one pole of its axis (the primitive mouth). The wall of the gut consists of two strata of cells, and these are the primary germinal layers, the animal skin layer (ectoderm) and vegetal gut-layer (m).

Ontogenetic development of strata from the blastula furnishes evidence as to the phylogenetic of the *Gastrula* from the *Blastula*.

In the end this invagination goes so far that the invaginated part of the blastoderm lies close on the inner or non-invaginated part (Fig. 29 J). In explaining the phylogenetic origin of the *gastrula* in the light of this ontogenetic process, we may assume that the one-layered cell-community of the blastula began to take in food more largely at one particular part of its surface. Natural selection would gradually lead to the

formation of a depression or pit at this alimentary spot on the surface of the ball. The depression would grow deeper and deeper. In time the vegetal function of taking in and digesting food would be confined to the cells that lined this hole; the



FIG. 57.—The Norwegian *Magnaphora* phylla, swimming about by means of the tentacles at the surface.

other cells would see to the animal functions of locomotion, sensation, and protection. This was the first division of labour among the originally homogeneous cells of the blastula.

The effect, then, of this earliest histological differentiation was to produce two different kinds of cells—nutritive cells in the depression and locomotive cells on the surface outside. But this involved the severance of the two primary germinal layers—a most important process. When we remember that in an man's body, with all its various parts, and the body of all the other higher animals, are built up originally out of these two simple layers, we cannot lay too much stress on the phylogenetic significance of this gastrulation. In the simple primitive gut or gastric cavity of the gastrula and its rudimentary mouth we have the first real organ of the animal frame in the morphological sense; all the other organs were developed afterwards from these. In reality, the whole body of the gastrula is merely a "primitive gut." I have shown already (Chapters VIII. and IX.) that the two-layered embryo of all the Metazoa can be reduced to this typical gastrula. This important fact justifies us in concluding, in accordance with the biogenetic

law, that their ancestors also were phylogenetically developed from a similar stem-form. This ancient stem-form is the gastrula.

The gastrula probably lived in the sea during the Laurentian period, swimming about in the water by means of its ciliary coat much as free ciliated gastrulae do to-day. Probably it differed from the existing gastrula only in one essential point, though extinct millions of years ago. We have reason, from comparative anatomy and ontogeny, to believe that it multiplied by sexual generation, not solely asexually (by cleavage, gemma-

becan
perm. We base these hypotheses on the fact that we do to-day find the simple or sexual reproduction of living gastrulae and other lower animals, especially the sponges.

kinds of gastrula
a with an organoblast
at of the hypothetical

theory.
There are not very many species of living gastrulae; but their morphologic and phylogenetic interest is so great, an



FIG. 58.—A gastrula, showing the position of the germinal cells. The cells in the centre of the gastrula are connected by a fibrous process. Each cell has a contractile vacuole as well as a nucleus.

their intermediate position between the Protozoa and Metazoa so instructive, that I proposed long ago (1876) to make a special class of them, I distinguished three orders in this class—the *Gastrularia*, *Phaenaria*, and *Cyrenaria* (or *Dicyo-*

mida). But we might also regard these three orders as so many independent classes in a primitive gastræid stem.

The *Gastremaria* and *Cyemaria*, the chief of these living gastræids, are small Metazoa that live parasitically inside other Metazoa, and are, as a rule, $\frac{1}{4}$ to $\frac{1}{2}$ of an inch long, often much less (Fig. 233, 1-15). Their soft body, devoid of skeleton, consists of two simple strata of cells, the primary germinal layers; the outer of these is thickly clothed with long hair-like lashes, by which the parasites swim about in the various cavities of their host. The inner germinal layer furnishes the sexual products. The pure type of the original gastrula (or *archygastrella*, Fig. 29, 1) is seen in the *Pemmatodiscus gastrulaceus*, which Monticelli discovered in the umbrella of a large medusa (*Physalia physalis*) in 1895; the convex surface of this gelatinous umbrella was covered with numbers of clear vesicles, of $\frac{1}{4}$ to $\frac{1}{2}$ inch in diameter, in the fluid contents of which the little parasites were swimming. The cup-shaped body of the *Pemmatodiscus* (Fig. 233, 1) is sometimes rather flat, and shaped like a hat or cone, at other times almost curved into a semi-circle. The simple hollow of the cup, the primitive gut (*g*), has a narrow opening (*o*). The skin layer (*e*) consists of long slender cylindrical cells, which bear long vibratory hairs; it is separated by a thin structureless, gelatinous plate (*f*) from the visceral or gut layer (*i*), the primitive cells of which are much smaller and have no cilia. *Pemmatodiscus* propagates asexually, by simple longitudinal cleavage; on this account it has recently been regarded as the representative of a special order of gastræids (*Metagastria*).

Probably a near relative of the *Pemmatodiscus* is the *Kanellaria Grunthi* (Fig. 233, 2). It lives in the body-cavity of *Vermalia* (*Siphonactis*), and differs from the former in having no lashes either on the large ectodermic cells (*e*) or the small entodermic (*i*); the germinal layers are separated by a thick, cup-shaped, gelatinous mass, which has been called the "clear vesicle" (*f*). The primitive mouth is surrounded by a dark ring that bears very strong and long vibratory lashes, and effects the swimming movements.

Pemmatodiscus and *Kanellaria* may be included in the family of the *Gastremaria*. To these gastræids with open gut are closely related the *Orthostomida* (*Rhabdaria*, Fig. 233, 3-5). They live parasitically

in the body-cavity of echinoderms (*Ophiura*) and vermetals; they are distinguished by the fact that their primitive gut-cavity is not empty, but filled with entodermic cells, from which the sexual cells are developed. These gastræids are of both sexes, the male (Fig. 3) being smaller and of a somewhat different shape from the oval female (Fig. 4).

The somewhat similar *Dicymeda* (Fig. 6) are distinguished from the preceding by the fact that their primitive gut-cavity is occupied by a single large entodermic cell instead of a crowded group of sexual cells. This cell does not yield sexual products, but afterwards divides into a number of cells (spores), each of which, without being impregnated, grows into a small embryo. The *Dicymeda* live parasitically in the body-cavity, especially the renal cavities, of the cuttle-fishes. They fall in several genera, some of which are characterised by the possession of special polar cells; the body is sometimes roundish, oval, or club-shaped, at other times long and cylindrical. The genus *Conocyema* (Figs. 7-15) differs from the ordinary *Dicymeda* in having four polar pimple in the form of a cross, which may be incipient tentacles.

The classification of the *Cyemaria* is much disputed; sometimes they are held to be parasitic infusoria (like the *Ophania*), sometimes platodes or vermetals, related to the suctorial worms or rotifers, but having degenerated through parasitism. I adhere to the phylogenetically important theory that I advanced in 1876, that we have here real gastræids, primitive survivors of the common stem-group of all the Metazoa. In the struggle for life they have found shelter in the body-cavity of other animals.

The small *Coelenteria* attached to the floor of the sea that I have called the *Physemaria* (*Halophysma* and *Gastrophysma*) probably form a third order (or class) of the living gastræids. The genus *Halophysma* (Figs. 234, 235) is externally very similar to a large rhizopod (described by the same name in 1862) of the family of the *Rhabdammidae*, which was at first taken for a sponge. In order to avoid confusion with these, I afterwards gave them the name of *Prophysma*. The whole mature body of the *Prophysma* is a simple cylindrical or oval tube, with a two-layered wall. The hollow of the tube is the gastric cavity, and the upper opening of it the mouth (Fig. 235, m).




FIG. 25.—Modern gastrulae. FIG. 1. *Fammatolopus gastrulensis* (*Nemertis*) in longitudinal section. FIG. 2. *Amphipoda* (*Amphipoda*) in longitudinal section. (From *Kessler and Gower*). FIG. 3. *Amphipoda* (*Amphipoda*) in longitudinal section. FIG. 4. *Amphipoda* (*Amphipoda*) in longitudinal section. FIG. 5. *Amphipoda* (*Amphipoda*) in longitudinal section. FIG. 6. *Amphipoda* (*Amphipoda*) in longitudinal section. FIG. 7. *Amphipoda* (*Amphipoda*) in longitudinal section. FIG. 8. *Amphipoda* (*Amphipoda*) in longitudinal section. FIG. 9. *Amphipoda* (*Amphipoda*) in longitudinal section. FIG. 10. *Amphipoda* (*Amphipoda*) in longitudinal section. FIG. 11. *Amphipoda* (*Amphipoda*) in longitudinal section. FIG. 12. *Amphipoda* (*Amphipoda*) in longitudinal section. FIG. 13. *Amphipoda* (*Amphipoda*) in longitudinal section. FIG. 14. *Amphipoda* (*Amphipoda*) in longitudinal section. FIG. 15. *Amphipoda* (*Amphipoda*) in longitudinal section. a, primitive gut, b, primitive mouth, c, calcareous, d, cuticular, e, gillanous plate between a and b, supporting plate, f, denture.

The two strata of cells that form the wall of the tube are the primary germinal layers. These rudimentary zoophytes differ from the swimming gastraea chiefly in being attached at one end (the end opposite to the foot) to the floor.

In *Prophyrama* the primitive gut is a simple oral cavity, but in the closely related *Gastrophysma* it is divided into two chambers by a transverse constriction, the hind and smaller chamber above furnishes the sexual products, the anterior one being for digestion.

The simplest sponges (*Olynthus*, Fig. 238) have the same organization as the *Physemaria*. The only material difference between them is that in response to the thin two-layered body-wall is pierced by numbers of pores. When these are closed they resemble the *Physemaria*. Possibly the gastraea that we call *Physemaria* are only sponges with the pores closed. The *Ammonocoids*, or the simple tubular sand-sponges of the deep-sea (*Ammonothus*, etc.), do not differ from

— very important point when




FIG. 234.

FIG. 235.

FIGS. 234 AND 235.—*Prophyrama primordialis*, a living gastraea. FIG. 234.—The whole of the spindle-shaped animal (attached below to the floor of the sea). FIG. 235.—The same in longitudinal section. The primitive gut (*d*) opens above at the primitive mouth (*m*). Between the clustered cells (*g*) are the anal cavities (*a*). The outer layer (*e*) consists of clustered cells (*g*) are the anal cavities (*a*). The




FIG. 236.—Anal cavity of *Gastrophysma*, in floor of the sea. FIG. 237.—Anal cavity of *Gastrophysma*, in transverse section.

the pores are closed. In my *Monograph on the Sponges* (with sixty plates) I endeavored to prove analytically that all the species of this class can be traced phylogenetically to a common stem-form (*Calcolynthus*).

The lowest form of the Cnidaria is also not far removed from the gastraea. In the interesting common fresh-water polyp (*Hydra*) the whole body is simply an oval tube with a double wall; only in this case the mouth has a crown of tentacles. Before these develop the hydra resembles an acula (Figs. 236, 237). Afterwards there are slight histological differentiations in its ectoderm, though the endoderm remains

a single stratum of cells. We find the first differentiation of epithelial and



FIG. 27A.—*Gyalium*, a very rudimentary sponge. A pore just opening in front.

stinging cells, or of muscular and neural cells, in the thick ectoderm of the hydra.

In all these rudimentary living coelenterates the sexual cells of both kinds—ova and sperm cells—are formed by the same individual; it is possible that the oldest gastraea were hermaphroditic. It is clear from comparative anatomy that hermaphroditism—the combination of both kinds of sexual cells in one individual—is the earliest form of sexual differentiation; the separation of the sexes (gonochorism) was a much later phenomenon. The sexual cells originally proceeded from the edge of the primitive mouth of the gastraeid.

CHAPTER XX.

OUR WORM-LIKE ANCESTORS

THE gastraea theory has now convinced us that all the Metazoa or multicellular animals can be traced to a common stem-form, the *Gastraea*. In accordance with the biogenetic law, we find solid proof of this in the fact that the two-layered embryos of all the Metazoa can be reduced to a primitive common type, the gastrula just as the countless species of the Metazoa do actually develop in the individual from the simple embryonic form of the gastrula; so they have all descended in past time from the common stem-form of the *Gastraea*. In this fact, and the fact we have already established that the *Gastraea* has been evolved from the hollow vesicle of the one-layered *Blasozoa*, and this again from the original unicellular stem-form, we have obtained a solid basis for our study of evolution. The clear path from the stem-cell to the gastrula represents the first section of our human stem-history (Chapters VIII., IX., and XIX.).

The second section, that leads from the *Gastraea* to the *Prochordonia*, is much more difficult and obscure. By the *Prochordonia* we mean the ancient and long-extinct animals which the important

embryonic form of the chordula proves to have once existed (cf. Figs. 83-86). The nearest of living animals to this embryonic structure are the lowest Tunicates, the Cephalata (*Appendicularia*) and the larvae of the Ascidæ. As both the Tunicates and the Vertebrates develop from the same chordula, we may infer that there was a corresponding common ancestor of both stems. We may call this the *Chordæon*, and the corresponding stem-group the *Prochordonia* or *Prochordata*.

From this important stem-group of the unarticulated *Prochordonia* (or "primitive chorda-animals") the stems of the Tunicates and Vertebrates have been divergently evolved. We shall see presently how this conclusion is justified in the present condition of morphological science.

We have first to answer the difficult and much-discussed question of the development of the Chordæon from the *Gastraea*; in other words, "How and by what transformations were the characteristic animals, resembling the embryonic chordula, which we regard as the common stem-forms of all the Chordæonia, both

Tunicates and Vertebrates, evolved from the simplest two-layered Metazoa?"

The descent of the Vertebrates from the Articulata has been maintained by a number of zoologists during the last thirty years with more zeal than discernment; and, as a vast amount has been written on the subject, we must deal with it to some extent. All three classes of Articulata in succession have been awarded the honour of being considered the "real ancestors" of the Vertebrates: first, the Annelids (earth-worms, leeches, and the like), then the Crustacea (crabs, etc.), and, finally, the Tracheata (spiders, insects, etc.). The most popular of these hypotheses was the annelid theory, which derived the Vertebrates from the Worms. It was almost simultaneously (1873) formulated by Carl Sapper, of Wurzburg, and Anton Dohrn, of Naples. The latter advanced this theory originally in favour of the falling degeneration theory, with which I dealt in my work, *Aims and Methods of Modern Embryology*.

This interesting degeneration theory—much discussed at that time, but almost forgotten now—was formed in 1873 with the aim of harmonising the results of evolution and ever-advancing Darwinism with religious belief. The spirited struggle that Darwin had occasioned by the reformation of the theory of descent in 1859, and that lasted for a decade with varying fortunes in every branch of biology, was drawing to a close in 1870-1873, and soon ended in the complete victory of transformism. To most of the disputants the chief point was not the general question of evolution, but the particular one of "man's place in nature"—"the question of questions," as Huxley rightly called it. It was soon evident to every clear-headed thinker that this question could only be answered in the sense of our anthropogeny, by admitting that man had descended from a long series of Vertebrates by gradual modification and improvement.

In this way the real affinity of man and the Vertebrates came to be admitted on all hands. Comparative anatomy and ontogeny spoke too clearly for their testimony to be ignored any longer. But in order still to save man's unique position, and especially the dogma of personal immortality, a number of natural philosophers and theologians discovered an admirable way of escape in the "theory of degeneration." Granting the affinity,

they turned the whole evolutionary theory upside down, and boldly contended that "man is not the most highly developed animal, but the animals are degenerate men." It is true that man is closely related to the ape, and belongs to the vertebrate stem; but the chain of his ancestry goes upward instead of downward. In the beginning "God created man in his own image," as the prototype of the perfect vertebrate; but, in consequence of original sin, the human race sank so low that the apes branched off from it, and afterwards the lower Vertebrates. When this theory of degeneration was consistently developed, its supporters were bound to hold that the entire animal kingdom was descended from the debased children of men.

This theory was most strenuously defended by the Catholic priest and natural philosopher, Micheli, in his *Haeckelogeny: An Academic Protest against Haeckel's Anthropogeny* (1875). In still more "academic" and somewhat mystic form the theory was advanced by a natural philosopher of the older Jena school—the mathematician and physicist, Carl Smoll. But it received its chief support on the zoological side from Anton Dohrn, who maintained the anthropocentric ideas of Smoll with particular ability. The Amphioxus, which modern science now almost unanimously regards as the real Primitive Vertebrate, the ancient model of the original vertebrate structure, is, according to Dohrn, a late, degenerate descendant of the stem, the "prodigal son" of the vertebrate family. It has descended from the Cyclostoma by a profound degeneration, and these in turn from the fishes; even the Annelids and the whole of the Tunicates are merely degenerate fishes! Following out this curious theory, Dohrn came to contest the general belief that the Coelenterata and Worms are "lower animals"; he even declared that the unicellular Protozoa were degenerate Coelenterata. In his opinion "degeneration is the great principle that explains the existence of all the lower forms."

If this Micheli-Dohrn theory were true, and all animals were really degenerate descendants of an originally perfect humanity, man would assuredly be the true centre and goal of all terrestrial life; his anthropocentric position and his immortality would be saved. Unfortunately, this trustful theory is in such

flagrant contradiction to all the known facts of paleontology and embryology that it is no longer worth serious scientific consideration.

But the case is no better for the much-discussed descent of the Vertebrates from the Annelids, which Dohrn afterwards maintained with great zeal. Of late years this hypothesis, which raised so much dust and controversy, has been entirely abandoned by most competent zoologists, even those who once supported it. Its chief supporter, Dohrn, admitted in 1890 that it is "dead and buried," and made a blushing retraction at the end of his *Studies of the Early History of the Vertebrate*.

Now that the annelid-hypothesis is "dead and buried," and other attempts to derive the Vertebrates from Molluscs, Echinoderms, or Molluscs, have been equally unsuccessful, there is only one hypothesis left to answer the question of the origin of the Vertebrates—the hypothesis that I advanced thirty-six years ago and called the "chordoma-hypothesis." In view of its sound establishment and its profound significance, it may very well claim to be a *theory*, and so should be described as the chordoma or chordoma theory.

I first advanced this theory in a series of university lectures in 1867, from which the *History of Creation* was composed. In the first edition of this work (1868) I endeavoured to prove, on the strength of Kowalevsky's epoch-making discoveries, that "of all the animals known to us (the Tunicates are undoubtedly the nearest blood-relatives of the Vertebrates), they are the most closely related to the Vermalia, from which the Vertebrates have been evolved. Naturally, I do not mean that the Vertebrates have descended from the Tunicates, but that the two groups have sprung from a common root. It is clear that the real Vertebrates (primarily the Acrania) were evolved in very early times from a group of Worms, from which the degenerate Tunicates also descended in another and retrogressive direction." This common extinct stem-group are the Prochordonia; we still have a silhouette of them in the chordula-embryos of the Vertebrates and Tunicates; and they still exist independently, in very modified form, in the class of the Copelata (*Appendicaria*, Fig. 325).

The chordoma-theory received the most valuable and competent support from

Carl Gegenbaur. This able comparative morphologist defended it in 1870, in the second edition of his *Elements of Comparative Anatomy*; at the same time he drew attention to the important relations of the Tunicates to a curious worm, *Balanoglossus*: he rightly regards this as the representative of a special class of worms, which he called "gut-breathers" (*Enteropneusta*). Gegenbaur referred on many other occasions to the close blood-relationship of the Tunicates and Vertebrates, and luminously explained the reasons that justify us in framing the hypothesis of the descent of the two stems from a common ancestor, an unsegmented worm-like animal with an axial chorda between the dorsal nerve-tube and the ventral gut-tube.

The theory afterwards received a good deal of support from the research made by a number of distinguished zoologists and anatomists, especially C. Kupffer, B. Hatschek, F. Balfour, E. Van Beneden, and Julin. Since Hatschek's *Studies of the Development of the Amphioxus* gave us full information as to the embryology of this lowest vertebrate, it has become so important for our purpose that we must consider it a document of the first rank for answering the question we are dealing with.

The ontogenetic facts that we gather from this sole survivor of the Acrania are the more valuable for phylogenetic purposes, as paleontology, unfortunately, throws no light whatever on the origin of the Vertebrates. Their invertebrate ancestors were soft organisms without skeleton, and thus incapable of fossilization, as is still the case with the lowest vertebrates—the Acrania and Cyclostoma. The same applies to the greater part of the Vermalia or worm-like animals, the various classes and orders of which differ so much in structure. The isolated groups of this rich stem are living branches of a huge tree, the greater part of which has long been dead, and we have no fossil evidence as to its earlier form. Nevertheless, some of the surviving groups are very instructive, and give us clear indications of the way in which the Chordonia were developed from the Vermalia, and these from the Coelenteria.

While we seek the most important of these palaeogenetic forms among the groups of Coelenteria and Vermalia, it is understood that not a single one of them

must be regarded as an unchanged, or even little changed, copy of the extinct stem-form. One group has retained one feature, another a different feature, of the original organisation, and other organs have been further developed and characteristically modified. Hence here, more than in any other part of our genealogical tree, we have to keep before our mind the full picture of development, and separate the unessential secondary phenomena from the essential and primary. It will be useful first to point out the chief advances in organisation by which the simple Gastraea gradually became the more developed Chordata.

We find our first solid datum in the gastrula of the Amphioxus (Fig. 38). Its bilateral and tri-axial type indicates that the Gastraea—the common ancestor of all the Metazoa—divided at an early stage into two divergent groups. The uniaxial Gastraea became sessile, and gave rise to two stems, the Sponges and the Chordata (the latter all reducible to simple polyps like the hydra). But the tri-axial Gastraea assumed a certain pose or direction of the body on account of its swimming or creeping movement, and in order to sustain this it was a great advantage to share the burden equally between the two halves of the body (right and left). Thus arose the typical bilateral form, which has three axes. The same bilateral type is found in all our artificial means of locomotion—crabs, ships, etc.; it is by far the best for the movement of the body in a certain direction and steady position. Hence natural selection early developed this bilateral type in a section of the Gastraea, and thus produced the stem-forms of all the bilateral animals.

The *Gastraea bifida*, of which we may conceive the bilateral gastrula of the amphioxus to be a perigenetic reproduction, represented the two-celled organism of the earliest Metazoa in its simplest form. The vegetal ectoderm (that lined their simple gut-cavity served for nutrition; the ciliated ectoderm that formed the external skin attended to locomotion and sensation; finally, the two primitive mesodermic cells, that lay to the right and left at the ventral border of the primitive mouth, were sexual cells, and effected reproduction. In order to understand the further development of the gastrula, we must pay particular attention to: (1) the careful study of the embryonic stages of the amphioxus that lie between the

gastrula and the chordula; (2) the morphological study of the simplest Platodes (*Platodaria* and *Turbellaria*) and several groups of unarticulated Vermalia (*Gastrellia*, *Nemertea*, *Euterpneuria*).

We have to consider the Platodes first, because they are on the border between the two principal groups of the Metazoa, the Coelentera and the Coelomaria. With the former they share the lack of body-cavity, anus, and vascular system; with the latter they have in common the bilateral type, the possession of a pair of nephridia or renal canals, and the formation of a vertical brain or cerebral ganglion. It is now usual to distinguish four classes of Platodes: the two free-living classes of the primitive worms (*Platodaria*) and the coiled-worms (*Turbellaria*), and the two parasitic classes of the suctorial worms (*Trematoda*) and the tape-worms (*Cestoda*). We have only to consider the first two of these classes, the other two are parasites, and have descended from the former by adaptation to parasitic habits and consequent degeneration.

The primitive worms (*Platodaria*) are very small flat worms of simple construction, but of great morphological and phylogenetic interest. They have been hitherto, as a rule, regarded as a special order of the *Turbellaria*, and associated with the *Aphroditidae*, but they differ considerably from these and all the other Platodes (flat worms) in the absence of renal canals and a special central nervous system; the structure of their tissues is also simpler than in the other Platodes. Most of the Platodes of this group (*Aphroditomem*, *Amphistoma*, *Corrodula*, *Achisopora*, etc.) are very soft and delicate animals, swimming about in the sea by means of a ciliary coat, and very small ($\frac{1}{16}$ to $\frac{1}{8}$ inch long). Their oval body, without appendages, is sometimes spindle-shaped or cylindrical, sometimes flat and leaf-shaped. Their skin is merely a layer of ciliated ectodermic cells. Under this is a soft medullary substance, which consists of endodermic cells with vacuoles. The food passes through the mouth directly into this digestive medullary substance, in which we do not generally see any permanent gut-cavity (it may have entirely collapsed); hence these primitive Platodes have been called *Atrichs* (without gut-cavity or coelom), or, more correctly, *Cryptocela*, or *Pseudocela*. The sexual organs of these hermaphrodite

Platodaria are very simple—two pairs of strings of cells, the inner of which (the ovaries, Fig. 239 *e*) produce ova, and the outer (the spermata, *s*) sperm-cells. These gonads are not yet independent sexual glands, but sexually differentiated cell-groups in the medullary substance, or, in other words, parts of the gut-wall. Their products, the sex-cells, are conveyed out behind by two pairs of short canals;




FIG. 239.—*Aphroditomorpha* Langh (*Turbellaria*), a primitive worm of the platodaria class, of the order *Cyrtodaria* or *Arctia*. The two aspects of the genus *Aphroditomorpha*, named after Professor Arnold Lang of Zürich, was found in September, 1895, at Ajaccio in Corsica (crawping between *Mytilus*). It is one-twelfth of an inch long, one-twenty-fifth of an inch broad, and visible in colour: *a* mouth, *b* auditory vesicle, *c* ectoderm, *d* cuticular, *e* ovary, *f* spermarium, *g* female aperture, *h* male aperture.

the male opening (*m*) lies just behind the female (*f*). Most of the Platodaria have not the muscular pharynx, which is very advanced in the *Turbellaria* and *Trentoloda*. On the other hand, they have, as a rule, before or behind the mouth, a bulbous sense-organ (auditory vesicle or organ of equilibrium, *g*), and many of them have also a couple of simple optic spots. The cell-pit of the ectoderm

that lies underneath is rather thick, and represents the first rudiment of a neural ganglion (vertical brain or neuropilium).

The *Turbellaria*, with which the simpler *Platodaria* were formerly classed, differ materially from them in the more advanced structure of their organs, and especially in having a central nervous system (vertical brain) and excretory renal canals (nephridia); both originate from the ectoderm. But between the two germinal layers a mesoderm is developed, a soft mass of connective tissue, in which the organs are embedded. The *Turbellaria* are still represented by a number of different forms, in both fresh and sea-water. The oldest of these are the very rudimentary and tiny forms that are known as *Rhabdocoela* on account of the simple construction of their gut; they are, as a rule, less than a quarter of an inch long, and of a simple oval or lanceolate shape (Fig. 240). The surface is covered with ciliated epithelium, a stratum of ectodermic cells. The digestive gut is still the simple primitive gut of the gastraea (*d*), with a single aperture that is both mouth and anus (*m*). There is, however, an invagination of the ectoderm at the mouth, which has given rise to a muscular pharynx (*ph*). It is noteworthy that the mouth of the *Turbellaria* (like the primitive mouth of the Gastraea) may, in this class, change its position considerably in the middle line of the ventral surface; sometimes it lies behind (*Ophiodontomorpha*), sometimes in the middle (*Meosostomum*), sometimes in front (*Prostomum*). This displacement of the mouth from front to rear is very interesting, because it corresponds to a phylogenetic displacement of the mouth. This probably occurred in the Platodaria ancestors of most (or all) of the *Coelomaria*; in these the permanent mouth (*metastoma*) lies at the fore end (oral pole), whereas the primitive mouth (*prostoma*) lay at the hind end of the bilateral body.

In most of the *Turbellaria* there is a narrow cavity, containing a number of secondary organs, between the two primary germinal layers, the outer or animal layer of which forms the epidermis and the inner vegetal layer the visceral epithelium. The culdest of these organs are the sexual organs; they are very variously constructed in the Platodaria; in the simplest case there are merely two pairs of gonads or sexual glands—a pair of testicles (Fig. 241a)

and a pair of ovaries (*o*). They open externally, sometimes by a common aperture (*Neogonopora*), sometimes by separate ones, the female behind the male (*Digonopora*, Fig. 241). The sexual glands develop originally from the two pronephroses or primitive mesodermic cells (Fig. 83 p). As these earliest mesodermic structures extended, and became spacious sexual pouches in the later descendants of the Platodes, probably the two custom-pouches were formed from them, the first trace of the real body-cavity of the higher Metazoa (*Enterocœlia*).

The gonads are among the oldest organs, the few other organs that we find in the Platodes between the gut-wall and body-wall being later evolutionary products. One of the oldest and most important of these are the kidneys or *nephridia*, which remove unusable matter from the body (Fig. 240 *nc*). These urinary or excretory organs were originally enlarged skin-glands—a couple of canals that run the length of the body, and have a separate or common external aperture (*na*). They often have a number of branches. These special excretory organs are not found in the other Coelenterata (Gastræads, Sponges, Cnidaria) or the Cryptocœlia. They are first met in the *Turbellaria*, and have been transmitted direct from these to the *Tremæta*, and from these to the higher stems.

Finally, there is a very important new organ in the *Turbellaria*, which we do not find in the *Cryptocœlia* (Fig. 239) and their gastræad ancestors—the rudimentary nervous system. It consists of a couple of simple cerebral ganglia (Fig. 241 *g*) and fine nervous fibres that radiate from them; these are partly voluntary nerves (or motor fibres) that go to the thin muscular layer developing under the skin; and partly sensory nerves that proceed to the sense-cells of the ciliated epidermis (*f*). Many of the *Turbellaria* have also special sense-organs; a couple of ciliated smell pits (*sa*), rudimentary eyes (*se*), and, less frequently, auditory vesicles.

On these principles I assume that the oldest and simplest *Turbellaria* arose from Platodaria, and these directly from bilateral Gastræads. The chief advances were the formation of gonads and nephridia, and of the rudimentary brain. On this hypothesis, which I advanced in 1872

in the first sketch of the gastræa-theory (*Monograph on the Sponges*), there is no direct affinity between the Platodes and the Cnidaria.

Next to the ancient stem-group of the *Turbellaria* come a number of more recent chordoma ancestors, which we class with the *Vermæta* or *Helminthes*, the unarticulated worms. These true




FIG. 240.—A simple *Turbellaria* (*Planorbis*), as shown, *ad* guttural epithelium, *gc* guttural chamber, *g* cerebral ganglion, *nc* nephridial canal, *na* renal aperture, *se* eye, *sa* olfactory pit. (Diagram.)

FIG. 241.—The same, showing the other organs, *g* brain, *se* eye, *sa* olfactory pit, *f* nervous, *nc* nephridial, *na* male aperture, *o* female aperture, *o* ovary, *f* ciliated epidermis. (Diagram.)

worms (*Tremæta*, lately also called *Scelerozoa*) are the difficulty or the lumber-room of the zoological classifier, because the various classes have very complicated relations to the lower Platodes on the one hand and the more advanced animals on the other. But if we exclude the Platodes and the Annelids from this stem, we find a fairly satisfactory unity of organisation

in the remaining classes. Among these worms we find some important forms that show considerable advances in organization from the platode to the chordonia stage. Three of these phenomena are particularly instructive: (1) The formation of a true (secondary) body-cavity (coeloma), (2) the formation of a second aperture of the gut, the anus, and (3) the formation of a vascular system. The great majority of the Vermalia have these

of an anus at the posterior end (Fig. 243 a). Further, the cilia that cover the whole surface of the Turbellaria are confined in the Gastrotricha to two ciliated bands (*f*) on the ventral surface of the oval body, the dorsal surface having bristles. Otherwise the organization of the two classes is the same. In both the gut consists of a muscular gullet (*c*) and glandular primitive gut (*d*). Over the gullet is a double brain (acroganglion, *g*;




FIG. 242.




FIG. 243.

FIGS. 242 and 243.—Chordonotus, a rudimentary Vermalian, of the group of Gastrotricha. *a*, mouth; *c*, gullet; *d*, gut; *e*, anus; *g*, brain; *h*, nephrite; *i*, nephrite; *j*, nephrite; *k*, nephrite; *l*, nephrite; *m*, nephrite; *n*, nephrite; *o*, nephrite; *p*, nephrite; *q*, nephrite; *r*, nephrite; *s*, nephrite.

three features, and they are all wanting in the Platodes; in the rest of the worms at least one or two of them are developed.

Next and very close to the Platodes we have the Ichthyodina (*Gastrotricha*), little marine and fresh-water worms, about $\frac{1}{16}$ to $\frac{1}{8}$ inch long. Zoologists differ as to their position in classification. In my opinion, they approach very close to the Rhabdocoela (Figs. 242, 241), and differ from them chiefly in the possession

At the side of the gut are two serpentine prorenal canals (water-vessels or pro-nephridia, *ns*), which open on the ventral side (*am*). Behind are a pair of simple sexual glands or gonads (Fig. 243 *e*).

While the Ichthyodina are thus closely related to the Platodes, we have to go farther away for the two classes of Vermalia which we unite in the group of the "snout-worms" (*Frontonia*). These are the *Nemertina* and the *Enicopneusta*

Both classes have a complete ciliary coat on the epidermis, a heritage from the Turbellaria and the Gavreans; also, both have two openings of the gut, the mouth and anus, like the Gastrotricha. But we find also an important organ that is wanting in the preceding forms—the vascular system. In their more advanced mesoderm we find a few contractile longitudinal canals which force the blood through the body by their contractions; these are the first blood-vessels.




FIG. 144.

FIG. 144.—A simple Metartemate. m mouth of gut. a anus, b brain, n nerves, c ciliary coat, or sensory pits (head-shield), e eyes, f dorsal vessel, l lateral vessels. (Diagram.)

FIG. 145.—A young Euberozoan (Belonoglossa). (From Alexander, *Argona*.) p acorn-shaped soma, h hook, g gill-chute and gill-barrel of the fore-gut, m long runs on each side, d digestive body-gut, filling the greater part of the body-cavity, v ventral vein or ventral vessel, lying between the parallel folds of the skin, n parapodia.




FIG. 145.

The Nemertina were formerly classed with the much less advanced Turbellaria. But they differ essentially from them in having an anus and blood-vessels, and several other marks of higher organization. They have generally long and narrow bodies, like a more or less flattened cord; there are, besides several small species, giant-forms with a width of $\frac{1}{2}$ to 1 inch and a length of several yards (even ten to fifteen). Most of them live in the sea, but some in fresh water and moist earth. In their internal structure they approach the Turbellaria on the one hand and the higher Vermalia (especially the Enteroptneusta) on the other. They have a good deal of interest as the lowest and oldest of all animals with blood. In them we find blood-vessels for the first time, distributing real blood through the body.




FIG. 246.—Transverse section of the branchial gut. *A* of *Nemertina*, *B* of *Chordone*. *r* branchial gut, *s* pharyngeal groove, *v* ventral fold between the two. Diagrammatic illustration from Gegenbaur, as above the relation of the dorsal branchial cavity (*r*) to the pharyngeal or hypobranchial groove (*s*).

The blood is red, and the red colouring-matter is hæmoglobin, connected with elliptic discoid blood-cells, as in the Vertebrates. Most of them have two or three parallel blood-canals, which run the whole length of the body, and are connected in front and behind by loops, and often by a number of ring-shaped pieces. The chief of these primitive blood-vessels is the one that lies above the gut in the middle line of the back (Fig. 244 *r*); it may be compared to either the dorsal vessel of the Articulata or the aorta of the Vertebrates. To the right and left are the two serpentine lateral vessels (Fig. 244 *l*).

After the Nemertina, I take (as distant relatives) the Enteroptneusta; they may be classed together with them as *Promela* or *Rhynchocela* (snout-worms). There is now only one genus of this class, with several species (*Halargaster*); but it

is very remarkable, and may be regarded as the last survivor of an ancient and long-extinct class of Vermalia. They are related, on the one hand, to the Nemertina and their immediate ancestors, the Pluteles, and to the lowest and oldest forms of the Chordonia on the other.

The Enteroptneusta (Fig. 245) live in the sea sand, and are long worms of very simple shape, like the Nemertina. From the latter they have inherited: (1) The bilateral type, with incomplete segmentation; (2) the ciliary coat of the soft epidermis; (3) the double rows of gastric pouches, alternating with a single or double row of grooves; (4) separation of the sexes (the Plutele ancestors were hermaphrodite); (5) the ventral mouth, underneath a protruding snout; (6) the anus terminating the simple gut-tube; and (7) several parallel blood-canals, running the length of the body, a dorsal and a ventral principal stem.

On the other hand, the Enteroptneusta differ from their Nemertine ancestors in several features, some of which are important, that we may attribute to adaptation. The chief of these is the branchial gut (Fig. 245 *B*). The anterior portion of the gut is converted into a respiratory organ, and pierced by two rows of gill-clefts; between these there is a branchial (gill) skeleton, formed of rods and plates of chitine. The water that enters

at the mouth makes its exit by these clefts. They lie in the dorsal half of the fore-gut, and this is completely separated from the ventral half by two longitudinal folds (Fig. 245 *A*). This ventral half, the glandular walls of which are clothed with ciliary epithelium and secrete mucus, corresponds to the pharyngeal or hypobranchial groove of the Chordonia (*A*), the important organ from which the later thyroid gland is developed in the Craniota (cf. p. 184). The agreement in the structure of the branchial gut of the Enteroptneusta, Tunicata, and Vertebrates was first recognised by Gegenbaur (1878); it is the more significant as at first we find only a couple of gill-clefts in the young animals of all three groups; the number gradually increases. We can infer from this the common descent of the three groups with all the more

confidence when we find the *Balanoglossus* approaching the Chordonia in other respects. Thus, for instance, the chief part of the central nervous system is a long dorsal neural string that runs above the gut and corresponds to the medullary tube of the Chordonia. Bateson believes he has detected a rudimentary chorda between the two.

Of all extant invertebrate animals the Enteropneusts come nearest to the Chordonia in virtue of these peculiar characters, hence we may regard them as the survivors of the ancient gut-breathing Vernalia from which the Chordonia also have descended. Again, of all the chorda-animals the Cephalata (Fig. 225) and the tailed larvae of the ascidia approach nearest to the young *Balanoglossus*. Both are, on the other hand, very closely related to the *Ampelousa*, the Primitive Vertebrate of which we have considered the importance (Chapters XVI. and XVII.). As we saw there, the unarticulated Tunicates and the articulated Vertebrates must be regarded as two independent stems, that have developed in divergent directions. But the common root of the two stems, the extinct group of the Prochordonia, must be sought in the vernalia stem; and of all the living Vernalia those we have considered give us the safest clue to their origin. It is true that the actual representatives of the important groups of the Cephalata, *Balanoglossus*, Nemertina, Ichthyodonta, etc., have more or less departed from the primitive model owing to adaptation to special environment. But we may just as confidently affirm that the main features of

their organization have been preserved by heredity.

We must grant, however, that in the whole stem-history of the Vertebrates the long stretch from the Gastraea and Planodes up to the oldest Chordonia remains by far the most obscure section. We might frame another hypothesis to raise the difficulty—namely, that there was a long series of very different and totally extinct forms between the Gastraea and the Chordonia. Even in this modified chordata-theory the six fundamental organs of the chordula would retain their great value. The medullary tube would be originally a chemical sensory organ, a dorsal olfactory tube, taking in respiratory-water and food by the neuropore in front and conveying them by the neurenteric canal into the primitive gut. This olfactory tube would afterwards become the nervous centre, while the expanding gonads (lying to right and left of the primitive mouth) would form the columns. The chorda may have been originally a digestive glandular groove in the dorsal middle line of the primitive gut. The two secondary gut-openings, mouth and anus, may have arisen in various ways by change of functions. In any case, we should ascribe the same high value to the chordula as we did before to the gastrula.

In order to explain more fully the chief stages in the advance of our race, I add the hypothetical sketch of man's ancestry that I published in my *Last Leaf* [a translation by Dr. Gadow of the paper read at the International Zoological Congress at Cambridge in 1898].—

**A.—Man's Genealogical Tree, First Half:
EARLIER SERIES OF ANCESTORS, WITHOUT
FOSSIL EVIDENCE.**

Chief Name.	Assumed Stem-group.	Living Relations of Ancestors.
Stages 1-5 Protist ancestors. Unicellular organisms. 1-5 Phytopl. Im. 2-4 Protozoa	1. Eucarya. Without nucleus. 2. Algae. Unicellular algae. 3. Lobos. Unicellular (multicellular) rhizoids. 4. Jellies. Unicellular. 5. Multicellular. Multicellular hollow spheres	1. Chlamydomonas. (Chlamydomonas?) 2. Volvox. 3. Eukaryotes. 4. Eukaryotes. 5. Eukaryotes. 6. Eukaryotes. 7. Eukaryotes. 8. Eukaryotes. 9. Eukaryotes. 10. Eukaryotes. 11. Eukaryotes. 12. Eukaryotes. 13. Eukaryotes. 14. Eukaryotes. 15. Eukaryotes. 16. Eukaryotes. 17. Eukaryotes. 18. Eukaryotes. 19. Eukaryotes. 20. Eukaryotes. 21. Eukaryotes. 22. Eukaryotes. 23. Eukaryotes. 24. Eukaryotes. 25. Eukaryotes. 26. Eukaryotes. 27. Eukaryotes. 28. Eukaryotes. 29. Eukaryotes. 30. Eukaryotes. 31. Eukaryotes. 32. Eukaryotes. 33. Eukaryotes. 34. Eukaryotes. 35. Eukaryotes. 36. Eukaryotes. 37. Eukaryotes. 38. Eukaryotes. 39. Eukaryotes. 40. Eukaryotes. 41. Eukaryotes. 42. Eukaryotes. 43. Eukaryotes. 44. Eukaryotes. 45. Eukaryotes. 46. Eukaryotes. 47. Eukaryotes. 48. Eukaryotes. 49. Eukaryotes. 50. Eukaryotes. 51. Eukaryotes. 52. Eukaryotes. 53. Eukaryotes. 54. Eukaryotes. 55. Eukaryotes. 56. Eukaryotes. 57. Eukaryotes. 58. Eukaryotes. 59. Eukaryotes. 60. Eukaryotes. 61. Eukaryotes. 62. Eukaryotes. 63. Eukaryotes. 64. Eukaryotes. 65. Eukaryotes. 66. Eukaryotes. 67. Eukaryotes. 68. Eukaryotes. 69. Eukaryotes. 70. Eukaryotes. 71. Eukaryotes. 72. Eukaryotes. 73. Eukaryotes. 74. Eukaryotes. 75. Eukaryotes. 76. Eukaryotes. 77. Eukaryotes. 78. Eukaryotes. 79. Eukaryotes. 80. Eukaryotes. 81. Eukaryotes. 82. Eukaryotes. 83. Eukaryotes. 84. Eukaryotes. 85. Eukaryotes. 86. Eukaryotes. 87. Eukaryotes. 88. Eukaryotes. 89. Eukaryotes. 90. Eukaryotes. 91. Eukaryotes. 92. Eukaryotes. 93. Eukaryotes. 94. Eukaryotes. 95. Eukaryotes. 96. Eukaryotes. 97. Eukaryotes. 98. Eukaryotes. 99. Eukaryotes. 100. Eukaryotes.

**B.—Man's Genealogical Tree, Second Half:
LATER ANCESTORS, WITH FOSSIL EVIDENCE.**

Geological Periods.	Assumed Stem-group.	Living Relations of Ancestors.
Silurian.	1. Silurian. Protist Im.	1. Silurian. Protist Im.
Devonian.	2. Devonian. Protist Im.	2. Devonian. Protist Im.
Carboniferous.	3. Carboniferous. Protist Im.	3. Carboniferous. Protist Im.
Permian.	4. Permian. Protist Im.	4. Permian. Protist Im.
Triassic.	5. Triassic. Protist Im.	5. Triassic. Protist Im.
Jurassic.	6. Jurassic. Protist Im.	6. Jurassic. Protist Im.
Cretaceous.	7. Cretaceous. Protist Im.	7. Cretaceous. Protist Im.
Older Eocene.	8. Older Eocene. Protist Im.	8. Older Eocene. Protist Im.
Mid-Eocene.	9. Mid-Eocene. Protist Im.	9. Mid-Eocene. Protist Im.
Oligocene.	10. Oligocene. Protist Im.	10. Oligocene. Protist Im.
Older Miocene.	11. Older Miocene. Protist Im.	11. Older Miocene. Protist Im.
New Miocene.	12. New Miocene. Protist Im.	12. New Miocene. Protist Im.
Pliocene.	13. Pliocene. Protist Im.	13. Pliocene. Protist Im.
Quaternary.	14. Quaternary. Protist Im.	14. Quaternary. Protist Im.

CHAPTER XXI.

OUR FISH-LIKE ANCESTORS

Our task of detecting the extinct ancestors of our race among the vast numbers of animals known to us encounters very different difficulties in the various sections of man's stem-history. These were very great in the series of our invertebrate ancestors; they are much slighter in the subsequent series of our vertebrate ancestors. Within the vertebrate stem there is, as we have already seen, so complete an agreement in structure and embryology that it is impossible to doubt their phylogenetic unity. In this case the evidence is much clearer and more abundant.

The characteristics that distinguish the Vertebrates as a whole from the Invertebrates have already been discussed in our description of the hypothetical Primitive Vertebrate (Chapter XI. Figs. 98-102). The chief of these are (1) The evolution of the primitive brain into a dorsal medullary tube, (2) the formation of the chorda between the medullary tube and the gut, (3) the division of the gut into branchial (gill) and hepatic (liver) gut, and (4) the internal articulation or metameres. The first three features are shared by the Vertebrates with the Ascidiata and the Prochordonia, the fourth is peculiar to them. Thus the chief advantage in organisation by which the earliest Vertebrates took precedence of the unsegmented Chordonia consisted in the development of internal segmentation.

This whole vertebrate stem divides first into the two chief sections of Acrania and Craniota. The Amphioxus is the only surviving representative of the older and lower section, the Acrania ("skull-less"). All the other vertebrates belong to the second division, the Craniota ("skull-animals"). [The Craniota descend directly from the Acrania, and these from the primitive Chordonia. The exhaustive study that we made of the comparative anatomy and ontogeny of the Ascidia and the Amphioxus has proved these relations for us. (See Chapters XVI. and XVII.)

The Amphioxus, the lowest Vertebrate, and the Ascidia, the nearest related Invertebrate, descend from a common extinct stem-form, the Chordonia; and this must have had, substantially, the organisation of the chordula.

However, the Amphioxus is important not merely because it fills the deep gulf between the Invertebrates and Vertebrates, but also because it shows us to-day the typical vertebrate in all its simplicity. We owe to it the most important data that we proceed on in reconstructing the gradual historical development of the whole stem. All the Craniota descend from a common stem-form, and this was substantially identical in structure with the Amphioxus. This stem-form, the Primitive Vertebrate (*Protophyllus*, Figs. 98-102), had the characteristics of the vertebrate as such, but not the important features that distinguish the Craniota from the Acrania. Though the Amphioxus has many peculiarities of structure and has much degenerated, and though it cannot be regarded as an unchanged descendant of the Primitive Vertebrate, it must have inherited from it the specific characters here enumerated above. We may not say that "Amphioxus is the ancestor of the Vertebrates"; but we can say: "Amphioxus is the nearest relation to the ancestor of all the animals we know." Both belong to the same small family, or lowest class of the Vertebrates, that we call the Acrania. In our genealogical tree this group forms the twelfth stage, or the first stage among the vertebrate ancestors (p. 228). From this group of Acrania both the Amphioxus and the Craniota were evolved.

The vast division of the Craniota embraces all the Vertebrates known to us, with the exception of the Amphioxus. All of them have a head clearly differentiated from the trunk, and a skull enclosing a brain. The head has also three pairs of higher sense-organs (nose, eyes, and ears). The brain is very rudimentary at first, a mere bulbous enlargement of the



FIG. 247.

fore end of the medullary tube. But it is soon divided by a number of transverse constrictions into, first three, then five successive cerebral vesicles. In this formation of the head, skull, and brain, with further development of the higher sense-organs, we have the advance that the Craniota made beyond their skull-less ancestors. Other organs also attained a higher development; they acquired a compact centralised heart with valves and a more advanced liver and kidneys, and made progress in other important respects.

We may divide the Craniota generally into *Cyclostoma* ("round-mouthed") and *Gnathostoma* ("jaw-mouthed"). There are only a few groups of the former in existence now, but they are very interesting, because in their whole structure they stand midway between the Acrania and the Gnathostoma. They are much more advanced than the Acrania, much less so than the fishes, and thus form a very welcome connecting-link between the two groups. We may therefore consider them a special intermediate group, the fourteenth and fifteenth stages in the series of our ancestors.

The few surviving species of the *Cyclostoma* are divided into two orders—the *Myxinoidei* and the *Petromyzontes*. The former, the hag-

fishes, have a long, cylindrical, worm-like body. They were classed by Linné with the worms, and by later zoologists with the fishes, or the amphibia, or the mollusca. They live in the sea, usually as parasites of fishes, into the skin of which they bore with their round suctional mouths and their tongues, armed with horny teeth. They are sometimes found alive in the body-cavity of fishes (such as the loach or sturgeon); in these cases they have passed through the skin into the interior. The second order consists of the *Petromyzontes* or lampreys; the small river lamprey (*Petromyzon fluviatilis*) and the large marine lamprey (*Petromyzon marinus*, Fig. 247). They also have a round suctional mouth, with horny teeth inside it; by means of this they attach themselves by sucking to fishes, stones, and other objects (hence the name *Petromyzus* = stone-sucker). It seems that this habit was very widespread among the earlier Vertebrates; the larvae of many of the Ganoids and frogs have suctional disks near the mouth.

The class that is formed of the *Myxinoidei* and *Petromyzontes* is called the *Cyclostoma* (round-mouthed), because their mouth has a circular or semi-circular aperture. The jaws (upper and lower) that we find in all the higher Vertebrates are completely wanting in the *Cyclostoma*, as in the *Amphioxus*. Hence the other Vertebrates are collectively opposed to them as *Gnathostoma* (jaw-mouthed). The *Cyclostoma* might also be called *Monorhina* (single-nosed), because they have only a single nasal passage, while all the *Gnathostoma* have two nostrils (*Amphirhina* = double-nosed). But apart from these peculiarities the *Cyclostoma* differ more widely from the fishes in other special features of their structure than the fishes do from man. Hence they are obviously the last survivors of a very ancient class of Vertebrates, that was far from attaining the advanced organization of the true fish. To mention only the chief points, the *Cyclostoma* show no trace of pairs of limbs. Their mucous skin is quite naked and smooth and devoid of scales. There is no bony skeleton. A very rudimentary skull is developed at the foremost end of their chorda. At this point a soft membranous (partly turning into cartilage), small skull-capule is formed, and encloses the brain.

FIG. 247.—The large marine lamprey (*Petromyzon marinus*), much reduced. Behind the eye there is a row of seven suckers used to attach to the fish, or to stones, etc.

or double-boned Vertebrates (*Gnathostomes* or *Amphibians*). We have to consider the fishes carefully as the class which, on the evidence of paleontology,




FIG. 100.—Embryo of a shark (*Squalus fetus*), seen from the ventral side. *a*, head-form (in front five pairs of gill-slits); *b*, belly-form; *c*, eye; *d*, tail-form; *e*, external gill-rakers; *f*, gill-arch (removed for most part); *g*, eye; *h*, nose; *i*, mouth-form.

comparative anatomy, and ontogeny, may be regarded with absolute certainty as the

stem-class of all the higher Vertebrates or Gnathostomes. Naturally, none of the actual fishes can be considered the direct ancestor of the higher Vertebrates. But it is certain that all the Vertebrates or Gnathostomes, from the fishes to man, descend from a common, extinct, fish-like ancestor. If we had this ancient stem-form before us, we would undoubtedly class it as a true fish. Fortunately, the comparative anatomy and classification of the fishes are now so far advanced that we can get a very clear idea of these interesting and instructive features.

In order to understand properly the genealogical tree of our race within the vertebrate stem, it is important to bear in mind the characteristics that separate the whole of the Gnathostomes from the Cyclostomes and Crustacea. In these respects the fishes agree entirely with all the other Gnathostomes up to man, and it is on this that we base our claim of relationship to the fishes. The following characteristics of the Gnathostomes are anatomic features of this kind: (1) The internal gill-arch apparatus with the jaw-arches; (2) the pair of nostrils; (3) the floating bladder or lungs; and (4) the two pairs of limbs.

The peculiar formation of the framework of the branchial (gill) arches and the connected maxillary (jaw) apparatus is of importance to the whole group of the Gnathostomes. It is inherited in a rudimentary form by all of them, from the earliest fishes to man. It is true that the primitive transformation (which we find even in the Ascidia) of the fore gut into the branchial gut can be traced in all the Vertebrates to the same simple type; in this respect the gill-clefts, which pierce the walls of the branchial gut in all the Vertebrates and in the Ascidia, are very characteristic. But the *external*, superficial branchial skeleton that supports the gill-grate in the Cyclostoma is replaced in the Gnathostomes by an *internal* branchial skeleton. It consists of a number of successive cartilaginous arches, which lie in the wall of the gullet between the gill-clefts, and run round the gullet from both sides. The foremost pair of gill-arches become the maxillary arches, from which we get our upper and lower jaws.

The olfactory organs are at first found in the same form in all the Gnathostomes, as a pair of depressions in the fore part of the skin of the head, above the mouth; hence, they are also called the *Amphibians*.

or *Tetacanth* was developed, to which the great majority of living fishes belong (especially nearly all our river fishes). Comparative anatomy and ontogeny show clearly that the Ganoids descended from the Selachii, and the Teleostei from the Ganoids. On the other hand, a collateral



FIG. 252.—Fossil angui-shark (*Squalodon albirostris*), from the upper Jurassic of Bavaria. (From Zittel.) The cartilaginous skull is clearly seen at the broad head, and the vertebrae behind. The wide bristled and the narrow jelly-fin have a number of canals; between these and the vertebrae are a number of ribs.

line, or rather the advancing chief line of the vertebrate stem, was developed from the earlier Ganoids, and this leads us through the group of the Dipneusts to the important division of the Amphibia.

The earliest fossil remains of Vertebrates that we know were found in the

Upper Silurian (p. 205), and belong to two groups—the Selachii and the Ganoids. The most primitive of all known representatives of the earliest fishes are probably the remarkable *Platyrhynchids*, the genera *Platyrhynchus*, *Neuracanthus*, *Orthacanthus*, etc. (Fig. 248). These ancient cartilaginous fishes agree in most points of structure with the real sharks (Figs. 249, 250); but in other respects they seem to be so much simpler in organization that many paleontologists separate them altogether, and regard them as *Protoselachii*; they are probably closely related to the extinct ancestors of the Gnathostomes. We find well-preserved remains of them in the Permian period. Well-preserved impressions of other sharks are found in the Jurassic schist, such as of the angui-fish (*Squalius*, Fig. 251). Among the extinct earlier sharks of the Tertiary period there were some twice as large as the biggest living fishes; *Carcharodon* was more than 100 feet long. The sole surviving species of this genus (*C. Romuleus*) is eleven yards long, and has teeth two inches long, but among the fossil species we find teeth six inches long (Fig. 252).

From the primitive fishes or Selachii, the earliest Gnathostomes, was developed the legion of the Ganoids. There are very few genera now of this interesting and varied group—the ancient sturgeons (*Acipenser*), the eggs of which are eaten as caviare, and the sturged pikes (*Polypterus*, Fig. 253), in African rivers, and bowpikes (*Lepidosteus*) in the rivers of North America. On the other hand, we have a great variety of specimens of this group in the fossil state, from the Upper Silurian onward. Some of these fossil Ganoids approach closely to the Selachii; others are nearer to the Dipneusts; others again represent a transition to the Teleostei. For our genealogical purposes the most interesting are the intermediate forms between the Selachii and the Dipneusts. Huxley, to whom we owe particularly important works on the fossil Ganoids, classed them in the order of the *Crossopterygii*. Many genera and species of this order are found in the Devonian and Carboniferous strata (Fig. 253); a single, greatly modified survivor of the group is still found in the large rivers of Africa (*Polypterus*, Fig. 253, and the closely related *Colemanichthys*). In many impressions of the *Crossopterygii* the floating bladder seems to be ossified,

and therefore well in the *Urosteus* (behind the head).

Part of these *Crossopterygii* very closely in their chief anatomical characters to the *Dipneusts*, and thus represent phylogenetically the transition from the Devonian Ganoids to the earliest air-breathing vertebrates. This important advance was made in the Devonian period. The numerous fossils that we see in the first two geological

the Laurentian and Cambrian periods, belong exclusively to aquatic plants and animals. From this paleontological fact, in conjunction with important geological and biological indications, we may infer with some confidence that there were no terrestrial animals at that time. During the whole of the vast archæozoic period—many millions of years—the living population of our planet consisted almost exclusively of aquatic organisms; this is a very remarkable fact, when we remember that this period embraces the larger half of the whole history of life. The lower animal-stems are wholly (or with very few exceptions) aquatic. But the higher stems also remained in the water during the primordial epoch. It was only towards its close that some of them came to live on land. We find isolated fossil remains of terrestrial animals first in the Upper Silurian, and in larger numbers in the Devonian strata, which were deposited at the beginning of the second chief section of geology (the paleozoic age). The number increases considerably in the Carboniferous and Permian deposits. We find many species both of the articulate and the vertebrate

in land and breathed the atmosphere; their aquatic ancestors of the Silurian period only breathed water. This important change in respiration is the chief modification that the animal organism underwent in passing from the water to the solid land. The first modification was the formation of lungs for breathing air; up to that time the gills alone had served for respiration. But there was at the same time a great change in the circulation and its organs; these are always very closely correlated to the respiratory organs. Moreover, the limbs and other organs were also more

or less modified, either in consequence of remote correlation to the preceding or owing to new adaptations.

In the vertebrate stem it was unquestionably a branch of the fishes—in fact, of the Ganoids—that made the first fortunate experiment during the Devonian period of adapting themselves to terrestrial life and breathing the atmosphere. This

modification of the heart and the lungs. The true fishes have merely a pair of blind olfactory pits on the surface of the head; but a connection of these with the cavity of the mouth was now formed. A canal made its appearance on each side, and led directly from the nasal depression




FIG. 39.—Tooth of a gigantic shark (*Carcharodon megalodon*), from the Pliocene at Wahn. (Half natural size. From Zittel.)

into the mouth-cavity, atmospheric air to the

lungs. Further, in all true fishes the heart has only two sections—an atrium that receives the venous blood from the veins, and a ventricle that propels it through a conical artery to the gills; the atrium was now divided into two halves, or right and left auricles, by an incomplete partition. The right auricle alone now received the venous blood from the body, while the left auricle received the venous blood that flowed from the lungs and gills to the heart. Thus the double circulation of the higher vertebrates was evolved from the simple

circulation of the true fishes, and, in accordance with the laws of correlation, this advance led to others in the structure of other organs.

The vertebrate class, that thus adapted

they retained the earlier gill-respiration along with the new pulmonary (lung) respiration, like the lowest actinopteria. This class was represented during the paleozoic age (or the Devonian, Car-




FIG. 233.




FIG. 234.



FIG. 235.

FIG. 233.—A Devonian Crossopterygian (*Haptophycin scintillans*), 6 (From Huxley.)

FIG. 234.—A Jurensis Crossopterygian (*Urdina pinnulata*), from the *Urdina* (Zittel.) Pectoral plates, & fibrous structure.

FIG. 235.—A living Crossopterygian, from the Upper Nile (*Polypterus*).

itself to breathing the atmosphere, and was developed from a branch Gensids, takes the name of the *Dipneusti* or *Dipna* ("double-breathers"), because

boniferous, and Permian number of different genera only three genera of the to-day: *Protipterus amecion*

of tropical Africa (the White Nile, the Niger, Quellinan, etc.), *Lepidosteus paradoxus* in tropical South America (in the tributaries of the Amazon), and *Ceratodus Forsteri* in the rivers of East Australia. This wide distribution of the three isolated survivors proves that they represent a group that was formerly very large. In their whole structure they

though most now associate them with the fishes. As a matter of fact, the characters of the two classes are so far united in the Dipneusts that the answer to the question depends entirely on the definition we give of "fish" and "amphibian." In habits they are true amphibians. During the tropical winter, in the rainy season, they swim in the water like the fishes, and



FIG. 256.



FIG. 257.

FIG. 256.—Fossil Dipneust (*Dipterus T'ienmensis*) A. from the old red sandstone (Devon). (From Pander.)

FIG. 257.—The Australian Dipneust (*Ceratodus Forsteri*). A. view from the right, B. lower side of the snout, C. lower jaw (Mandibula), D. quadruple bone, E. first paracanthium, F. first pectoralium, G. lower side of the head, H. mandibula, I. branched fin ray, J. first fin. B. larger than life of the fossil *Ceratodus Forsteri* (from the Transvaal).

form a transition from the fishes to the amphibians. The transitional formation between the two classes is so pronounced in the whole organisation of these remarkable animals that zoologists had a lively controversy over the question whether they were really fishes or amphibians. Several distinguished zoologists classed them with the amphibians,

breathe water by gills. During the dry season they bury themselves in the dry mud, and breathe the atmosphere through lungs, like the amphibians and the higher vertebrates. In this double respiration they resemble the lower amphibians, and have the same characteristic formation of the heart; in this they are much superior to the fishes. But in most other features

they approach nearer to the fishes, and are inferior to the amphibia. Externally they are entirely fish-like.

In the Dipneusts the head is not marked off from the trunk. The skin is covered with large scales. The skeleton is soft, cartilaginous, and at a low stage of development, as in the lowest Selachii and the earliest Ganoids. The chorda is completely retained, and surrounded by an unsegmented sheath. The two pairs of limbs are very simple fins of a primitive

skeleton; the cartilaginous skeleton of its two pairs of fins, for instance, has still the original form of a bi-serial or feathered leaf, and was on that account described by Gegenbaur as a "primitive fin-skeleton." On the other hand, the skeleton of the pairs of fins is greatly reduced in the African dipneust (*Protopterus*) and the American (*Lepidosteus*). Further, the lungs are double in these modern dipneusts, as in all the other air-breathing vertebrates; they



FIG. 148.



FIG. 149.

FIG. 148.—Young *Ceratodus*, shortly after issuing from the egg, magnified ten times. *E. G. Cooper, J. Hyatt.* (From Richard Semon.)

FIG. 149.—Young *Ceratodus*, six weeks after issuing from the egg. *a* spiral fold of gut. *b* rudimentary belly-fin. (From Richard Semon.)

type, like those of the lowest Selachii. The formation of the brain, the gut, and the nasal organs is also the same as in the Selachii. Thus the Dipneusts have preserved by heredity many of the less advanced features of our primitive fish-like ancestors, and at the same time have made a great step forward in adaptation to air-breathing by means of lungs and the correlative improvement of the heart.

Ceratodus is particularly interesting on account of the primitive build of its

have on that account been called "double-lunged" (*Dipneustes*) in contrast to the *Ceratodus*; the latter has only a single lung (*Monopneustes*). At the same time the gills also are developed as water-breathing organs in all these lung-fishes. *Protopterus* has external as well as internal gills.

The paleozoic Dipneusts that are in the direct line of our ancestry, and form the connecting-bridge between the Ganoids and the Amphibia, differ in many respects

from their living descendants, but agree with them in the above essential features. This is confirmed by a number of interesting facts that have lately come to our knowledge in connection with the

embryonic development of the *Ceratodus* and *Leptodus*; they give us important information as to the stem-history of the lower Vertebrates, and therefore of our early ancestors of the paleozoic age.

CHAPTER XXII.

OUR FIVE-TOED ANCESTORS

WITH the phylogenetic study of the four higher classes of Vertebrates, which must now engage our attention, we reach much firmer ground and more light in the construction of our genealogy than we have, perhaps, enjoyed up to the present. In the first place, we owe a number of very valuable data to the very interesting class of Vertebrates that come next to the Dipneusta and have been developed from them—the Amphibia. To this group belong the salamander, the frog, and the toad. In earlier days all the reptiles were, on the example of Linné, classed with the Amphibia (lizards, serpents, crocodiles, and tortoises). But the reptiles are much more advanced than the Amphibia, and are nearer to the birds in the chief points of their structure. The true Amphibia are nearer to the Dipneusta and the fishes; they are also much older than the reptiles. There were plenty of highly-developed (and sometimes large) Amphibia during the Carboniferous period; but the earliest reptiles are only found in the Permian period. It is probable that the Amphibia were evolved even earlier—during the Devonian period—from the Dipneusta. The extinct Amphibia of which we have fossil remains from that remote period (very numerous especially in the Triassic strata) were distinguished for a graceful scaly coat or a powerful bony armour on the skin (like the crocodile), whereas the living amphibians have usually a smooth and slippery skin.

The earliest of these armoured Amphibia (*Phacelomphalia*) form the order of *Stegomphala* ("roof-headed") (Fig. 266). It is among these, and not among the actual Amphibia, that we must look for the forms that are directly related to the genealogy of our race, and are the ancestors of the three higher

classes of Vertebrates. But even the existing Amphibia have such important relations to us in their anatomic structure, and especially their embryonic development, that we may say: Between the Dipneusta and the Amphibia there was a series of extinct intermediate forms which we should certainly class with the Amphibia if we had them before us. In their whole organisation even the actual Amphibia seem to be an instructive transitional group. In the important respects of respiration and circulation they approach very closely to the Dipneusta, though in other respects they are far superior to them.

This is particularly true of the development of their limbs or extremities. In them we find them for the first time as five-toed feet. The thorough investigations of Gegenbaur have shown that the fish's fin, of which very erroneous opinions were formerly held, are many-toed feet. The various cartilaginous or bony radii that are found in large numbers in each fin correspond to the fingers or toes of the higher Vertebrates. The axial joints of each fin-radius correspond to the various parts of the toe. Even in the Dipneusta the fin is of the same construction as in the fishes; it was afterwards gradually evolved into the five-toed form, which we first encounter in the Amphibia. This reduction of the number of the toes to six, and then to five, probably took place in the second half of the Devonian period—at the latest, in the subsequent Carboniferous period—in those Dipneusta which we regard as the ancestors of the Amphibia. We have several fossil remains of five-toed Amphibia from this period. There are numbers of fossil impressions of them in the Triassic of Thuringia (*Chirotherium*).

The fact that the toes number five is of great importance, because they have clearly been transmitted from the Amphibia to all the higher Vertebrates. Man entirely resembles his amphibian

is well known that this hereditary number of the toes has assumed a very great practical importance from remote times; on it our whole system of enumeration (the decimal system applied to measure-




FIG. 26.—Fossil amphibian from the Permian, found in the Phosphorite near Dresden (*Temnodactylus amplexus*). (From Greiner) A skeleton of a young larva. B larva, restored, with teeth. C the adult form, natural size.

ancestors in this respect, and indeed in the whole structure of the bony skeleton of his five-toed extremities. A careful comparison of the skeleton of the frog with our own is enough to show this. It

ment of time, mass, weight, etc.) is based. There is absolutely no reason why there should be five toes in the fore and hind feet in the lowest Amphibia, the reptiles, and the higher Vertebrates, unless we

ascribe it to inheritance from a common stem-form. Heredity alone can explain it. It is true that we find less than five toes in many of the Amphibia and of the higher Vertebrates. But in all these cases we can prove that some of the toes atrophied, and were in time lost altogether.

The cause of this evolution of the five-toed foot from the many-toed fin in the amphibian ancestor must be sought in adaptation to the entire change of function that the limbs experienced in passing from an exclusively aquatic to a partly terrestrial life. The many-toed fin had been used almost solely for motion in the water; it had now also to support the body in creeping on the solid ground. This led to a modification both of the skeleton and the muscles of the limbs. The number of the fin-radial was gradually reduced, and sank finally to five. But these five remaining radii became much stronger. The soft cartilaginous radii became bony rods. The rest of the skeleton was similarly strengthened. Thus from the one-armed lever of the many-toed fish-fin arose the improved many-armed lever system of the five-toed amphibian limbs. The movements of the body gained in variety as well as in strength. The various parts of the skeletal system and correlated muscular system began to differentiate more and more. In view of the close correlation of the muscular and nervous systems, this also made great advance in structure and function. Hence we find, as a matter of fact, that the brain is much more developed in the higher Amphibia than in the fishes, the Dipneusta, and the lower Amphibia.

The first advance in organization that was occasioned by the adoption of life on land was naturally the construction of an organ for breathing air—a lung. This was formed directly from the floating-bladder inherited from the fishes. At first its function was insignificant beside that of the gills, the older organ for water-respiration. Hence we find in the lowest Amphibia, the gilled Amphibia, that, like the Dipneusta, they pass the greater part of their life in the water, and breathe water through gills. They only come to the surface at brief intervals, or creep on to the land, and then breathe air by their lungs. But some of the gilled Amphibia—the salamanders—remain entirely in the water when they are young, and afterwards spend most of their time on

land. In the adult state they only breathe air through lungs. The same applies to the most advanced of the Amphibia, the Batrachia (frogs and toads); some of them have entirely lost the gill-bearing larva form.* This is also the case with certain small, serpentine Amphibia, the Caecilia (which live in the ground like earth-worms).

The great interest of the natural history of the Amphibia consists especially in their intermediate position between the lower and higher Vertebrates. The lower Amphibia approach very closely to the



FIG. 41.—Larva of the Spotted Salamander (*Desmognathus maculatus*), seen from the ventral side. In the centre a pale-out gill hangs from the gill. The external gills are gracefully concealed. The two pairs of legs are still very small.

Dipneusta in their whole organization, live mainly in the water, and breathe by gills; but the higher Amphibia are just as close to the Amniotes, live mainly on land, and breathe by lungs. But in their younger state the latter resemble the former, and only reach the higher stage by a complete metamorphosis. The embryonic development of most of the

* The tree-frog of Macaronesia (*Hyla molleri* Steud.) keeps the gills on the mouth, and the tail and yellow on the eighth day of fetal life. On the ninth or tenth day after fecundation the frog emerges from the egg.

higher Amphibia still faithfully reproduces the stem-history of the whole class, and the various stages of the advance that was made by the lower Vertebrates in passing from aquatic to terrestrial life during the Devonian or the Carboniferous period are repeated in the spring by every frog that develops from an egg in our ponds.

The common frog leaves the egg in the shape of a larva, like the tailed salamander (Fig. 261), and this is altogether different



FIG. 261.—Larva of the common grass-frog (*Rana temporaria*), or "tadpole." *a*, mouth; *b*, a pair of suckers for fastening on to stones, if attached from which the gill-cover descends; *c*, behind it the gill-clefs, from which the branching gills (*d*) protrude; *e*, tail-muscles; *f*, cutaneous linings of the tail.

from the mature frog (Fig. 262). The short trunk ends in a long tail, with the form and structure of a fish's tail (*e*). There are no limbs at first. The respiration is exclusively branchial, first through external (*d*) and then internal gills. In harmony with this the heart has the same structure as in the fish, and consists of two sections—an atrium that receives the venous blood from the body, and a ventricle that forces it through the arteries into the gills.

We find the larva of the frog (or tadpole, *Gyrus*) in great numbers in our ponds every spring in this fish-form, using their muscular tails in swimming, just like the fishes and young Ascidia. When they have reached a certain size, the remarkable metamorphosis from the fish-form to the frog begins. A blind sac grows out of the gullet, and expands into a couple of spacious sacs: these are the lungs. The simple chamber of the heart is divided into two sections by the development of a partition, and there are at the same time considerable changes in the structure of the chief arteries. Previously all the blood went from the auricle through the aortic arches into the gills, but now only part of it goes to the gills, the other part passing to the lungs through the new-formed pulmonary artery. From this point arterial blood returns to the left auricle of the heart, while the venous blood gathers in the right auricle. As both auricles open into a single ventricle, this contains mixed blood. The dipneust form has now succeeded to the fish-form. In the further course of the metamorphosis the gills and the branchial vessels entirely disappear, and the respiration becomes exclusively pulmonary. Later, the long swimming tail is lost, and the frog now hops to the land with the legs that have grown meantime.

This remarkable metamorphosis of the Amphibia is very instructive in connection with our human genealogy, and is particularly interesting from the fact that the various groups of actual Amphibia have remained at different stages of their stem-history, in harmony with the biogenetic law. We have first of all a very low order of Amphibia—the *Sorobranchea* ("gilled-amphibia"), which retain their gills throughout life, like the fishes. In a second order of the salamanders the gills are lost in the metamorphosis, and when fully grown they have only pulmonary respiration. Some of the tailed Amphibia still retain the gill-clefs in the side of the neck, though they have lost the gills themselves (*Neurichthys*). If we force the larva of our salamanders (Fig. 261) and tribons to remain in the water, and prevent them from reaching the land, we can in favourable circumstances make them retain their gills. In this fish-like condition (they reach sexual maturity, and remain throughout life at the lower stage of the gilled Amphibia.

We have the reverse of this experiment in a Mexican gilled salamander, the fish-like axolotl (*Siredon pauciformis*). It was formerly regarded as a permanent gilled amphibian persisting throughout life at the fish-stage. But some of the hundreds of these animals that are kept in the Botanical Garden at Paris got on to the land for some reason or other, lost their gills, and changed into a form closely resembling the salamander (*Ambystoma*). Other species of the genus became sexually mature for the first time in this condition. This has been regarded as an astounding phenomenon, although every common frog and salamander repeats the metamorphosis in the spring. The whole change from the aquatic and gill-breathing animal to the terrestrial lung-breathing form may be followed step by step in this case. But what we

Their ancestors also had long tails and gills like the gilled Amphibia, as the tail and the gill-arches of the human embryo clearly show.

For comparative anatomical and ontogenetic reasons, we must not seek these amphibian ancestors of ours—as one would be inclined to do, perhaps—among the tail-less Batrachia, but among the tailed lower Amphibia.

The vertebrate form that comes next to the Amphibia in the series of our ancestors is a lizard-like animal, the earliest existence of which can be confidently deduced from the facts of comparative anatomy and ontogeny. The living *Hattersia* of New Zealand (Fig. 264) and the extinct *Eryonophis* of the Permian period (Fig. 265) are closely related to this important stem-form; we may call them the *Protamniotes*, or Primitive Amniotes.



FIG. 265.—Fossil tailed amphibian, from the Bobonian Cadeuriform (*Sedonia*). (From Frahm.) The body coat is retained on the left.

see here in the development of the individual has happened to the whole class in the course of its stem-history.

The metamorphosis goes farther in a third order of Amphibia, the *Batrachia* or *Anura*, than in the salamander. To this belong the various kinds of toads, ringed snakes, water-frogs, tree-frogs, etc. These lose, not only the gills, but also (sooner or later) the tail, during metamorphosis.

The ontogenetic loss of the gills and the tail in the frog and toad can only be explained on the assumption that they are descended from long-tailed Amphibia of the salamander type. This is also clear from the comparative anatomy of the two groups. This remarkable metamorphosis is, however, also interesting because it throws a certain light on the phylogeny of the tail-less apes and man.

All the Vertebrates above the Amphibia—or the three classes of reptiles, birds, and mammals—differ so much in their whole organisation from all the lower Vertebrates we have yet considered, and have so great a resemblance to each other, that we put them all together in a single group with the title of *Amniotes*. In these three classes alone we find the remarkable embryonic membrane, already mentioned, which we called the *amnion*; a xenogenetic adaptation that we may regard as a result of the sinking of the growing embryo into the yolk-sac.

All the Amniotes known to us—all reptiles, birds, and mammals (including man)—agree in so many important points of internal structure and development that their descent from a common ancestor can be affirmed with tolerable certainty. If the evidence of comparative

anatomy and ontogeny is ever entirely beyond suspicion, it is certainly the case here. All the peculiarities that accompany and follow the formation of the amnion, and that we have learned in our consideration of human embryology; all the peculiarities in the development of the organs which we will presently follow in detail; finally, all the principal special features of the internal structure of the full-grown Amniotes—prove so clearly the common origin of all the Amniotes from a single extinct stem-form that it is difficult to entertain the idea of their evolution from several independent stems. This unknown common stem-form is our Primitive Amniote (*Protamnion*). In outward appearance it was probably something between the salamander and the lizard.

It is very probable that some part of the Permian period was the age of the origin of the Protamnion. This follows from the fact that the Amphibia are not fully developed until the Carboniferous period, and that the first fossil reptiles (*Plesiosaurs*, *Homocentrus*, *Protosaurus*) are found towards the close of the Permian period. Among the important changes of the vertebrate organization that marked the rise of the first Amniotes from salamandrine Amphibia during this period the following three are especially noteworthy: the entire disappearance of the water-breathing gills and the conversion of the gill-arches into other organs, the formation of the allantois or primitive urinary sac, and the development of the amnion.

One of the most salient characteristics of the Amniotes is the complete loss of the gills. All Amniotes, even if living in water (such as sea-serpents and whales), breathe air through lungs, never water through gills. All the Amphibia (with very rare exceptions) retain their gills for some time when young, and have for a time (if not permanently) bronchial respiration; but after these there is no question of bronchial respiration. The Protamnion itself must have entirely abandoned water-breathing. Nevertheless, the gill-arches are preserved by heredity, and develop into totally different (in part rudimentary) organs—various parts of the bone of the tongue, the frame of the jaws, the organ of hearing, etc. But we do not find in the embryos of the Amniotes any trace of gill-leaves, or of real respiratory organs on the gill-arches.

With this complete abandonment of the gills is probably connected the formation of another organ, to which we have already referred in embryology—namely, the allantois or primitive urinary sac (cf. p. 166). It is very probable that the urinary bladder of the Dipneusts is the first structure of the allantois. We find in these a urinary bladder that proceeds from the lower wall of the hind end of the gut, and serves as receptacle for the renal secretions. This organ has been transmitted to the Amphibia, as we can see in the frog.

The formation of the amnion and the complete disappearance of the gills are the chief characteristics that distinguish the Amniotes from the lower Vertebrates we have hitherto considered. To these we may add several subordinate features that are transmitted to all the Amniotes, and are found in these only. One striking embryonic character of the Amniotes is the great curve of the head and neck in the embryo. We also find an advance in the structure of several of the internal organs of the Amniotes which raises them above the highest of the animals. In particular, a partition is formed in the simple ventricle of the heart, dividing into right and left chambers. In connection with the complete metamorphosis of the gill-arches we find a further development of the accessory organs. Also, there is a great advance in the structure of the brain, skeleton, muscular system, and other parts. Finally, one of the most important changes is the reconstruction of the kidneys. In all the earlier Vertebrates we have found the primitive kidneys as excretory organs, and these appear at an early stage in the embryos of all the higher Vertebrates up to man. But in the Amniotes these primitive kidneys cease to act at an early stage of embryonic life, and their function is taken up by the permanent or secondary kidneys, which develop from the terminal section of the pronephal ducts.

Taking all these peculiarities of the Amniotes together, it is impossible to doubt that all the animals of this group—all reptiles, birds, and mammals—have a common origin, and form a single blood-related stem. Our own race belongs to this stem. Man is, in every feature of his organization and embryonic development, a true Amniote, and has descended from the Protamnion with all the other

Amniotes. Though they appeared at the end (possibly even in the middle) of the Paleozoic age, the Amniotes only reached their full development during the Mesozoic age. The birds and mammals made their first appearance during this period. Even the reptiles show their greatest growth at this time, so that it is called "the reptile age." The extinct Protamniote, the ancestor of the whole group,

and only comes in contact with the Mammalia at its root, is the combined group of the reptiles and birds; these two classes may, with Huxley, be conveniently grouped together as the *Sauropsida*. Their common stem-form is an extinct lizard-like reptile of the order of the Rhynchocephalia. From this have been developed in various directions the serpents, crocodiles, tortoises, etc.—in a



FIG. 66.—The lizard (*Notoptis parvula*—*Sphenodon parvulus*) of New Zealand. The sole surviving rhynchocephale. (From Huxley.)

belongs in its whole organisation to the reptile class.

The genealogical tree of the amniote group is clearly indicated in its chief lines by their paleontology, comparative anatomy, and ontogeny. The group succeeding the Protamniote divided into two branches. The branch that will claim our whole interest is the class of the Mammalia. The other branch, which developed in a totally different direction,

word, all the members of the reptile class. But the remarkable class of the birds has also been evolved directly from a branch of the reptile group, as is now established beyond question. The embryos of the reptiles and birds are identical until a very late stage, and have an astonishing resemblance even later. Their whole structure agrees so much that no anatomist now questions the descent of the birds from the reptiles. On the other

hand, the mammal line has descended from the group of the Sauromammalia, a different branch of the Proreptilia. It is connected at its deepest roots with the reptile line, but it then diverges completely from it and follows a distinctive development. Man is the highest outcome of this class, the "crown of creation." The hypothesis that the three higher Vertebrate classes represent a single Amniote-stem, and that the common root of this stem is to be found in the amphibian class, is now generally admitted.

The instructive group of the Permian *Trocasauria*, the common root from which the divergent stems of the Sauropsids and mammals have issued, merits our particular attention as the stem-group of all the Amniotes. Fortunately a living representative of this extinct ancestral

263), of which well-preserved skeletons are found in the Solenhofen schists, is perhaps still more closely related to them.

Unfortunately, the numerous fossil remains of Permian and Triassic *Trocasauria* that we have found in the last two decades are, for the most part, very imperfectly preserved. Very often we can make only precarious inferences from these skeletal fragments as to the anatomic characters of the soft parts that went with the bony skeleton of the extinct *Trocasauria*. Hence it has not yet been possible to arrange these important fossils with any confidence in the ancestral series that descend from the Protamniotes to the Sauropsids on the one side and the Mammals on the other. Opinions are particularly divided as to the place in classification and the phylogenetic significance of the remarkable *Theromorpha*



FIG. 263.—*Heterosaurus paludicola*, a Jurassic proreptile from Kellheim. (From Kellheim.)

group has been preserved to our day; this is the remarkable lizard of New Zealand, *Hatteria punctata* (Fig. 264). Externally it differs little from the ordinary lizard; but in many important points of internal structure, especially in the primitive construction of the vertebral column, the skull, and the limbs, it occupies a much lower position, and approaches its amphibian ancestors, the *Stegocophala*. Hence *Hatteria* is the phylogenetically oldest of all living reptiles, an isolated survivor from the Permian period, closely resembling the common ancestor of the Amniotes. It must differ so little from this extinct form, our hypothetical Protamniote, that we put it next to the Proreptilia. The remarkable Permian *Palaeohatteria*, that Cresson discovered in the Plauen terrain at Dresden in 1882, belongs to the same group (Fig. 266). The Jurassic genus *Heterosaurus* (Fig.

Cope gives this name to a very interesting and extensive group of extinct terrestrial reptiles, of which we have only fossil remains from the Permian and Triassic strata. Forty years ago some of these *Theromorpha* (fresh-water animals) were described by Owen as *Amniotes*. But during the last twenty years the distinguished American paleontologists, Cope and Osborn, have greatly increased our knowledge of them, and have claimed that the stem-forms of the Mammals must be sought in this order. As a matter of fact, the *Theromorpha* are nearer to the Mammals in the chief points of structure than any other reptiles. This is especially true of the *Therodontia*, to which the *Purcosaurus* and *Pelycosaurus* belong (Fig. 267). The whole structure of their pelvis and hind-feet has attained the same form as in the Monotremes, the lowest Mammals. The formation of the

scapula and the quadrate bone shows an approach to the Mammals such as we find in no other group of reptiles. The teeth also are already divided into incisors, canines, and molars. Nevertheless, it is very doubtful whether the Theromorphs really are in the ancestral line of the Sauromammals, or lead direct from the Tocosaurs to the earliest Mammals. Other experts on this group believe that it is an independent legion of the reptiles, connected, perhaps, at its lowest root, with the Sauromammals, but developed quite independently of the Mammals—though parallel to them in many ways.

One of the most important of the zoological facts that we rely on in our investigation of the genealogy of the human race is the position of man in the Mammal class. However different the views of zoologists may have been as to this position in detail, and as to his relations to the apes, no scientist has ever doubted that man is a true mammal in his whole organization and development. Linné drew attention to this fact in the first edition of his famous *Systema Naturæ* (1735). As will be seen in any museum of anatomy or any manual of comparative anatomy, the human frame has all the characteristics that are common to the Mammals and distinguish them conspicuously from all other animals.

If we examine this undoubted fact from the point of view of phylogeny, in the light of the theory of descent, it follows at once that man is of a common stem with all the other Mammals, and comes from the same root as they. But the various features in which the Mammals agree and by which they are distinguished are of such a character as to make a polyphyletic hypothesis quite inadmissible. It is impossible to entertain the idea that all the living and extinct Mammals come from a number of separate roots. If we accept the general theory of evolution, we are bound to admit the monophyletic hypothesis of the descent of all the Mammals (including man) from a single mammalian stem-form. We may call this long extinct root-form and its earliest descendants (a few genera of one family) "primitive mammals" or "stem-mammals" (*Protomammalia*). As we

have already seen, this root-form developed from the primitive *Proreptile* stem in a totally different direction from the birds, and soon separated from the main stem of the reptiles. The differences between the Mammals and the reptiles and birds are so important and characteristic that we can assume with complete confidence this division of the vertebrate stem at the commencement of the development of the Amniotes. The reptiles and birds, which we group together as the *Sauropseds*, generally agree in the characteristic structure of the skull and brain, and this is notably different from that of the Mammals. In most of the reptiles and birds the skull is connected with the first cervical vertebra (the *axis*)



FIG. 464.—Skull of a Permian lizard (*Palaeopterus lucasensis*). (From Osborn.) a, nasal bone; b, frontal bone; c, lacrimal bone; d, postorbital bone; e, surangular bone; f, dentary bone; g, vomer; h, intermaxillary.

by a single, and in the Mammals (and Amphibia) by a double, condyle at the back of the head. In the former the lower jaw is composed of several pieces, and connected with the skull so that it can move by a special maxillary bone (the *quadrate*); in the Mammals the lower jaw consists of one pair of bony pieces, which articulate directly with the temporal bone. Further, in the Sauropsids the skin is clothed with scales or feathers; in the Mammals with hair. The red blood-cells of the former have a nucleus; those of the latter have not. In fine, two quite characteristic features of the Mammals, which distinguish them not only from the birds and reptiles, but from all other animals, are the possession of a

complete diaphragm and of mammary glands that produce the milk for the nutrition of the young. It is only in the Mammals that the diaphragm forms a transverse partition of the body-cavity, completely separating the pectoral from the abdominal cavity. It is only in the mammals that the mother suckles its young, and this rightly gives the name to the whole class (*mammae* = breasts).

From these pregnant facts, of comparative anatomy and ontogeny it follows absolutely that the whole of the Mammals belong to a single natural stem, which

and the brain; (3) the development of a hairy coat; (3) the complete formation of the diaphragm; and (4) the construction of the mammary glands and adaptation to suckling. Other important changes of structure proceeded step by step with these.

The epoch at which these important advances were made, and the foundation of the Mammal class was laid, may be put with great probability in the first section of the Mesozoic or secondary age—the Triassic period. The oldest fossil remains of mammals that we know were

found in strata that belong to the earliest Triassic period—the upper Keuper. One of the earliest forms is the genus *Docuatherium*, from the North American Triassic (Fig. 268). Their teeth still strikingly recall those of the Polycaurids. Hence we may assume that this small and probably insectivorous mammal belonged to the stem-group of the Placentalia. We do not find any positive trace of the third and most advanced division of the Mammalia—the Placentals. These (including man) are much younger, and we do not find indisputable fossil remains of them until the Cenozoic age, or the Tertiary period. This paleontological fact is very important, because it fully harmonises with the evolutionary succession of the Mammal orders that is deduced from their comparative anatomy and ontogeny.

The later science teaches us that the whole Mammal class divides into three main groups or sub-classes, which correspond to three successive phylogenetic stages. These three stages, which also represent three important

stages in our human genealogy, were first distinguished in 1816 by the eminent French zoologist, Geoffroy, and received the names of *Ornithodelphia*, *Didelphia*, and *Monodelphia*, according to the construction of the female organ (*delphis* = uterus or womb). Huxley afterwards gave them the names of *Prototheria*, *Metatheria*, and *Eutheria*. But the three sub-classes differ so widely from each other, not only in the construction of the sexual organs, but in many other respects also, that we may confidently draw up the following




FIG. 268.—Skull of a Triassic theromorph (*Giveterium glenensis*). a, from the right; b, from below; c, from above; d, enlarged tooth. N, nostril; Mx, maxilla; Pt, prefrontal; A, frontal bone; Pa, parietal; Pl, postorbital; Ml, lower jaw.

branched off at an early date from the reptile-root. It follows further with the same absolute certainty that the human race is also a branch of this stem. Man shares all the characteristics I have described with all the Mammals, and differs in them from all other animals. Finally, from these facts we deduce with the same confidence those advances in the vertebrate organisation by which one branch of the Sauriomammals was converted into the stem-form of the Mammals. Of these advances the chief were: (1) The characteristic modification of the skull

important phylogenetic thesis: The Monodelphia or Placentals descend from the Didelphia or Marsupials; and the latter, in turn, are descended from the Monotremes or Ornithodelphia.

Thus we must regard as the twenty-first stage in our genealogical tree the earliest and lowest chief group of the Mammalia—the sub-class of the Monotremes ("cloaca-animals," Ornithodelphia, or Prototheria, Figs 269 and 270). They take their name from the cloaca which they share with all the lower Vertebrates. This cloaca is the common outlet for the passage of the excrement, the urine, and the sexual products. The urinary ducts and sexual canals open into the hindmost part of the gut, while in all the other Mammals they are separated from the rectum and anus. The latter have a special urogenital outlet (*Foramen urogenitale*). The bladder also opens into the cloaca in the Monotremes, and, indeed, apart from the two urinary ducts; in all the other Mammals the latter open directly into the bladder. It was proved by Hawke and Caldwell in 1883 that the Monotremes lay large eggs like the reptiles, while all the other Mammals are viviparous. In 1844 Richard Brown further proved that these large eggs, rich in food-yolk, have a partial segmentation and discoid gastrulation, as I had hypothetically assumed in 1879, here again they resemble their reptilian ancestors. The construction of the mammary gland is also peculiar in the Monotremes. In these the glands have no teats for the young animal to suck, but there is a special part of the breast pierced with holes like a sieve, from which the milk issues, and the young Monotreme must lick it off. Further, the brain of the Monotremes is very little advanced. It is smaller than that of any of the other Mammals. The fore-brain or cerebrum, in particular, is so small that it does not cover the cerebellum. In the skeleton (Fig. 270) the formation of the scapula among other parts is curious; it is quite different from that of the other Mammals, and rather agrees with that of the reptiles and Amphibia. Like these, the Monotremes have a strongly developed *caracululum*. From these and other less prominent characteristics it follows absolutely that the Monotremes occupy the lowest place among the Mammals, and represent a transitional group between the *Tocostauria* and the

rest of the Mammals. All these remarkable reptilian characters must have been possessed by the stem-form of the whole mammal class, the *Protomammal* of the Triassic period, and have been inherited from the *Proreptiles*.

During the Triassic and Jurassic periods the sub-class of the Monotremes was represented by a number of different stem-mammals. Numerous fossil remains of them have lately been discovered in the Mesozoic strata of Europe, Africa, and America. To-day there are only two surviving specimens of the group, which we place together in the family of the duck-bills, *Ornithosoma*. They are confined to Australia and the neighbouring island of Van Diemen's Land (or Tasmania); they become scarcer every year, and will soon, like their blood-relatives, be counted among the extinct animals. One form lives in the rivers,



FIG. 268.—Lower jaw of a Primitive Mammal or Protomammal (*Ornithosoma valens*) from the North American Triassic. *a* molar, *b* canine, *c* premolar, *d* incisor, *e* *f* *Foramen urogenitale*.

and builds subterranean dwellings on the banks; this is the *Ornithosoma paradoxum*, with webbed feet, a thick soft fur, and broad flat jaws, which look very much like the bill of a duck (Figs 269, 270). The other form, the land duck-bill, or spiny ant-eater (*Echidna hyaline*), is very much like the anteaters in its habits and the peculiar construction of its thin snout and very long tongue; it is covered with needles, and can roll itself up like a hedgehog. A cognate form (*Peraiasma Bryant*) has lately been found in New Guinea.

These modern *Ornithosoma* are the scattered survivors of the vast Mesozoic group of Monotremes; hence they have the same interest in connection with the stem history of the Mammals as the living stem-reptiles (*Hatteria*) for that of the reptiles, and the isolated *Acrania* (*Amphioxus*) for the phylogeny of the Vertebrate stem.

The Australian duck-bills are distinguished externally by a toothless bird-

like beak or snout. This absence of real bony teeth is a late result of adaptation, as in the toothless Placentals (*Edentata*, armadillos and ant-eaters). The extinct Monotremes, to which the *Proganimalia* belonged, must have had developed teeth, inherited from the reptiles. Lastly small rudiments of real molars have been discovered in the young of the Ornitho-

Marsupials retain a great part of the characteristics of the Monotremes, they




FIG. 279.—The Ornithorhynchus or Duck-bill.
(*Ornithorhynchus anatinus*.)

rhynchus, which has horny plates in the jaws instead of real teeth.

The living Ornithostoma and the stem-forms of the Marsupials (or *Didelphes*) must be regarded as two widely diverging lines from the *Proganimalia*. This second sub-class of the Mammalia is very interesting as a perfect intermediate stage between the other two. While the




FIG. 280.—Skeleton of the preceding.

have also acquired some of the chief features of the Placentals. Some features

are also peculiar to the Marsupials, such as the construction of the male and female sexual organs and the form of the lower jaw. The Marsupials are distinguished by a peculiar hook-like bony process that bends from the corner of the lower jaw and points inwards. As most of the Placentals have not this process, we can, with some probability, recognise the Marsupial from this feature alone. Most of the mammal remains that we have from the Jurassic and Cretaceous deposits are merely lower jaws, and most of the jaws found in the Jurassic deposits at Stonesfield and Purbeck have the peculiar hook-like process that characterises the lower jaw of the Marsupial. On the strength of this paleontological fact, we may suppose that they belonged to Marsupials. Placentals do not seem to have existed at the middle of the Mesozoic age—not until towards its close (in the Cretaceous period). At all events, we have no fossil remains of indubitable Placentals from that period.

The existing Marsupials, of which the plant-eating kangaroo and the carnivorous opossum (Fig. 272) are the best known, differ a good deal in structure, shape, and size, and correspond in many respects to the various orders of Placentals. Most of them live in Australia, and a small part of the Australian and East Malayan islands. There is now not a single living Marsupial on the mainland of Europe, Asia, or Africa. It was very different during the Mesozoic and even during the Cenozoic age. The sedimentary deposits of these periods contain a great number and variety of marsupial remains, sometimes of a colossal size, in various parts of the earth, and even in Europe. We may infer from this that the existing Marsupials are the remnant of an extensive earlier group that was distributed all over the earth. It had to give way in the struggle for life to the more powerful Placentals during the Tertiary period. The survivors of the group were able to keep alive in Australia and South America because the one was completely separated from the other parts of the earth during the whole of the Tertiary period, and the other during the greater part of it.

From the comparative anatomy and nomenclature of the existing Marsupials we may draw very interesting conclusions as to their intermediate position between

the earlier Monotremes and the later Placentals. The defective development of the brain (especially the cerebrum), the possession of marsupial bones, and the simple construction of the alveolar (without any placenta as yet) were inherited by the Marsupials, with many other features, from the Monotremes, and preserved. On the other hand, they have lost the independent bone (coracodromus) at the shoulder-blade. But we have a more important advance in the disappearance of the cloaca; the rectum and anus are separated by a partition from the uro-genital opening (*anus sequestratus*). Moreover, all the Marsupials have teats on the mammary glands, at which the new-born animal sucks. The teats pass into the cavity of a pouch or pocket on the ventral side of the mother, and this is supported by a couple of marsupial bones. The young are born in a very imperfect condition, and carried



FIG. 271.—Lower jaw of a *Promastomys* (*Dipodops pringi*), from the Jurassic of the Eocene series. (From Mehl.)

by the mother for some time longer in her pouch, until they are fully developed (Fig. 272). In the giant kangaroo, which is as tall as a man, the embryo only develops for a month in the uterus, is then born in a very imperfect state, and finishes its growth in the mother's pouch (*marsupium*); it remains in this about nine months, and at first hangs continually on to the teat of the mammary gland.

From these and other characteristics (especially the peculiar construction of the internal and external sexual organs in male and female) it is clear that we must conceive the whole sub-class of the Marsupials as one stem group, which has been developed from the *Promastomids*. From one branch of these Marsupials (possibly from more than one) the stem-forms of the higher Mammals, the Placentals, were afterwards evolved. Of the existing forms of the Marsupials,

which have undergone various modifications through adaptation to different environments, the family of the opossums (*Didelphidae* or *Prosimiidae*) seems to be the oldest and nearest to the common stem-form of the whole class. To this family belong the crab-eating opossum of Brazil (Fig. 272) and the opossum of

Lessors, were evolved directly from the opossum. We must not forget, however, that the conversion of the five-toed foot into a prehensile hand is polyphyletic. By the same adaptation to climbing trees the habit of grasping their branches with the feet has in many different cases brought about that opposition of the



FIG. 272.—The crab-eating Opossum (*Phylander carnifex*). The female has three young in the pouch. (From Bodin.)

Virginia, on the embryology of which Selenka has given us a valuable work (cf. Figs. 63-7 and 732-5). These *Didelphids* climb trees like the apes, grasping the branches with their hand-shaped hind feet. We may conclude from this that the stem-forms of the *Primates*, which we must regard as the earliest

thumb or great toe to the other toes which makes the hand prehensile. We see this in the climbing lizards (*Chamaeleon*), the birds, and the tree-dwelling mammals of various orders.

Some zoologists have lately advanced the opposite opinion, that the *Marsupials* represent a completely independent sub-

class of the Mammals, with no direct relation to the Placentals, and developing independently of them from the Monotremes. But this opinion is untenable if we examine carefully the whole organization of the three sub-classes, and do not lay the chief stress on incidental features and secondary adaptations (such as the

formation of the marsupium). It is then clear that the Marsupials—viviparous Mammals without placenta—are a necessary transition from the oviparous Monotremes to the higher Placentals with chorion-villi. In this sense the Marsupial class certainly contains some of man's ancestors.

CHAPTER XXIII.

OUR APE ANCESTORS

THE long series of animal forms which we must regard as the ancestors of our race has been confined within narrower and narrower circles as our phylogenetic inquiry has progressed. The great majority of known animals do not fall in the line of our ancestry, and even within the vertebrate stem only a small number are found to descend. In the most advanced class of the stem, the mammals, there are only a few families that belong directly to our genealogical tree. The most important of these are the apes and their predecessors, the half-apes, and the earliest Placentals (*Prochordata*).

The Placentals (also called *Chordata*, *Monodelphia*, *Anthra* or *Epithemia*) are distinguished from the lower mammals we have just considered, the Monotremes and Marsupials, by a number of striking peculiarities. Man has all these distinctive features, that is a very significant fact. We may, on the ground of the most careful comparative-anatomical and ontogenetic research, formulate the thesis: "Man is in every respect a true Placental." He has all the characteristics of structure and development that distinguish the Placentals from the two lower divisions of the mammals, and, in fact, from all other animals. Among these characteristics we must especially notice the more advanced development of the brain. The fore-brain or cerebrum especially is much more developed in them than in the lower animals. The *corpus callosum*, which forms a sort of wide bridge connecting the two hemispheres of the cerebrum, is only fully formed in the

Placentals; it is very rudimentary in the Marsupials and Monotremes. It is true that the lowest Placentals are not far removed from the Marsupials in cerebral development, but within the placental group we can trace an unbroken gradation of progressive development of the brain, rising gradually from this lowest stage up to the elaborate psychic organ of the ape and man. The human soul—a physiological function of the brain—is in reality only a more advanced apowoul.

The mammary glands of the Placentals are provided with teats like those of the Marsupials, but we never find in the Placentals the pouch in which the latter carry and suckle their young. Nor have they the marsupial bones in the ventral wall at the anterior border of the pelvis, which the Marsupials have in common with the Monotremes, and which are formed by a partial ossification of the sinews of the inner oblique abdominal muscle. There are merely a few insignificant remnants of them in some of the Carnivora. The Placentals are also generally without the hook-shaped process at the angle of the lower jaw which is found in the Marsupials.

However, the feature that characterizes the Placentals above all others, and that has given its name to the whole subclass, is the formation of the placenta. We have already considered the formation and significance of this remarkable embryonic organ when we traced the development of the chorion and the allantois in the human embryo (pp. 165-9). The tertiary sac or the allantois, the

curious vesicle that grows out of the hind part of the gut, has essentially the same structure and function in the human embryo as in that of all the other Amniotes (cf. Figs. 194-6). There is a quite secondary difference, on which great stress has wrongly been laid, in the fact that in man and the higher apes the original cavity of the allantois quickly degenerates, and the rudiment of it sticks out as a solid projection from the primitive gut. The thin wall of the allantois consists of the same two layers or membranes as the wall of the gut—the gut-gland layer within and the gut-fibre

and remarkable structure that we call the *placenta*.

The placenta is formed by the branches of the blood-vessels in the wall of the allantois growing into the hollow ectodermic tufts (villi) of the chorion, which run into corresponding depressions in the mucous membrane of the womb. The latter also is richly permeated with blood-vessels which bring the mother's blood to the embryo. As the partition in the villi between the maternal blood-vessels and those of the fetus is extremely thin, there is a direct exchange of fluid between the two, and this is of the greatest importance in the nutrition of the young mammal. It is true that the maternal vessels do not entirely pass into the fetal vessels, so that the two kinds of blood are simply mixed. But the partition between them is so thin that the nutritive fluid easily transudes through it. By means of this transudation or diosmosis the exchange of fluids takes place without difficulty. The larger the embryo is in the placenta, and the longer it remains in the womb, the more necessary it is to have special structures to meet its great consumption of food.

In this respect there is a very conspicuous difference between the lower and higher mammals. In the Marsupials, in which the embryo is only a comparatively short time in the womb and is born in a very immature condition, the vascular arrangements in the yolk-sac and the allantois suffice for its nutrition, as we find them in the Monotremes, birds, and reptiles. But in the Placentals, where gestation lasts a long time, and the embryo reaches its full development under the protection of its enveloping membranes, there has to be a new mechanism for the direct supply of a large quantity of food, and this is admirably met by the formation of the placenta.

Branches of the blood-vessels penetrate into the chorion-villi from within, starting from the gut-fibre layer of the allantois, and bringing the blood of the fetus through the umbilical vessels (Fig. 273 *2A*). On the other hand, a thick network of blood-vessels develops in the mucous membrane that clothes the inner surface of the womb, especially in the region of the depressions into which the chorion-villi penetrate (*2B*). This network of arteries contains maternal blood, brought by the uterine vessels. As the connective tissue between the enlarged capillaries of



FIG. 273.—Fetal membranes of the human embryo (diagrammatic), as the thick muscular wall of the womb. *pl*, placenta (the inner layer (*pl* *int*) of which penetrates into the chorion-villi (*ch*) with its processes). *al*, allantois, the smooth chorion, a network of blood-vessels, an amniotic sheet of the umbilical cord (which pumps blood into the vessel of the embryo—this gives *2A*), the villous part, the yolk sac, *2B*, the chorion (inner and outer). The uterine cavity (*ut*) opens later into the vagina and above on the right into an oviduct (*ov*). (From Abderhalden.)

layer without. In the gut-fibre layer of the allantois there are large blood-vessels, which serve for the nutrition, and especially the respiration, of the embryo—the umbilical vessels (*n* 170). In the reptiles and birds the allantois enlarges into a spacious sac, which encloses the embryo with the amnion, and does not combine with the outer fetal membrane (the chorion). This is the case also with the lowest mammals, the oviparous Monotremes and most of the Marsupials. It is only in some of the later Marsupials (*Peramelids*) and all the Placentals that the allantois develops into the distinctive

the uterus disappears, wide cavities filled with maternal blood appear, and into these the chorion-villi of the embryo penetrate. The sum of these vessels of both kinds, that are so intimately correlated at this point, together with the connective and enveloping tissue, is the *placenta*. The placenta consists, therefore, properly speaking, of two different though intimately connected parts—the fetal placenta (Fig. 273 *ch*) within and the maternal or uterine placenta (*pl*) without. The latter is made up of the mucous coat of the uterus and its blood-vessels, the former of the tufted chorion and the umbilical vessels of the embryo (cf. Fig. 196).

The manner in which these two kinds of vessels combine in the placenta, and the structure, form, and size of it, differ a good deal in the various Placentals; to some extent they give us valuable data

at birth the fetal placenta alone comes away; the uterine placenta is not torn away with it.

The formation of the placenta is very different in the second and higher section of the Placentals, the *Deciduata*. Here again the whole surface of the chorion is thickly covered with the villi in the beginning. But they afterwards disappear from one part of the surface, and grow proportionately thicker on the other part. We thus get a differentiation between the smooth chorion (*chorion laeve*, Fig. 273 *ch*) and the thickly-tufted chorion (*chorion frondosum*, Fig. 273 *ch*). The former has only a few small villi or none at all, the latter is thickly covered with large and well-developed villi; this alone now constitutes the placenta. In the great majority of the *Deciduata* the placenta has the same shape as in man (Fig. 197, 200)—namely



FIG. 274.—Skull of a fossil lemur (*Adapis parvus*), from the Miocene of Quarry A, Leptot, near the right, half natural size. B, lower jaw. C, lower molar. D, incisor. E, canine. F, preincisor or molars.

for the natural classification, and therefore the phylogeny, of the whole of this sub-class. On the ground of these differences we divide it into two principal sections; the lower Placentals or *Indecidua*, and the higher Placentals or *Deciduata*.

To the *Indecidua* belong three important groups of mammals: the Lemurs (*Prosimia*), the Ungulates (tapirs, horses, pigs, ruminants, etc.), and the Cetacea (dolphins and whales). In these *Indecidua* the villi are distributed over the whole surface of the chorion (or its greater part), either singly or in groups. They are only loosely connected with the mucous coat of the uterus, so that the whole fetal membrane with its villi can be easily withdrawn from the uterine depressions like a hand from a glove. There is no real coalescence of the two placentas at any part of the surface of contact. Hence

a thick, circular disk like a cake; so we find in the *Insectivora*, *Chiroptera*, *Rodentia*, and *Apes*. This *discoplacenta* lies on one side of the chorion. But in the *Sarcotheria* (both the *Carnivora* and the seals, *Pinnipedia*) and in the elephant and several other *Deciduata* we find a *somoplacenta*; in these the rich mass of villi runs like a girdle round the middle of the elliptical chorion, the two poles of it being free from them.

Still more characteristic of the *Deciduata* is the peculiar and very intimate connection between the *chorion frondosum* and the corresponding part of the mucous coat of the womb, which we must regard as a real coalescence of the two. The villi of the chorion push their branches into the blood-filled spaces of the coat of the uterus, and the vessels of each loop together so intimately that it is no longer possible to separate the fetal

from the maternal placenta; they form henceforth a compact and apparently simple placenta. In consequence of this coalescence, a whole piece of the lining of the womb comes away at birth with the foetal membrane that is interlaced with it. This piece is called the "falling-away" membrane (*decidua*). It is also called the serous (spongy) membrane, because

the Deciduata. (Cf. Figs. 199, 200, pp. 168-70.)

In the various orders of the Deciduata the placenta differs considerably both in outer form and internal structure. The extensive investigations of the last ten years have shown that there is more variation in these respects among the higher mammals than was formerly supposed. The physiological work of this important embryonic organ, the nutrition of the foetus during its long sojourn in the womb, is accomplished in the various groups of the Placentals by very different and sometimes very elaborate structures. They have lately been fully described by Hans Strahl.

The phylogeny of the placenta has become more intelligible from the fact that we have found a number of transitional forms of it. Some of the Marsupials (*Peromyscus*) have the beginning of a placenta. In some of the Lemurs (*Lemur*) a divided placenta with decidua is developed.

While these important results of comparative embryology have been throwing further light on the close blood-relationship of man and the anthropoid apes in the last few years (p. 172), the great advance of paleontology has at the same time been affording us a deeper insight into the stem-history of the Placental group. In the seventh chapter of my *Systematic Phylogeny of the Terrestrial Vertebrates* I advanced the hypothesis that the Placentals form a single stem with many branches, which has been evolved from an older group of the Marsupials (*Pro-*

dolphia). The four great legions of the Placentals—Rodents, Ungulates, Carnivora, and Primates—are sharply separated to-day by important features of organization. But if we consider their extinct ancestors of the Tertiary period, the differences gradually disappear, the deeper we go in the Cenozoic deposits; in the end we find that they vanish altogether.



FIG. 275.—The Slender Lemur (*Stenops gracilis*) of Cochin, a tail-less lemur.

it is pierced like a sieve or sponge. All the higher Placentals that have this decidua are classed together as the "Deciduata." The tearing away of the decidua at birth naturally causes the mother to lose a quantity of blood, which does not happen in the Indocidua. The lost part of the uterine coat has to be replaced by a new growth after birth in

The primitive stem-forms of the Rodents (*Euroscytodes*), the Ungulates (*Chondylarthrus*), the Carnassals (*Isiobondus*), and the Primates (*Lemurinae*) are so closely related at the beginning of the Tertiary period that we might group them together as different families of one order, the Proplacentalia (*Mallotheria* or *Prochazalia*).

Hence the great majority of the Placentalia have no direct and close relationship to man, but only the legroom of the Primates. This is now generally divided into three orders—the half-apes (*Prosimiæ*), apes (*Simiæ*), and man (*Anthropi*). The lemurs or half-apes are the stem-group, descending from the older *Mallotheria* of the Cretaceous period. From them the apes were evolved in the Tertiary period, and man was formed from these towards its close.

The Lemurs (*Prosimiæ*) have few living representatives. But they are very interesting, and are the last survivors of a once extensive group. We find many fossil remains of them in the older Tertiary deposits of Europe and North America, in the Eocene and Miocene. We distinguish two sub-orders, the fossil *Lemurinae* and the modern *Lemurogona*. The earliest and most primitive forms of the Lemurinae are the Pachylemurs (*Hypomadenia*), they come next to the earliest Placentalia (*Prochazalia*), and have the typical full dentition, with forty-four teeth ($\frac{2}{1} \frac{1}{1} \frac{2}{1}$). The Neotelmurs (*Adapida*, fig. 274) have only forty teeth, and have lost an incisor in each jaw ($\frac{2}{1} \frac{1}{1} \frac{2}{1}$). The dentition is still further reduced in the Lemurogona (*Autotelmurs*), which usually have only thirty-six teeth ($\frac{2}{1} \frac{1}{1} \frac{2}{1}$). These living varieties are scattered far over the southern part of the Old World. Most of the species live in Madagascar, some in the Sunda Islands, others on the mainland of Asia and Africa. They are generally and melancholic animals, they live a quiet life, climbing trees, and eating fruit and insects. They are of different kinds. Some are closely related to the Marsupials (especially the opossum). Others (*Macrocterus*) are nearer to the Insectivora, others, again (*Chomomyx*) to the Rodents. Some of the lemurs (*Homocidus*) approach closely to the true apes. The numerous fossil remains of half-apes and apes that have been recently found in the Tertiary deposits justify us in thinking that man's ancestors were represented by several different

species during this long period. Some of these were almost as big as man, such as the diluvial Lemurogona *Megaladaps* of Madagascar.

Next to the lemurs come the true apes (*Simiæ*), the twenty-ninth stage in our ancestry. It has been beyond question for some time now that the apes approach nearest to man in every respect of all the animals. Just as the lowest apes come close to the lemurs, so the highest come next to man. When we carefully study the comparative anatomy of the apes and man, we can trace a gradual and uninterrupted advance in the organization of the ape up to the purely human frame, and, after impartial examination of the "ape-



FIG. 276.—The white-nosed ape (*Ceropithecus*).

problem" that has been discussed of late years with such passionate interest, we come infallibly to the important conclusion, first formulated by Huxley in 1863: "Whatever systems of organs we take, the comparison of their modifications in the series of apes leads to the same result: that the anatomic differences that separate man from the gorilla and chimpanzee are not as great as those that separate the gorilla from the lower apes." Translated into phylogenetic language, this "pithomacra-law," formulated in such masterly fashion by Huxley, is quite equivalent to the popular saying "Man is descended from the apes."

In the very first exposition of his profound natural classification (1735) Linné

placed the anthropoid mammals at the head of the animal kingdom, with three genera: man, the ape, and the sloth. He afterwards called them the "Primates"—the "lords" of the animal world; he then also separated the lemur from the true ape, and rejected the sloth. Later zoologists divided the order of Primates

and *Quadrumanus* was retained by Cuvier and most of the subsequent zoologists. It seems to be extremely important, but, as a matter of fact, it is totally wrong. This was first shown in 1863 by Huxley, in his famous *Man's Place in Nature*. On the strength of careful comparative-anatomical research he proved that the



FIG. 277.—The gibbon (*Cynopithecus brevirostris*). (From Broder.)

First the Göttingen anatomist, Blumenbach, founded a special order for man, which he called *Bimana* ("two-handed"); in a second order he united the apes and lemurs under the name of *Quadrumanus* ("four-handed"); and a third order was formed of the distantly-related *Chiroptera* (bats, etc.). The separation of the *Bimana*

apes are just as truly "two-handed" as man; or, if we prefer to reverse it, that man is as truly four-handed as the ape. He showed convincingly that the ideas of hand and foot had been wrongly defined, and had been improperly based on physiological instead of morphological grounds. The circumstance that we oppose the

thumb to the other four fingers in our hand, and so can grasp things, seemed to be a special distinction of the hand in contrast to the foot, in which the corresponding great toe cannot be opposed in this way to the others. But the apes can grasp with the hind-foot as well as the fore, and so were regarded as quadrumanous. However, the inability to grasp that we find in the foot of civilized man is a consequence of the habit of clothing it with tight coverings for thousands of years. Many of the bare-footed lower races of men, especially among the negroes, use the foot very freely in the same way as the hand. As a result of early habit and continued practice, they can grasp with the foot (in climbing trees, for instance) just as well as with the hand. Even new-born infants of our own race can grasp very strongly with the great toe, and hold a spoon with it as firmly as with the hand. Hence the physiological distinction between hand and foot can neither be pressed very far, nor has it a scientific basis. We must look to morphological characters.

As a matter of fact, it is possible to draw such a sharp morphological distinction—a distinction based on anatomic structure—between the fore and hind extremity. In the formation both of the bony skeleton and of the muscles that are connected with the hand and foot before and behind there are material and constant differences, and these are found both in man and the ape. For instance, the number and arrangement of the smaller bones of the hand and foot are quite different. There are similar constant differences in the muscles. The hind extremity always has three muscles (a short flexor muscle, a short extensor muscle, and a long calf-muscle) that are not found in the fore extremity. The arrangement of the muscles also is different before and behind. These characteristic differences between the fore and hind extremities are found in man as well as the ape. There can be no doubt, therefore, that the ape's foot deserves that name just as much as the human foot does, and that all true apes are just as "bimanous" as man. The common distinction of the apes as "quadrumanous" is altogether wrong morphologically.

But it may be asked whether, quite apart from this, we can find any other features that distinguish man more sharply from the ape than the various

species of apes are distinguished from each other. Huxley gave as complete and demonstrative a reply to this question that the opposition still raised on many sides is absolutely without foundation. On the ground of careful comparative-anatomical research, Huxley proved that in all morphological respects the differences between the highest and lowest apes are greater than the corresponding differences between the highest apes and man. He thus restored Liéna's order of the Primates (excluding the bats), and divided it into three sub-orders, the first composed of the half-apes (*Lemnidae*), the second of the true apes (*Simidae*), the third of men (*Anthropidae*).

But, as we wish to proceed quite consistently and impartially on the laws of systematic logic, we may, on the strength of Huxley's own law, go a good deal farther in this division. We are justified in going at least one important step further, and assigning man his natural place within one of the sections of the order of apes. All the features that characterize this group of apes are found in man, and not found in the other apes. We do not seem to be justified, therefore, in founding for man a special order distinct from the apes.

The order of the true apes (*Simiæ* or *Plathecæ*)—excluding the lemurs—has long been divided into two principal groups, which also differ in their geographical distribution. One group (*Hæmaphyshecæ*, or western apes) live in America. The other group, to which man belongs, are the *Eopithecæ* or eastern apes; they are found in Asia and Africa, and were formerly in Europe. All the eastern apes agree with man in the features that are chiefly used in zoological classification to distinguish between the two simian groups, especially in the dentition. The objection might be raised that the teeth are too subordinate an organ physiologically for us to lay stress on them in so important a question. But there is a good reason for it; it is with perfect justice that zoologists have for more than a century paid particular attention to the teeth in the systematic division and arrangement of the orders of mammals. The number, form, and arrangement of the teeth are much more faithfully inherited in the various orders than most other characters.

Hence the form of dentition in man is very important. In the fully developed

condition we have thirty-two teeth; of these eight are incisors, four canines, and twenty molars. The eight incisors, in the middle of the jaws, have certain

Next to these, at each side of both jaws, is a canine (or "eye tooth"), which is larger than the incisors. Sometimes it is very prominent in man, as it is in most




Fig. 1st-5th—Skeletons of a man (Fig. 1st) and the four anthropoid apes (Fig. 2nd Gibbon, Fig. 3rd Chimpanzee, Fig. 4th Gorilla, Fig. 5th Orang: Fig. 1st, Hens; Fig. 2nd, Jeffries; Fig. 3rd, Anthropologist; Fig. 4th, Garth; Fig. 5th, Hens).

characteristic differences above and below. In the upper jaw the inner incisors are larger than the outer; in the lower jaw the inner are the smaller.

apes and many of the other mammals, and forms a sort of tusk. Next to this there are five molars above and below on each side, the first two of which (the

"pre-molars") are small, have only one root, and are included in the change of tooth; the three back ones are much larger, have two roots, and only come with the second teeth. The apes of the Old World, or all the living or fossil apes of Asia, Africa, and Europe, have the same dentition as man.

On the other hand, all the American apes have an additional pre-molar in each half of the jaw. They have six molars above and below on each side, or thirty-six teeth altogether. This characteristic difference between the eastern and western apes has been so faithfully inherited that it is very instructive for us. It is true that there seems to be an exception in the case of a small family of South American apes. The small silky apes (*Acrotrichus* or *Leptailurus*), which include the tamarin (*Midas*) and the brush-monkey (*Jacchus*), have only five molars in each half of the jaw (instead of six), and so seem to be nearer to the eastern apes. But it is found, on closer examination, that they have three pre-molars, like all the western apes, and that only the last molar has been lost. Hence the apparent exception really confirms the above distinction.

Of the other features in which the two groups of apes differ, the structure of the nose is particularly instructive and conspicuous. All the eastern apes have the same type of nose as man—a comparatively narrow partition between the two halves, so that the nostrils run downwards. In some of them the nose protrudes as far as in man, and has the same characteristic structure. We have already alluded to the curious long-nosed apes, which have a long, fleshy-curved nose. Most of the western apes have, it is true, rather flat noses, like, for instance, the white-nosed monkey (Fig. 276); but the nasal partition is thin and narrow in them all. The American apes have a different type of nose. The partition is very broad and thick at the bottom, and the wings of the nostrils are not developed, so that they point outwards instead of downwards. This difference in the form of the nose is so constantly inherited in both groups that the apes of the New World are called "flat-nosed" (*Platyrrhines*), and those of the Old World "narrow-nosed" (*Catarrhines*). The bony passage of the ear (at the bottom of which is the tympanum) is short and wide in all the *Platyrrhines*,

but long and narrow in all the *Catarrhines*; and in man this difference also is significant.

This division of the apes into *Platyrrhines* and *Catarrhines*, on the ground of the above hereditary features, is now generally admitted in zoology, and receives strong support from the geographical distribution of the two groups in the east and west. It follows at once, as regards the phylogeny of the apes, that two divergent lines proceeded from the common stem-form of the ape-order in the early Tertiary period, one of which spread over the Old, the other over the New, World. It is certain that all the *Platyrrhines* come of one stock, and also all the *Catarrhines*; but the former are phylogenetically older, and must be regarded as the stem-group of the latter.

What can we deduce from this with regard to our own genealogy? Man has just the same characters, the same form of dentition, auditory passage, and nose, as all the *Catarrhines*; in this he radically differs from the *Platyrrhines*. We are thus forced to assign him a position among the eastern apes in the order of Primates, or at least place him alongside of them. But it follows that man is a direct blood relative of the apes of the Old World, and can be traced to a common stem-form together with all the *Catarrhines*. In his whole organisation and in his origin man is a true *Catarrhine*; he originated in the Old World from an unknown, extinct group of the eastern apes. The apes of the New World, or the *Platyrrhines*, form a divergent branch of our genealogical tree, and this is only distantly related at its root to the human race. We must assume, of course, that the earliest Eocene apes had the full dentition of the *Platyrrhines*; hence we may regard this stem-group as a special stage (the twenty-sixth) in our ancestry, and deduce from it (as the twenty-seventh stage) the earliest *Catarrhines*.

We have now reduced the circle of our nearest relatives to the small and comparatively scanty group that is represented by the sub-order of the *Catarrhines*, and we are in a position to answer the question of man's place in this sub-order, and say whether we can deduce anything further from this position as to our immediate ancestors. In answering this question the comprehensive and able studies that Huxley gives of

the comparative anatomy of man and the various Catarrhines in his *Man's Place in Nature* are of great assistance to us. It is quite clear from these that the differences between man and the highest Catarrhines (gorilla, chimpanzee, and orang) are in every respect slighter than the corresponding differences between the highest and the lowest Catarrhines (white-nosed monkey, macaco, baboon, etc.). In fact, within the small group of the tail-less anthropoid apes the differences between the various genera are not less than the differences between them and man. This is seen by a glance at the skeletons that Huxley has put together (Figs. 278-282). Whether we take the skull or the vertebral column or the ribs or the fore or hind limbs, or whether we extend the comparison to the muscles, blood-vessels, brain, placenta, etc., we always reach the same result on impartial examination—that man is not more different from the other Catarrhines than the extreme forms of them (for instance, the gorilla and baboon) differ from each other. We may now, therefore, complete the Huxleyan law we have already quoted with the following thesis: "Whatever system of organs we take, a comparison of their modifications in the series of Catarrhines always leads to the same conclusion; the anatomic differences that separate man from the most advanced Catarrhines (orang, gorilla, chimpanzee) are not as great as those that separate the latter from the lowest Catarrhines (white-nosed monkey, macaco, baboon)."

We must, therefore, consider the descent of man from other Catarrhines to be fully proved. Whatever further information on the comparative anatomy and ontogeny of the living Catarrhines we may obtain in the future, it cannot possibly disturb this conclusion. Naturally, our Catarrhine ancestors must have passed through a long series of different forms before the human type was produced. The chief advances that effected this "creation of man," or his differentiation from the nearest related Catarrhines, were: the adoption of the erect posture and the consequent greater differentiation of the fore and hind limbs, the evolution of articulate speech and its organ, the larynx, and the further development of the brain and its function; the soul; sexual selection had a great influence in this, as Darwin showed in his famous work.

With an eye to these advances we can distinguish at least four important stages in our simian ancestry, which represent prominent points in the historical process of the making of man. We may take, after the Lemurs, the earliest and lowest Platyrrhines of South America, with thirty-six teeth, as the twenty-sixth stage of our genealogy; they were developed from the Lemurs by a peculiar modification of the brain, teeth, nose, and fingers. From these Eocene stem-apes were formed the earliest Catarrhines or eastern apes, with the human dentition (thirty-two teeth), by modification of the nose, lengthening of the bony channel of the ear, and the loss of four pre-molars. These oldest stem-forms of the whole Catarrhine group were still thickly coated with hair, and had long tails—baboons (*Cynopithecus*) or tailed apes (*Menacens*, Fig. 376). They lived during the Tertiary period, and are found fossilized in the Miocene. Of the actual tailed apes perhaps the nearest to them are the *Semnopithecus*.

If we take these *Semnopithecus* as the twenty-seventh stage in our ancestry, we may put next to them, as the twenty-eighth, the tail-less anthropoid apes. This name is given to the most advanced and man-like of the existing Catarrhines. They were developed from the other Catarrhines by losing the tail and part of the hair, and by a higher development of the brain, which found expression in the enormous growth of the skull. Of this remarkable family there are only a few genera to-day, and we have already dealt with them (Chapter XV.)—the gibbon (*Hylodactylus*, Fig. 203) and orang (*Sapientia*, Figs. 204, 205) in South-Eastern Asia and the Archipelago; and the chimpanzee (*Anthropithecus*, Figs. 206, 207) and gorilla (*Gorilla*, Fig. 208) in Equatorial Africa.

The great interest that every thoughtful man takes in these nearest relatives of ours has found expression recently in a fairly large literature. The most distinguished of these works for impartial treatment of the question of affinity is Robert Hartmann's little work on *The Anthropoid Apes*. Hartmann divides the primate order into two families: (1) *Primaries* (man and the anthropoid apes); and (2) *Simians* (true apes, Catarrhines and Platyrrhines). Professor Klatsch, of Heidelberg, has advanced a different view in his interesting and richly illustrated work on *The Origin and Development of the Human*

Race. This is a substantial supplement to my *Anthropogeny*, in so far as it gives the chief results of modern research on the early history of man and civilisation. But when Klatech declares the descent of man from the apes to be "irrational, narrow-minded, and false," in the belief that we are thinking of some living species of ape, we must remind him that no competent scientist has ever held so narrow a view. All of us look merely—in the sense of Lamarck and Darwin—to the original unity (admitted by Klatech) of the primate stem. This common descent of all the Primates (men, apes, and lemurs) from one primitive stem-form, from which the most far-reaching conclusions follow for the whole of anthropology and ethnology in admitted fact.

Klatech tells us by myself and all other competent zoologists who accept the theory of evolution in general. He says explicitly (p. 172). "The three anthropoid apes—gorilla, chimpanzee, and orang—seem to be branches from a common root, and this was not far from that of the gibbon and man." That is in the main the opinion that I have maintained (especially against Virchow) in a number of works ever since 1866. The hypothetical common ancestor of all the Primates, which must have lived in the earliest Tertiary period (more probably in the Cretaceous), was called by me *Archiprimus*; Klatech now calls it *Prosimia*. Dubois has proposed the private name of *Prohylobates* for common and much younger stem of the anthropomorpha (man and the anthropoid apes).

three points. None of these can be said to be absolutely the most man-like. The gorilla comes next to man in the structure of the hand and foot, the chimpanzee in the chief features of the skull, the orang in brain development, and the gibbon in the formation of the chest. None of these existing anthropoid apes is among the direct ancestors of our race, they are scattered survivors of an ancient branch of the Catarrhines, from which the human race developed in a particular direction.

Although man is directly connected with

this anthropoid family and originated from it, we may assign an important intermediate form between the *Prohylobates* and him (the twenty-ninth stage in our ancestry), the ape-man (*Pithecanthropus*). I gave this name in the *History of Creation* to the "speechless primitive man" (*Alah*), which were men in the ordinary sense as far as the general structure is concerned (especially in the differentiation of the limbs), but lacked one of the chief human characteristics, articulate speech and the higher intelligence that goes with it, and so had a less developed brain. The phylogenetic hypothesis of the organisation of this "ape-man" which I then advanced was brilliantly confirmed twenty-four years afterwards by the



FIG. 263.—Skull of the fossil ape-man of Java (*Pithecanthropus erectus*), collected by Eugène Dubois.

anthropus erectus by Eugène Dubois (then military surgeon in Java, afterwards professor at Amsterdam). In 1892 he found at Trinil, in the residency of Madiun in Java, in Pliocene deposits, certain remains of a large and very man-like ape (roof of the skull, femur, and teeth), which he described as "an erect ape-man" and a survivor of a "stem-form of man" (Fig. 263). Naturally, the *Pithecanthropus* excited the liveliest interest, as the long-sought transitional form between man and the ape: we seemed to have found "the missing link." There were very interesting scientific discussions of it at the last three International Congresses of Zoology (Leyden, 1895, Cambridge, 1898, and Berlin, 1901). I took an active part in the discussion at

Cambridge, and may refer the reader to the paper I read there on "The Present Position of Our Knowledge of the Origin of Man" (translated by Dr. Gadow with the title of *The Last Link*).

An extensive and valuable literature has grown up in the last ten years on the Pithecanthropus and the pithecoïd theory connected with it. A number of distinguished anthropologists, anatomists, paleontologists, and phylogenists have taken part in the controversy, and made use of the important data furnished by the new science of pre-historic research. Hermann Klintsch has given a good summary of them, with many fine illustrations, in the above-mentioned work. I refer the reader to it as a valuable supplement to the present work, especially as I cannot go any further here into these anthropological and pre-historic questions. I will only repeat that I think he is wrong in the attitude of hostility that he affects to take up with regard to my own views on the descent of man from the apes.

The most powerful opponent of the pithecoïd theory—and the theory of evolution in general—during the last thirty years (until his death in September, 1902) was the famous Berlin anatomist, Rudolf Virchow. In the speeches which he delivered every year at various congresses and meetings on this question, he was never tired of attacking the hated "ape theory." His constant categorical position was, "It is quite certain that man does not descend from the ape or any other animal." This has been repeated incessantly by opponents of the theory, especially theologians and philosophers. In the inaugural speech that he delivered in 1894 at the Anthropological Congress at Vienna, he said that "man might just as well have descended from a sheep or an elephant as from an ape." Absurd expressions like this only show that the famous pathological anatomist, who did so much for medicine in the establishment of cellular pathology, had not the requisite attainments in comparative anatomy and ontogeny, systematic zoology and paleontology, for sound judgment in the province of anthropology. The Strasbourg anatomist, Gustav Schwabe, deserved great praise for having the moral courage to oppose this dogmatic and ungrounded teaching of Virchow, and showing its untenability. The recent admirable works of Schwabe on the Pithecanthropus, the earliest races

of man, and the Neanderthal skull (1897-1901) will supply any candid and judicious reader with the empirical material with which he can convince himself of the baselessness of the erroneous dogmas of Virchow and his clerical friends (J. Ranke, J. Bumüller, etc.).

As the Pithecanthropus walked erect, and his brain (judging from the capacity of his skull, Fig. 383) was midway between the lowest man and the anthropoid apes, we must assume that the next great step in the advance from the Pithecanthropus to man was the further development of human speech and reason.

Comparative philology has recently shown that human speech is polyphyletic in origin; that we must distinguish several (probably many) different primitive tongues that were developed independently. The evolution of language also teaches us (both from its ontogeny in the child and its phylogeny in the race) that human speech proper was only gradually developed after the rest of the body had attained its characteristic form. It is probable that language was not evolved until after the dispersal of the various species and races of men, and this probably took place at the commencement of the Quaternary or Diluvial period. The speechless apes-men or *Alais* certainly existed towards the end of the Tertiary period, during the Pliocene, possibly even the Miocene, period.

The third, and last, stage of our animal ancestry is the true or speaking man (*Homo*), who was gradually evolved from the preceding stage by the advance of animal language into articulate human speech. As to the time and place of this real "creation of man" we can only express tentative opinions. It was probably during the Diluvial period in the hotter zone of the Old World, either on the mainland in tropical Africa or Asia, or on an earlier continent (Lamuria—now sunk below the waves of the Indian Ocean), which stretched from East Africa (Madagascar, Abyssinia) to East Asia (Sunda Islands, Further India). I have given fully in my *History of Creation* (chap. xxviii) the weighty reasons for claiming this descent of man from the anthropoid eastern ape, and shown how we may conceive the spread of the various races from this "Paradise" over the whole earth. I have also dealt fully with the relations of the various races and species of men to each other.

SYNOPSIS OF THE CHIEF SECTIONS OF OUR STEM-HISTORY

First stage: The Protists.

Man's ancestors are unicellular protozoa, originally un-nucleated *Monera* like the *Chromacea*, structureless green particles of *Plasma*; afterwards real nucleated cells (first planuloidous *Protophytas*, like the *Palmetta*, then plasmodiophagous *Prozoa*, like the *Amoeba*).

Second stage: The Blastozoa.

Man's ancestors are round *oponidia* or colonies of *Protozoa*, they consist of a close association of many homogeneous cells, and thus are individuals of the second order. They resemble the round cell-communities of the *Magrophorae* and *Volvocina*, equivalent to the ontogenetic blastula hollow globules, the wall of which consists of a single layer of isolated cells (blastoderm).

Third stage: The Gastraea.

Man's ancestors are *Gastraea*, like the simplest of the actual Metazoa (*Prophyllina*, *Olinthina*, *Hydra*, *Pennatulidina*). Their body consists merely of a primitive gut, the wall of which is made up of the two primary germinal layers.

Fourth stage: The Platodes.

Man's Ancestors have substantially the organization of simple *Platodes* (at first like the cryptozoic *Platodina*, later like the rhadocoele *Turbellaria*). The leaf-shaped bilateral-symmetrical body has only one gut-opening, and develops the first trace of a nervous centre (from the ectoderm in the middle line of the back (Figs. 239, 240).

Fifth stage: The Vermula.

Man's ancestors have substantially the organization of unarticulated *Vermula*, as first *Gavistotricha* (Ichthyodina), afterwards *Frontozia* (*Nemertina*, *Enteropneusta*). Four secondary germinal layers develop, two middle layers arising between the limiting layers (mesoderm). The dorsal ectoderm forms the vertical plate, acrogastric (Fig. 243).

Sixth stage: The Prochordonia.

Man's ancestors have substantially the organization of a simple unarticulated *Chordonia* (*Copepoda* and *Acordia*-larvae). The unsegmented chorda develops between the

dorsal medullary tube and the ventral gut-tube. The simple coelom-pouches divide by a frontal septum into two on each side: the dorsal pouch (epiviscum) forms a mesopleuric; the ventral pouch (hypoviscum) forms a gonad. Head-gut with gill-clefts.

Seventh stage: The Amphibia.

Man's ancestors are skull-less *Vertebrata*, like the *Amphioxus*. The body is a series of metamers, so several of the primitive segments are developed. The head contains in the ventral half the branchial gut, the trunk the hepatic gut. The medullary tube is still simple. No skull, jaws, or limbs.

Eighth stage: The Cyclostoma.

Man's ancestors are jaw-less *Craniozoa* (like the *Myxozoa* and *Petromyzontia*). The number of metamers increases. The fore-end of the medullary tube expands into a vesicle and forms the brain, which soon divides into five cerebral vesicles. In the sides of it appear the three higher sense-organs: nose, eyes, and auditory vesicles. No jaws, limbs, or floating bladder.

Ninth stage: The Ichthyoda.

Man's ancestors are fish-like *Craniozoa*: (1) Primitive fishes (*Selachii*), (2) plated fishes (*Ganoids*); (3) amphibian fishes (*Dipnoetia*), (4) scaled amphibians (*Stegoccephali*). The ancestors of this series develop two pairs of limbs, a pair of fore (breast-fins) and of hind (belly-fins) legs. The gill-arches are formed between the gill-clefts: the first pair forms the maxillary arches (upper and lower jaws). The floating bladder (lung) and pancreas grow out of the gut.

Tenth stage: The Amniotes.

Man's ancestors are *Amniotes* or gill-less *Vertebrata*: (1) Primitive *Amniotes* (*Proreptalia*), (2) *Sauromammalia*, (3) Primitive *Mammalia* (*Monotremata*), (4) *Marsupialia*, (5) *Lemurs* (*Prosimia*), (6) *Western apes* (*Platyrrhini*); (7) *Eastern apes* (*Catarrhini*); at first tailed *Cynopithecina*, then tail-less *Anthropoids*; later speechless *ape-men* (*Aiali*); finally speaking man. The ancestors of these *Amniotes* develop an amnion and allantois, and gradually assume the mammal, and finally the specifically human, form.

CHAPTER XXIV.

EVOLUTION OF THE NERVOUS SYSTEM

THE previous chapters have taught us how the human body as a whole develops from the first simple rudiment, a single layer of cells. The whole human race owes its origin, like the individual man, to a single cell. The unicellular stem-form of the race is reproduced daily in the unicellular embryonic stage of the individual. We have now to consider in detail the evolution of the various parts that make up the human frame. I must, naturally, confine myself to the most general and principal outlines, to make a special study of the evolution of each organ and tissue is both beyond the scope of this work, and probably beyond the anatomic capacity of most of my readers to appreciate. In tracing the evolution of the various organs we shall follow the method that has hitherto guided us, except that we shall now have to consider the ontogeny and phylogeny of the organs together. We have seen, in studying the evolution of the body as a whole, that phylogeny casts a light over the darker paths of ontogeny, and that we should be almost unable to find our way in it without the aid of the former. We shall have the same experience in the study of the organs in detail, and I shall be compelled to give simultaneously their ontogenetic and phylogenetic origin. The more we go into the details of organic development, and the more closely we follow the rise of the various parts, the more we see the inseparable connection of embryology and stem-history. The ontogeny of the organs can only be understood in the light of their phylogeny, just as we found of the embryology of the whole body. Each embryonic form is determined by a corresponding stem-form. This is true of details as well as of the whole.

We will consider first the animal and then the vegetal systems of organs of the body. The first group consists of the psychic and the motor apparatus. To the former belong the skin, the nervous system, and the sense-organs. The motor apparatus is composed of the

passive and the active organs of movement (the skeleton and the muscles). The second or vegetal group consists of the nutritive and the reproductive apparatus. To the nutritive apparatus belong the alimentary canal with all its appendages, the vascular system, and the renal (kidney) system. The reproductive apparatus comprises the different organs of sex (embryonic glands, sexual ducts, and copulative organs).

As we know from previous chapters (XI.-XIII.), the animal systems of organs (the organs of sensation and presentation) develop for the most part out of the outer primary germ-layer, or the cutaneous (skin) layer. On the other hand, the vegetal systems of organs arise for the most part from the inner primary germ-layer, the visceral layer. It is true that this antithesis of the animal and vegetal spheres of the body in man and all the higher animals is by no means rigid; several parts of the animal apparatus (for instance, the greater part of the muscles) are formed from cells that come originally from the ectoderm; and a great part of the vegetative apparatus (for instance, the mouth-cavity and the gonoducts) are composed of cells that come from the ectoderm.

In the more advanced animal body there is so much interlacing and displacement of the various parts that it is often very difficult to indicate the source of them. But, broadly speaking, we may take it as a positive and important fact that in man and the higher animals the chief part of the animal organs comes from the ectoderm, and the greater part of the vegetative organs from the endoderm. It was for this reason that Carl Ernst von Baer called the one the animal and the other the vegetative layer (see p. 26).

The solid foundation of this important thesis is the *gastrula*, the most instructive embryonic form in the animal world, which we still find in the same shape in the most diverse classes of animals. This form points demonstrably to a

common stem-form of all the Metazoa, the *Gastrea*; in this long-extinct stem-form the whole body consisted throughout life of the two primary germinal layers, as is now the case temporarily in the gastrula; in the *Gastrea* the simple cutaneous (skin) layer actually represented all the animal organs and functions, and the simple visceral (gut) layer all the vegetal organs and functions. This is the case with the modern *Gastrea* (Fig. 233); and it is also the case potentially with the *Scyphozoa*.

We shall easily see that the gastrula theory is thus able to throw a good deal of light, both morphologically and physiologically, on some of the chief features of embryonic development. If we take up first the consideration of the chief element in the animal sphere, the psychic apparatus or sensorium and its evolution. This apparatus consists of two very different parts, which seem at first to have very little connection with each other—the outer skin, with all its hairs,

system. The latter comprises the central nervous system (brain and spinal cord), the peripheral, cerebral, and spinal nerves, and the sense-organs. In the fully-formed vertebrate body these two chief elements of the sensorium lie far apart, the skin being external to, and the central nervous system in the very centre of, the body. The one is only connected with the other by a section of the peripheral nervous system and the sense-organs. Nevertheless, as we know from human embryology, the medullary tube is formed from the cutaneous layer. The organs that discharge the most advanced functions of the animal body—the organs of the soul, or of psychic life—develop from the external skin. This is a perfectly natural and necessary process. If we reflect on the historical evolution of the psychic and sensory functions, we are forced to conclude that the cells which accomplish them must originally have been local.

body. Only elementary organs in this superficial position could directly receive the influences of the environment. Afterwards, under the influence of natural selection, the cellular group in the skin which was specifically "sensitive" withdrew into the inner and more protected part of the body, and formed there the foundation of a central nervous organ. As a result of increased differentiation,

the skin and the central nervous system became further and further separated, — and in the end the two were only peripherally connected by the afferent peripheral sensory nerves.

The observations of the comparative anatomist are in complete accord with his view. He tells us that large numbers of the lower animals have no nervous system, though they exercise the functions of sensation and will like the higher animals. In the unicellular *Protozoa*, which do not form germinal layers, there is, of course, neither nervous system nor



FIG. 234.—The human skin, in vertical section (from *Strick*), highly magnified. *a*, horny layer of the epidermis; *b*, vascular layer of the epidermis; *c*, bundle of the cutis; *d*, blood-vessels of cutis; *e*, ducts of the sweat-glands; *f*, *g*, hair-follicles in the cutis; *h*, nerve, passing into a tactile corpuscle above.

skin. But in the second division of the animal kingdom also, the Metazoa, there is at first no nervous system. Its functions are represented by the simple cutaneous layer of the body. The lower Metazoa have inherited from the *Gastrea* (Fig. 30 *c*). We find this in the lowest Zoophytes—the *Gastrea*, *Physalia*, and *Spongia* (Figs. 233-238). The lowest *Cnidaria* (the hydroid polype) also are little superior to the *Gastrea* in structure. Their vegetative functions are accomplished by the simple visceral layer, and their animal functions by the simple cutaneous layer. In these

cases the simple cell-layer of the ectoderm is at once skin, locomotive apparatus, and nervous system.

When we come to the higher Metazoa, in which the sensory functions and their organs are more advanced, we find a division of labour among the ectodermic cells. Groups of sensitive nerve cells separate from the ordinary epidermal cells; they retain a protected

the mesoderm form special neural ganglia there. Even in the Platyodes, especially the *Turbellaria* we find an independent nervous system.

This is the "upper pharyngeal ganglion," or *acrogastron*, situated above the gullet (Fig. 241 p). From this rudimentary structure has been developed the elaborate central nervous system of the higher

In some of the earth-worm, the first rudimentary central nervous system



FIG. 241.—Epidermal cells of a human embryo at two months. (From Kolliker.)

(Fig. 74 a) is a local thickening of the skin-sense layer (*hs*), which afterwards separates altogether from the horny plate. In the earliest Platyodes (*Cryptocella*) and *Vermata* (*Gastrotrocha*) the *acrogastron* remains in the epidermis. But the medullary tube of the Vertebrates originates in the same way. Our embryology has taught us that this first structure of the central nervous system also develops originally from the outer germinal layer.

Let us now examine more closely the evolution of the human skin, with its various appendages, the hairs and glands. This external covering has, physiologically, a double and important part to play. It is, in the first place, the common integument that covers the whole surface of the body, and forms a protective — — — — — for the other organs. As such it also effects a certain exchange

matter between the body and the surrounding atmosphere (exhalation, perspiration). In the second place, it is the

1 origin
part of the temperature of the
the press

of bodies that

The human skin (like that of all the higher animals) is composed of two layers: the outer skin, or *epidermis*, and the inner skin, or *hypodermis*.

The outer skin, or *epidermis*, is a simple ectodermic cell-layer, and contains no blood-vessels (Fig. 284 a, b). It develops

the layer. The underlying skin (or *hypodermis*) consists chiefly of a

outermost parietal stratum of the middle germinal layer, or the *dermo-fibrous layer*. The corium is much thicker than the epidermis. In its deeper strata (the *subcutis*) there are clusters of fat-cells (Fig. 284 b). Its uppermost stratum (the *cutis proper*, or the *papillary stratum*) forms, over almost the whole surface of the body, a number of conical microscopic papillae (something like warts), which push into the overlying epidermis (c). These tactile or sensory papillae contain the finest sensory organs of the skin, the touch corpuscles. Others contain merely end-loops of the blood-vessels that nourish the skin (c, d). The various parts of the corium arise by division of labour from the originally homogeneous cells of the cutis-plate, the outermost lamina of the mesodermic skin-fibre layer (Fig. 145 *hbr.*, Figs. 284 a, b).

In the same way, all the parts and appendages of the epidermis develop by differentiation from the homogeneous cells of this horny plate (Fig. 285). At an early stage the simple cellular layer of this horny plate divides into two. The inner and softer stratum (Fig. 284 b) is known as the *mucous stratum*, the outer and harder (*κ*) as the *horny (corneous) stratum*. This horny layer is being constantly used up and rubbed away at the surface; new layers of cells grow up in their place out of the underlying mucous stratum. At first the epidermis is a simple covering of the surface of the body. Afterwards various appendages develop from it, some internally, others externally. The internal appendages are the cutaneous glands—sweat, fat, etc.

The external appendages are the hairs and nails.

The cutaneous glands are originally merely solid cone-shaped growths of the epidermis, which sink into the underlying corium (Fig. 286 1). Afterwards a canal (2, 3) is formed inside them, either by

is or by the secretion of fluid

Some of the glands, as the sudoriferous, do not ramify (Fig. 286 2). These glands, which secrete perspiration, are very long, and have a spiral coil at the end, but they never ramify, so also the wax-glands of the is glands give out buds and ramify; thus, for instance, the lachrymal glands of the upper eyelid that secrete tears (Fig. 286), and the sebaceous glands which secrete the fat in the skin and generally open into the hair-follicles. Sudoriferous and sebaceous glands are found only in mammals. But we find lachrymal glands in all the three classes of Amniotes—reptiles, birds, and mammals. They are wanting in the lower, aquatic vertebrates.

The mammary glands (Figs. 287 and

They secrete the milk for the feeding of the new-born mammal. In spite of their unusual size, these structures are nothing more than large sebaceous glands in the skin. The milk is formed by the liquefaction of the fatty milk-cells inside the branching mammary-gland tubes (Fig. 287 c), in the same way as the skin-grease or hair-fat, by the solution of fatty cells inside the sebaceous glands. The outlets of the mammary glands enlarge and form sac-like mammary ducts (d), these narrow again (e), and open in the teats or nipples of the breast by sixteen to

The first gland is epidermic, corium and

lobes (Fig. 288). These gradually ramify,

the lobes. Thus is formed the prominent female breast (*mammas*), on the top of which rises the teat or nipple (*mamilla*). The latter is only developed later on, when the mammary gland is fully formed, and this ontogenetic phenomenon is very interesting, because

mammals (the stem-forms of the whole class) have no teats. In them the milk comes out through a flat portion of the ventral skin that is pierced like a sieve. In we still find in the lowest living mammals the viviparous Monotremes of Australia. The young animal licks the

In many of the lower mammals we find a number of milk-glands at different parts of the ventral surface. In the human female there is usually only a pair of glands, at the breast; and it is the same with the apes, bats, elephants, and several other mammals. Some find successive pairs or



FIG. 286.—Proliferatory lachrymal glands from a human embryo of four months. (From Albinus) 1, solid structure, in the shape of a simple solid cone, 2 and 3 more advanced structures, ramifying and hollow buds, 4, solid buds, cellular coat of the hollow buds, 5, structure of the teat which afterwards forms the teat.

glands (or even more) female. Some have pairs of breasts, like pigs and hedgehogs (Fig. 289). This polymastism points back to an older stem-form. We often find many breasts

(Fig. 289 D). Sometimes, moreover, the normal mammary glands are fully developed and can suckle in the male, but the organs without function in the male. We have already (Chapter XI) dealt with this remarkable and interesting instance of atavism.

While the cutaneous glands are inner growths of the epidermis, the appendages

which we call hairs and nails are external local growths in it. The nails (*Ungues*)

which we

Instead of them; the ungulate
The stem-form of the mammals
had claws; we find them in a rudimentary
form even in the salamander. The horny
claws are highly developed in most of the
reptiles (Fig. 264, p. 245), and the mam-
mals have inherited them from the earliest
representatives of this class, the stem-
reptiles (*Tapejaria*). Like the hoofs



FIG. 264.—The female breast (mammula) in vertical section. *c*, mammary glandular tissue, *b*, enlarged milk ducts, *a*, narrower quills, which open into the nipple. (From H. Meyer.)

(*ungulae*) of the Ungulates the nails of apes and men have been evolved from the claws of the older mammals. In the human embryo the first rudiment of the nail is found (between the horny and the mucous stratum of the epidermis) in the fourth month. But their edges do not penetrate through until the end of the sixth month.

The most interesting and important appendages of the epidermis are the hairs; on account of their peculiar composition and origin we must regard them as highly characteristic of the whole mammalian class. It is true that we also find hairs in many of the lower animals,

hairs, like the hairs of plants, are thread-like appendages of the surface, and differ entirely from the hairs of the mammals in the details of their structure and develop-

Embryology of the hair

views as to their phylogeny. On the older view the hairs of the mammals are equivalent or homologous to the feathers of the bird or the horny scales of the reptile. As we deduce all three classes of Amniotes from a common stem-group, we must assume that these Pennsylvanian stem-reptiles had a complete scaly coat, inherited from their Carboniferous ancestors, the mailed amphibians (*Stegocéphala*). The bony scales of their corium were covered with horny scales. In passing from aquatic to terrestrial life the horny scales were further developed, and the bony scales degenerated in most of the reptiles. As regards the bird's feathers, it is certain that they are modifications of the horny scales of their reptilian ancestors. But it is otherwise with the hairs of the mammals. In their case the

has lately been advanced on the basis of very extensive research, by Friedrich Müller, that they evolved from the cutaneous sense-organs of amphibian ancestors by modification of functions; the epidermic structure is very similar in both in its embryonic rudiments. This modern view, which had the support of the greatest expert on the vertebrates, Carl Gegenbaur, can be harmonised with the older theory to an extent, in the sense that both the scales and hairs, were very closely connected originally. Probably the budding of the skin-sense in the *pander* the protection of the

touch subsequently by the cornification of the hairs; many hairs are still sensory organs (tactile hairs on the muzzle and choles of many mammals: pubic hairs).

Thus middle position of the genetic connection of scales and hairs was advanced in my *Systematic Phylogeny of the Vertebrates* (p. 433). It is confirmed by the similar arrangement of the two cutaneous formations. As Müller pointed out, the hairs, as well as the cutaneous sense-organs and the scales, are at first arranged in regular longitudinal series, and they afterwards break into alternate groups. In the embryo of a bear two

inches long, which I owe to the kindness of Herr von Schmerzing (of Arva Vasalla, Hungary), the back is covered with sixteen to twenty alternating longitudinal rows of scaly protuberances (Fig. 289). They are at the same time arranged in regular transverse rows, which converge at an acute angle from both sides towards the middle of the back. The tip of the scale-like wart is turned inwards. Between these larger hard scales (or groups of hairs) we find numbers of rudimentary smaller hairs.

The human embryo is, as a rule, entirely clothed with a thick coat of fine ring the last three or four weeks of gestation. This embryonic woolly coat (*Lanugo*) generally disappears in part during the last weeks of fetal life, but in any case, as a rule, it is lost immediately at birth, and is replaced by the thinner coat of the permanent hair. These permanent hairs grow out of hair-follicles, which are formed from the root-sheaths of the disappearing wool-fibres. The embryonic wool-coat usually, in the case of the human embryo, covers the whole body, with the exception of the palms of the hands and soles of the feet. The parts are always bare, as in the case of apes and of most other mammals. Sometimes the wool-coat of the embryo has a striking effect, by its colour, on the later permanent hair-coat. Hence it happens occasionally, for instance, among our Indo-Germanic races, that children of blond parents seem—to the distress of the latter—to be covered at birth with a dark brown or even a black woolly coat. Not until this has disappeared do we see the permanent hair which the child has inherited. Sometimes the darker coat remains for weeks, and even months, after birth. This remarkable woolly coat of the human embryo is a legacy from the apes, our ancient long-haired ancestors.

It is not less noteworthy that many of the higher apes approach man in the thickness of the hair on various parts of the body. With most of the apes, especially the higher Catarrhines (or narrow-nosed apes), the face is, mostly, or entirely, bare, or at least it has hair no longer or thicker than that of man. In their case, too, the back of the head is usually provided with a thicker growth of hair; this is lacking, however, in the case of the bald-headed chimpanzees (*Pan troglodytes*). The males of many species of apes have a considerable beard on the

cheeks and chin; this sign of the masculine sex has been acquired by sexual selection. Many species of apes have a very thin covering of hair on the breast and the upper side of the limbs—much thinner than on the back or the under side of the limbs. On the other hand, we are often astonished to find tufts of hair on the shoulders, back, and extremities of members of our Indo-Germanic and of the Semitic races. Exceptional hair on the face, as on the whole body, is hereditary in certain families of hairy men. The quantity and the quality of the hair on head and chin are also conspicuously transmitted in families. The extraordinary variations in the total or partial hairy coat of the body, which is so noticeable, not only in compar-



Fig. 291.—Mammary gland of a new-born infant: a original central gland; small black dots (from Lanugo).

different races of men, but also in comparing different families of the same race, can only be explained on the assumption that in man the hairy coat is, on the whole, a rudimentary organ, a useless inheritance from the more thickly-coated apes. In this man resembles the elephant, rhinoceros, hippopotamus, whale, and other mammals of various orders, which have also, almost entirely or for the most part, lost their hairy coats by adaptation.

The particular process of adaptation by which man lost the growth of hair on most parts of his body, and retained or augmented it at some points, was most probably sexual selection. As Darwin luminously showed in his *Descent of Man*, sexual selection has been very active

in this respect. As the male anthropoid apes chose the females with the least hair, and the females favoured the males with the finest growths on chin and head, the general coating of the body gradually degenerated, and the hair of the beard and head was more strongly developed. The growth of hair at other parts of the body (armpit, pubic region) was also probably due to sexual selection. Moreover, changes of climate, or habits, and other adaptations unknown to us, may

poid apes—gorilla, chimpanzee, orang, and several species of gibbons—besides man (Figs. 202, 207). In other species of gibbon the hairs are pointed towards the hand both in the upper and lower arm, as in the rest of the mammals. We can easily explain this remarkable peculiarity of the anthropoids and man on the theory that our common ancestors were accustomed (as the anthropoid apes are to-day) to place their hands over their heads, or across a branch above their



FIG. 216.—Embryos of a bear (*Ursus arctos*), twice natural size. A seen from ventral side, B from the left.

have assisted the disappearance of the hairy coat.

The fact that our coat of hair is inherited directly from the anthropoid apes is proved in an interesting way, according to Darwin, by the direction of the rudimentary hairs on our arms, which cannot be explained in any other way. Both on the upper and the lower part of the arm they point towards the elbow. Here they meet at an obtuse angle. This curious arrangement is found only in the anthro-

poids, during rain. In this position, the fact that the hairs point downwards helps the rain to run off. Thus the direction of the hair on the lower part of our arm reminds us to-day of that useful custom of our anthropoid ancestors.

The nervous system in man and all the other Vertebrates is, when fully formed, an extremely complex apparatus, that we may compare, in anatomic structure and physiological function, with an extensive telegraphic system. The chief station of

the system is the central marrow or central nervous system, the innumerable ganglionic cells or *neurones* (Fig. 9) of which are connected by branching processes with each other and with numbers of very fine conducting wires. The latter are the peripheral and ubiquitous nerve-fibres; with their terminal apparatus, the sense-organs, etc., they constitute the conducting marrow or peripheral nervous system. Some of them—the sensory nerve-fibres—conduct the impressions from the skin and other sense-organs to the central marrow; others—the motor nerve-fibres—convey the commands of the will to the muscles.

The central nervous system or central marrow (*medulla centralis*) is the real organ of psychic action in the narrower sense. However we conceive the intimate connection of this organ and its functions,

which we call sensation, we are inseparably dependent development of the man and all the higher

gives us most important information regarding the nature of the "soul," it should be full of interest. If the central marrow develops in just the same way in the human embryo as in the embryo of the other mammals, the evolution of the human psychic organ from the central

them from the lower vertebrates, must be beyond question. No one can doubt the momentous bearing of these embryonic phenomena.

In order to understand them fully we must first say a word or two of the general and the anatomic composition of the mature human central marrow. Like the central nervous system of all other Craniotes, it consists of two parts, the head-marrow or brain (*medulla capitis* or *encephalon*) and the spinal-marrow (*medulla spinalis* or *notomyelon*). The one is enclosed in the bony skull, the other in the bony vertebral column. Twelve pairs of cerebral nerves proceed from the brain, spinal nerves

rest of the body (Fig. 171). On an anatomic investigation the spinal marrow is found to be a cylindrical cord, not only both in the region of the last vertebra and the region of the loins (at

the first lumbar vertebra) below (Fig. 291). At the cervical bulb the strong nerves of the upper limbs, and at the lumbar bulb those of the lower limbs, proceed from the spinal cord. Above, the latter passes into the brain through the medulla oblongata (Fig. 291, *mo*). The spinal cord needs to be a thick mass of nervous matter, but it has a narrow canal at its axis, which passes into the further



FIG. 291.

FIG. 292.

FIG. 291.—Human embryo, three months old, from the dorsal side. Brain and spinal cord exposed. (From A. Müller.) A cerebral hemisphere (dorsal brain), or corpus quadrigemum; medulla (head brain), under B, the oblongata (after brain).

FIG. 292.—Central marrow of a human embryo, four months old, natural size, from the back. (From A. Müller.) A large hemisphere, b cerebellum, c medulla oblongata, underneath is the spinal cord.

cerebral ventricles above, and is filled, like these, with a clear fluid.

The brain is a large, complex structure, occupying the greater part of the skull, of an elaborate structure. On general examination it divides into two parts, the cerebrum and cerebellum. The cerebrum lies in front and above, and has the familiar characteristic convolutions and furrows on its surface (Figs. 292, 293). The upper side it is divided by a deep longitudinal fissure into two halves,

cerebral hemispheres; these are connected by the *corpus callosum*. The large cerebrum is separated from the small cerebellum by a deep transverse furrow. The latter lies behind and below, and has also numbers of furrows, but much finer and more regular, with convolutions between, at its surface. The cerebellum also is divided by a longitudinal fissure into two halves, the "small hemispheres"; these are connected by a worm-shaped piece, the *vermis cerebelli*, above, and by the broad *pons Varolii* below (Fig. 292 f').

But comparative anatomy and ontogeny teach us that in man and all the other




FIG. 292.—The human brain, seen from below (From *N. Meyer*). Above, (in front) is the cerebrum with its sulci and branching furrows. Below (behind) the cerebellum with its narrow parallel furrows. The human numbers indicate the roots of the twelve pairs of cerebral nerves as a series towards the rear.

Crinates the brain is at first composed, not of these two, but of three, and afterwards five, consecutive parts. These are found in just the same form—as five consecutive vesicles—in the embryo of all the Crinates, from the Cyclostoma and fishes to man. But, however much they agree in their rudimentary condition, they differ considerably afterwards. In man and the higher mammals the first of these ventricles, the cerebrum, grows so much that in its mature condition it is by far the largest and heaviest part of the brain. To it belong not only the large hemispheres, but also the corpus callosum that unites them, the olfactory lobes,

from which the olfactory nerves start, and most of the structures that are found at the roof and bottom of the large lateral ventricles inside the two hemispheres, such as the *corpora striata*. On the other hand, the *optic thalami*, which lie between the latter, belong to the second division, which develops from the "intermediate brain", to the same extent as the single third cerebral ventricle and the structures that are known as the *corpora geniculata*, the *Infundibulum*, and the pineal gland. Behind these parts we find, between the cerebrum and cerebellum, a small ganglion composed of two prominences, which is called the *corpus quadrigeminum* on account of a superficial transverse fissure cutting across (Fig. 290 w, 291 v). Although this quadrigeminum is very insignificant in man and the higher mammals, it forms a special third section, greatly developed in the lower vertebrates, the "middle brain." The fourth section is the "hind-brain" or little brain (cerebellum) in the narrower sense, with the single median part, the *vermis*, and the pair of lateral parts, the "small hemispheres." (Fig. 291 c). Finally, we leave the fifth and last section, the *medulla oblongata* (Fig. 291 me), which contains the single fourth cerebral cavity and the contiguous parts (pyramids, olivary bodies, corpora restiformia). The medulla oblongata passes straight into the medulla spinalis (spinal cord). The narrow central canal of the spinal cord continues above into the quadrangular fourth cerebral cavity of the medulla oblongata, the floor of which is the quadrangular depression. From here a narrow duct, called "the aqueduct of Sylvius," passes through the corpus quadrigeminum to the third cerebral ventricle, which lies between the two optic thalami; and this in turn is connected with the pair of lateral ventricles which lie to the right and left in the large hemispheres. Thus all the cavities of the central marrow are directly interconnected. All these parts of the brain have an infinitely complex structure in detail, but we cannot go into this. Although it is much more elaborate in man and the higher Vertebrates than in the lower classes, it develops in them all from the same rudimentary structure, the five simple cerebral vesicles of the embryonic brain.

But before we consider the development of the complicated structure of the brain from this simple series of vesicles, let

us glance for a moment at the lower unicells, which have no brain. Even in the skull-less vertebrate, the Amphioxus, we find no independent brain, as we have seen. The whole central marrow is merely a simple cylindrical cord which runs the length of the body, and ends equally simply at both extremities—a plain medullary tube. All that we can discover is a small vesicular bulb at the forward part of the tube, a degenerate rudiment of a primitive brain. We meet the same simple medullary tube in the first structure of the ascidian larva, in the same characteristic position, above the chorda. On closer examination we find here also a small vesicular swelling

at the fore end of the tube, the first trace of a differentiation of it into brain and spinal cord. It is probable that this differentiation was more advanced in the extinct Pterobranchs, and the brain-bulb more pronounced (Figs. 28-303). The brain is phylogenetically older than the spinal cord, as the trunk was not developed until after the head. If we consider the undeniable affinity of the Ascidaria to the Vermalia, and remember that we can trace all the Chordonia to lower Vermalia, it seems probable that the simple central marrow of the former is equivalent to the simple nervous ganglion, which lies above the gullet in the lower worms, and has long been known as the "upper pharyngeal ganglion" (*ganglion pharyngeum superior*); it would be better to call it the primitive or vertical brain (*acroganglion*).

Probably this upper pharyngeal ganglion of the lower worms is the structure from which the complex central marrow of the higher animals has been evolved. The medullary tube of the Chordonia has been formed by the lengthening of the vertical brain on the dorsal side. In all the other animals the central nervous system has been developed in a totally different way from the upper pharyngeal ganglion; in the Articulates, especially, a pharyngeal ring, with ventral marrow, has been added. The Molluscs also have a pharyngeal ring, but it is not found in the Vertebrates. In these the central

marrow has been prolonged down the dorsal side, in the Articulates down the ventral side. This fact proves of itself that there is no direct relationship between the Vertebrates and the Articulates. The unfortunate attempts to derive the dorsal marrow of the former from the ventral marrow of the latter have totally failed (cf. p. 219).

When we examine the embryology of the human nervous system, we must start from the important fact, which we have already seen, that the first structure of it in man and all the higher Vertebrates is the simple medullary tube, and that this separates from the outer germinal layer in the middle line of the so-called



FIG. 105.—The human brain, seen from the left. (From *H. Meyer*.) The furrows of the cerebrum are indicated by black, and those of the cerebellum by fine lines. Under the latter we can see the medulla oblongata. *f*, *f*, frontal convolution; *C*, central sulcus; *S*, fissure of Sylvius; *T*, temporal furrow; *P*, parietal lobes; *A*, angular gyrus; *B*, parieto-occipital fissure.

embryonic shield. As the reader will remember, the straight medullary furrow first appears in the middle of the sandal-shaped embryonic shield. At each side of it the parallel borders curve over in the form of dorsal or medullary swellings. These bend together with their free borders, and thus form the closed medullary tube (Figs. 133-137). At first this tube lies directly underneath the horny plate; but it afterwards travels forwards, the upper edges of the provertebral plates growing together between the horny plate and the tube, joining above the latter, and forming a completely closed canal. As Gegenbaur very properly observes, "this gradual imbedding in the

inner part of the body is a process acquired with the progressive differentiation and the higher potentiality that this secures; by this process the organ of greater value to the organism is buried within the frame" (Cf. Figs. 143-146).

In the *Cyclostomes*—a stage above the *Acrania*—the fore end of the cylindrical medullary tube begins early to expand into a pear-shaped vesicle; this is the first outline of an independent brain. In this way the central marrow of the Vertebrates divides clearly into its two chief sections, brain and spinal cord. The simple vesicular form of the brain, which persists for some time in the *Cyclostomes*, is found also at first in all the higher Vertebrates (Fig. 153 A). But in these it soon passes away, the one vesicle being divided into several successive parts by transverse constrictions. There are first two of these constrictions,



FIG. 154. FIG. 155. FIG. 156.

Fig. 154-156.—Central marrow of the human embryo from the neurula week. *a*, such long. (From *Reichert*). Fig. 154, back view of the whole neurula; *b*, brain and spinal cord exposed. Fig. 155, the brain with the uppermost part of the cord, from the left. Fig. 156, the brain from above. *r*, fore brain, *m*, intermediate brain, *h*, hind brain. A hind brain, *h*, after brain.

dividing the brain into three consecutive vesicles (fore brain, middle brain, and hind brain, Fig. 154 B, C, D). Then the first and third are sub-divided by fresh constrictions, and thus we get five successive sections (Fig. 155).

In all the *Cranioles*, from the *Cyclostomes* up to man, the same parts develop from these five original cerebral vesicles, though in very different ways. The first vesicle, the fore brain (Fig. 155 B), forms by far the largest part of the cerebrum—namely, the large hemispheres, the olfactory lobes, the corpora striata, the callosum, and the fornix. From the second vesicle, the intermediate brain (*m*), originate especially the optic thalami, the other parts that surround the third cerebral ventricle, and the infundibulum and pineal gland. The third vesicle, the middle brain (*m*), produces the corpora quadrigemina and

the aqueduct of Sylvius. From the fourth vesicle, the hind brain (*h*), develops the greater part of the cerebellum—namely, the vermis and the two small hemispheres. Finally, the fifth vesicle, the after brain (*a*), forms the medulla oblongata, with the quadrangular pit (the floor of the fourth ventricle), the pyramids, olivary bodies, etc.

We must certainly regard it as a comparative-anatomical and ontogenetic fact of the greatest significance that in all the *Cranioles*, from the lowest *Cyclostomes* and fishes up to the apex and man, the brain develops in just the same way in the embryo. The first rudiment of it is always a simple vesicular enlargement of the fore end of the medullary tube. In every case, first three, then five, vesicles develop from this bulb, and the permanent brain with all its complex anatomic structures, of so great a variety in the various classes of Vertebrates, is formed from the five primitive vesicles. When we compare the mature brain of a fish, an amphibian, a reptile, a bird, and a mammal, it seems incredible that we can trace the various parts of these organs, that differ so much internally and externally, to common types. Yet all these different *Craniole* brains have started with the same rudimentary structure. To continue ourselves of this we have only to compare the corresponding stages of development of the embryos of these different animals.

This comparison is extremely instructive. If we extend it through the whole series of the *Cranioles*, we soon discover this interesting fact: In the *Cyclostomes* (the *Myxinoidea* and *Petromyzontes*), which we have recognized as the lowest and earliest *Cranioles*, the whole brain remains throughout life at a very low stage, which is very brief and passing in the embryos of the higher *Cranioles*; they retain the five original sections of the brain unchanged. In the fishes we find an essential and considerable modification of the five vesicles; it is clearly the brain of the *Salichii* in the first place, and subsequently the brain of the *Canoids*, from which the brain of the rest of the fishes on the one hand and of the *Dipnoids* and *Amphibia*, and through these of the higher Vertebrates, on the other hand, must be derived. In the fishes and *Amphibia* (Fig. 300) there is a preponderant development of the middle brain, and also the after brain, the first, second, and

fourth sections remaining very primitive. It is just the reverse in the higher Vertebrates, in which the first and third sections, the cerebrum and cerebellum, are exceptionally developed; while the




FIG. 292.—Head of a chick embryo (shaded lightly from the back, magnified forty times. (From *Atababur*) the anterior wall of the fore brain, the optic vesicles, and middle brain, the hind brain, the after brain, the brain (from below), the vestibular sacs, the primitive segment, the spinal cord.

middle brain and after brain remain small. The corpora quadrigemina are mostly covered by the cerebrum, and the oblongata by the cerebellum. But we find a number of stages of development within the higher Vertebrates themselves. From the Amphibia upwards the brain (and with it the psychic life) develops in two different directions; one of these is followed by the reptiles and birds, and the other by the mammals. The development of the first section, the fore brain, is particularly characteristic of the mammals. It is only in them that the cerebrum becomes so large as to cover all the other parts of the brain (Figs. 293, 301-304).

There are also notable variations in the relative position of the cerebral vesicles. In the lower Craniotes they lie originally almost in the same plane. When we examine the brain laterally, we can cut through all five vesicles with a straight line. But in the Amniotes there is a considerable curve in the brain along with the bending of the head and neck; the whole of the upper dorsal surface of the

brain develops much more than the under ventral surface. This causes a curve, so that the parts come to lie as follows: The fore brain is right in front and below, the intermediate brain a little higher, and the middle brain highest of all; the hind brain lies a little lower, and the after brain lower still. We find this only in the Amniotes—the reptiles, birds, and mammals.

Thus, while the brain of the mammals agrees a good deal in general growth with that of the birds and reptiles, there are some striking differences between the two. In the Sauropsids (birds and reptiles) the middle brain and the middle part of the hind brain are well developed. In the mammals these parts do not grow, and the fore-brain develops so much that it overlies the other vesicles. As it continues to grow towards the rear, it at last covers the whole of the rest of the brain, and also encloses the middle parts from



FIG. 293.—Brain of three craniota embryos in vertical section. A of a shark (*Heterobranchius*), B of a lamprey (*Lampetra*), C of a gold (*Carassius*). a fore brain, b intermediate brain, c middle brain, d hind brain, e after brain, f primitive ducts. (From Gegenbaur.)

FIG. 294.—Brain of a shark (*Sphyrna*) back view. a cerebellum, b olfactory lobes, which send the large olfactory nerves to the nasal capsule (a & b intermediate brain, c middle brain, d hind brain, e after brain, f primitive ducts). (From Gegenbaur.)

the sides (Figs. 301-303). This process is of great importance, because the fore brain is the organ of the higher psychic life, and in it those functions of the nerve-cells are discharged which we sum up in

the word "soul." The highest achievements of the animal body—the wonderful manifestations of consciousness and the complex molecular processes of thought—have their seat in the fore brain. We can remove the large hemispheres, piece by




FIG. 30a.—Brain and spinal cord of the frog. *a* fore brain, *b* from the ventral pole, *c* oblongata, *d* from the (12) fore brain, *e* infundibulum at the base of the submedian brain, *f* middle brain, *g* hind brain, *h* quadrangular pit in the after brain, *i* spinal cord (very short in the frog), *j* roots of the spinal nerves, *k* terminal filum of the spinal cord [From Gegenbaur.]



FIG. 30b.—Brain of an ox-embryo, two inches in length. [From Alcock, *embryos*, magnified three times.] Left view, the lateral wall of the left hemisphere has been removed. *a* corpora striata, *b* Mammillary body, *c* Armenian horn, *d* middle brain, *e* cerebellum, *f* roof of the fourth ventricle, *g* pons Varoli, *h* medulla oblongata.

piece, from the mammal without killing it, and we then see how the higher functions of consciousness, thought, will, and sensation, are gradually destroyed, and in the end completely extinguished. If the animal is fed artificially, it may be

kept alive for a long time, as the destruction of the psychic organs by no means involves the extinction of the faculties of digestion, respiration, circulation, urination—in a word, the vegetative functions. It is only conscious sensation, voluntary movement, thought, and the combination of various higher psychic functions that are affected.

The fore brain, the organ of these functions, only attains this high level of development in the more advanced Placentals, and thus we have the simple explanation of the intellectual superiority of the higher mammals. The soul of most of the lower Placentals is not much above that of the reptiles, but among the higher Placentals we find an uninterrupted gradation of mental power up to the apes and man. In harmony with this we find an astonishing variation in the



FIG. 30c.—Brain of a human embryo, twelve weeks old. [From Alcock, *embryos*, natural size.] Seen from behind and above. *a* fore brain, *b* corpus quadrigenum (middle brain), *c* anterior median line, *d* cerebellum, *e* fourth ventricle, *f* medulla oblongata.

degree of development of their fore brain not only qualitatively, but also quantitatively. The mass and weight of the brain are much greater in modern mammals, and the differentiation of its various parts more important, than in their extinct Tertiary ancestors. This can be shown paleontologically in any particular order. The brains of the living ungulates are (relatively to the size of the body) four to six times (in the highest groups even eight times) as large as those of their earlier Tertiary ancestors, the well-preserved skulls of which enable us to determine the size and weight of the brain.

In the lower mammals the surface of the cerebral hemispheres is quite smooth and level, as in the rabbit (Fig. 30d). Moreover, the fore brain remains so small that it does not cover the middle brain. At a stage higher the middle

peripheral nerves grow out of the medullary tube (Fig. 171), and are, like it, products of the skin-sense layer.

The complete agreement in the structure and development of the psychic organs which we find between man and the highest mammals, and which can only be explained by their common origin, is of profound importance in the monistic psychology. This is only seen in its full light when we compare these morphological facts with the corresponding physiological phenomena, and remember that every psychic action requires the complete and normal condition of the correlative brain structure for its full and normal exercise. The very complex molecular movements inside the neural cells, which we describe comprehensively as "the life of the soul," can no more exist in the vertebrate, and therefore in man, without their organs than the circulation

without the heart and blood. And as the central marrow develops in man from the same medullary tube as that of the other vertebrates, and as man shares the characteristic structure of his cerebrum (the organ of thought) with the anthropoid apes, his psychic life also must have the same origin as theirs.

If we appreciate the full weight of these morphological and physiological facts, and put a proper phylogenetic interpretation on the observations of embryology, we see that the older idea of the personal immortality of the human soul is scientifically untenable. Death puts an end, in man as in any other vertebrate, to the physiological function of the cerebral neurons, the countless microscopic ganglionic cells, the collective activity of which is known as "the soul." I have shown this fully in the eleventh chapter of my *Riddle of the Universe*.

CHAPTER XXV.

EVOLUTION OF THE SENSE-ORGANS

THE sense-organs are indubitably among the most important and interesting parts of the human body; they are the organs by means of which we obtain our knowledge of objects in the surrounding world. *Nihil est in intellectu quod non prius fuerit in sensu*. They are the first sources of the life of the soul. There is no other part of the body in which we discover such elaborate anatomical structures, co-operating with a definite purpose; and there is no other organ in which the wonderful and purposive structure seems so clearly to compel us to admit a Creator and a preconceived plan. Hence we find special efforts made by dualists to draw our attention here to the "wisdom of the Creator" and the design visible in his works. As a matter of fact, you will discover, on mature reflection, that on this theory the Creator is at bottom only playing the part of a clever mechanic or watch-maker; all these familiar teleological ideas of Creator and creation are

based, in the long run, on a similar childlike anthropomorphism.

However, we must grant that at the first glance the teleological theory seems to give the simplest and most satisfactory explanation of these purposive structures. If we merely examine the structure and functions of the most advanced sense-organs, it seems impossible to explain them without postulating a creative act. Yet evolution shows us quite clearly that this popular idea is totally wrong. With its assistance we discover that the purposive and remarkable sense-organs were developed, like all other organs, without any preconceived design—developed by the same mechanical process of natural selection, the same constant correlation of adaptation and heredity, by which the other purposive structures in the animal frame were slowly and gradually brought forth in the struggle for life.

Like most other Vertebrates, man has six sensory organs, which serve for eight

different classes of sensations. The skin serves for sensations of pressure and temperature. This is the oldest, lowest, and vaguest of the sense-organs; it is distributed over the surface of the body. The other sensory activities are localised. The sexual sense is bound up with the skin of the external sexual organs, the sense of taste with the mucous lining of the mouth (tongue and palate), and the sense of smell with the mucous lining of the nasal cavity. For the two most advanced and most highly differentiated sensory functions there are special and very elaborate mechanical structures—the eye for the sense of sight, and the ear for the sense of hearing and space (equilibrium).

Comparative anatomy and physiology teach us that there are no differentiated sense-organs in the lower animals; all their sensations are received by the surface of the skin. The undifferentiated skin-layer or ectoderm of the *Gastrea* is the simple stratum of cells from which the differentiated sense-organs of all the Metazoa (including the Vertebrates) have been evolved. Starting from the assumption that necessarily only the superficial parts of the body, which are in direct touch with the outer world, could be concerned in the origin of sensations, we can see at once that the sense-organs also must have arisen there. This is really the case. The chief part of all the sense-organs originates from the plain-sense layer, partly directly from the horny plate, partly from the brain, the foremost part of the medullary tube, after it has separated from the horny plate. If we compare the embryonic development of the various sense-organs, we see that they all make their appearance in the simplest conceivable form; the wonderful contrivances that make the higher sense-organs among the most remarkable and elaborate structures in the body develop only gradually. In the phylogenetic explanation of them comparative anatomy and ontogeny achieve their greatest triumphs. But at first all the sense-organs are merely parts of the skin in which sensory nerves expand. These nerves themselves were originally of a homogeneous character. The different functions or specific energies of the differentiated sense-nerves were only gradually developed by division of labour. At the same time, their simple terminal expansions in the skin were converted

The great instructiveness of these historical facts in connection with the life of the soul is not difficult to see. The whole philosophy of the future will be transformed as soon as psychology takes cognisance of these genetic phenomena and makes them the basis of its speculations. When we examine impartially the manuals of psychology that have been published by the most distinguished speculative philosophers and are still widely distributed, we are astonished at the naïveté with which the authors raise their airy metaphysical speculations, regardless of the momentous embryological facts that completely refute them. Yet the science of evolution, in conjunction with the great advance of the comparative anatomy and physiology of the sense-organs, provides the one sound empirical basis of a natural psychology.




FIG. 105.—Head of a shark (*Sphyrna*) from the ventral side. *a*, mouth; *b*, secondary pits; *c*, nasal grooves; *d*, nasal fold in natural position; *e*, nasal fold shown expanded. (The dots are openings of the olfactory canals.) (From Giglioli.)

In respect of the terminal expansions of the sensory nerves, we can distribute the human sense-organs in three groups, which correspond to three stages of development. The first group comprises those organs the nerves of which spread out quite simply in the free surface of the skin itself (organs of the sense of pressure, warmth, and sex). In the second group the nerves spread out in the mucous coat of cavities which are at first depressions in or invaginations of the skin (organs of the sense of smell and taste). The third group is formed of the very elaborate organs, the nerves of which spread out in an internal vesicle, separated from the skin (organs of the sense of sight, hearing, and space).

There is little to be said of the development of the lower sense-organs. We

have already considered (p. 268) the organ of touch and temperature in the skin. I need only add that in the course of man and all the higher Vertebrates countless microscopic sense-organs develop, but the precise relation of these to the sensations of pressure or resistance, of warmth and cold, has not yet been explained. Organs of this kind, in or on which sensory cutaneous nerves terminate, are the "tactile corpuscles" (or the

canal to which these parts belong (Chapter XXVII.) I will only point out for the present that the mucous coat of the tongue and palate, in which the gustatory nerve ends, originates from a part of the outer skin. As we have seen, the whole of the mouth-cavity is formed, not as a part of the gut-tube proper, but as a pit-like fold in the outer skin (p. 139). Its mucous lining is therefore formed, not from the visceral, but from the cutaneous



FIGS. 266 and 267.—Head of a chick embryo, three days old. 266 from above, 267 from the right. *a*, rudimentary eye (olfactory pit); *b*, rudimentary eye (top); *c*, rudimentary ear (auditory pit); *d*, ear, brain, *e*, eye-socket, a process of upper jaw, *f*, process of lower jaw of the first gill-arch.

FIG. 268.—Head of a chick embryo, four days old, from below. *a*, rudimentary eye; *b*, upper-jaw process of the first gill-arch; *c*, lower-jaw process of same; *d*, second gill-arch; *e*, pharyngeal pouch of eye; *f*, gullet.

FIGS. 269 and 270.—Heads of chick embryos: 269 from the end of the fourth, 270 from the beginning of the fifth week. Letters as in Fig. 266, except *h*, inner, *i*, outer, nasal process of nasal furrow, *j*, frontal process, *k*, mouth. (From Haeckel.) FIGS. 266–270 are magnified to the same extent.

Pacinian corpuscles) and end-bulbs. We find similar corpuscles in the organs of the sexual sense; the male penis and the female clitoris; they are processes of the skin, the development of which we will consider later (together with the rest of the sexual parts, Chapter XXIX.). The evolution of the organ of taste, the tongue and palate, will also be treated later, together with that of the alimentary

layer, and the taste-cells at the surface of the tongue and palate are not products of the gut-fibre layer, but of the skin-mucous layer.

This applies also to the mucous lining of the olfactory organ, the nose. However, the development of this organ is much more interesting. Although the nose seems superficially to be simple and single, it really consists, in man and all

other Gnathostomes, of two completely separated halves, the right and left cavities. They are divided by a vertical partition, so that the right nostril leads into the right cavity alone and the left nostril into the left cavity. They open internally (and separately) by the posterior nasal apertures into the pharynx, so that we can get direct into the gullet through the nasal passages without touching the mouth. This is the way the air usually passes in respiration; the mouth being closed, it goes through the nose into the gullet, and through the larynx and bronchial tubes into the lungs. The nasal cavities are separated from the mouth by the horizontal bony palate, to which is attached behind (as a dependent process) the soft palate with the uvula. In the upper and hinder parts of the nasal cavities the olfactory nerve, the first pair of cerebral nerves, expands in the mucous coat which clothes them. The terminal branches of it spread partly over the septum (partition), partly on the side-walls of the internal cavities, to which are attached the turbinated bones. These bones are much more developed in many of the higher mammals than in man, but there are three of them in all mammals. The sensation of smell arises by the passage of a current of air containing odorous matter over the mucous lining of the cavities, and stimulating the olfactory cells of the nerve-endings.

Man has all the features which distinguish the olfactory organ of the mammals from that of the lower Vertebrates. In all essential points the human nose entirely resembles that of the Catarrhine apes, some of which have quite a human external nose (compare the face of the long-nosed apes). However, the first structure of the olfactory organ in the human embryo gives an indication of the future simple proportions of our Catarrhine nose. It has the form in which we find it permanently in the fishes—a couple of simple depressions in the skin at the outer surface of the head. We find these blind olfactory pits in all the fishes; sometimes they lie near the eyes, sometimes more forward at the point of the snout, sometimes lower down, near the mouth (Fig. 449).

This first rudimentary structure of the double nose is the same in all the Gnathostomes; it has no connection with the primitive mouth. But even in a section

begins to make its appearance, a furrow in the surface of the skin running from each side of the nasal pit to the nearest corner of the mouth. This furrow, the nasal groove or furrow (Fig. 305 *r*), is very important. In many of the sharks, such as the *Squalus*, a special process of the frontal skin, the nasal fold or internal nasal process, is formed internally over the groove (*s*, *s'*). In contrast to this the outer edge of the furrow rises in an "external nasal process." As the two processes meet and coalesce over the



FIG. 31.—Frontal section of the mouth and throat of a human embryo, back half-inch long. (Lentened by Wilhelm His. The vertical section (in the frontal plane, from left to right) is so constructed that we see the nasal pits in the upper third of the figure and the eyes at the sides. In the middle third the primitive gullet with the gill-arches (gill-arches as sections), in the lower third the pharynx and the broadened trachea and the rudimentary lungs.

nasal groove in the Dipneusts and Amphibia, it is converted into a canal, the nasal canal. Henceforth we can penetrate from the external pits through the nasal canals direct into the mouth, which has been formed quite independently. In the Dipneusts and the lower Amphibia the internal aperture of the nasal canals lies in front (behind the lips); in the higher Amphibia it is right behind. Finally, in the three higher classes of Vertebrates the primary mouth-cavity is

palate-roof into two distinct cavities—the upper (secondary) nasal cavity and the lower (secondary) mouth-cavity. The nasal cavity in turn is divided by the construction of the vertical septum into two halves—right and left.

Comparative anatomy shows us to-day, in the series of the double-nosed Vertebrates, from the fishes up to man, all the different stages in the development of the nose, which the advanced olfactory organ of the higher mammals has passed through at various periods in the course of its phylogeny. It first appears in the embryo of man and the higher Vertebrates, in which the double fish-nose persists throughout life. At an early stage, before there is any trace of the characteristic human face, a pair of small pits are formed in the head over the original mouth-cavity; these were first discovered by Baer, and rightly called the "olfactory pits" (Figs. 306 *m*, 307 *n*).



FIG. 311.—Diagrammatic section of the mouth-nose cavity. While the palato-pharynx (*p*) divides the original mouth-cavity into the lower secondary mouth (*m*) and the upper nasal cavity, the latter in turn is divided by the vertical partition (*v*) into two halves (*n*, *n*). (From Gegenbaur.)

These primitive nasal pits are quite separate from the rudimentary mouth, which also originates as a pit-like depression in the skin, in front of the blind fore end of the gut. Both the pair of nasal pits and the single mouth-pit (Fig. 310 *m*) are clothed with the horny plate. The original separation of the former from the latter is, however, presently abolished, a process forming above the mouth-pit—the "frontal process" (Fig. 309 *n*). Its outer edge rises to the right and left in the shape of two lateral processes; these are the inner nasal processes or folds (*n*). Opposite to these a parallel ridge is formed on either side between the eye and the nasal pit; these are the outer nasal processes (*sn*). Thus between the inner and outer nasal processes a groove-like depression is formed on either side, which leads from the nasal

pit towards the mouth-pit (*m*); this groove is, as the reader will guess, the same nasal furrow or groove that we have already seen in the shark (Fig. 309 *r*). As the parallel edges of the inner and outer nasal processes bend towards each other and join above the nasal groove, this is converted into a tube, the primitive nasal canal. Hence the nose of man and all the other Amniotes consists at this embryonic stage of a couple of narrow tubes, the nasal canals, which lead from the outer surface of the forehead into the rudimentary mouth. This transitional condition resembles that in which we find the nose permanently in the Dipnoans and Amphibia.

A cone-shaped structure, which grows from below towards the lower end of the two nasal processes and joins with them, plays an important part in the construction of the open nasal groove into the closed canal. This is the upper-jaw process (Fig. 306-310 *o*). Below the mouth-pit are the gill-arches, which are separated by the gill-clefts. The first of these gill-arches, and the most important for our purpose, which we may call the maxillary (jaw) arch, forms the skeleton of the jaws. Above at its basis a small process grows out of this first gill-arch; this is the upper-jaw process. The first gill-arch itself develops a cartilage at one of its inner sides, the "Meckel cartilage" (named after its discoverer), on the outer surface of which the lower jaw is formed (Fig. 306-310 *u*). The upper-jaw process forms the chief part of the skeleton of that jaw, the palate bone, and the pterygoid bone. On its outer side is afterwards formed the upper-jaw bone, in the narrower sense, while the middle part of the skeleton of the upper jaw, the intermaxillary, develops from the foremost part of the frontal process.

The two upper-jaw processes are of great importance in the further development of the face. From them is formed, growing into the primitive mouth-cavity, the important horizontal partition (the palate) that divides the former into two distinct cavities. The upper cavity, into which the nasal canals open, now develops into the nasal cavity, the air-passage and the organ of smell. The lower cavity forms the permanent secondary mouth (Fig. 312 *m*), the food-passage and the organ of taste. Both the upper and lower cavities open behind into the gullet (pharynx). The hard

palate that separates them is formed by the joining of two lateral halves, the horizontal plates of the two upper-jaw processes, or the palate-plates (*p*). When these do not, sometimes, completely join in the middle, a longitudinal cleft remains, through which we can penetrate from the mouth straight into the nasal cavity. This is the malformation known as "wolf's throat." "Hare-lip" is the lesser form of the same defect. At the same time as the horizontal partition of the hard palate a vertical partition is formed by which the single nasal cavity is divided into two sections—a right and left half (Fig. 312 *m, n*).

skull, growing forwards from behind. The characteristic human nose is formed very late. Much stress is at times laid on this organ as an exclusive privilege of man. But there are apes that have similar noses, such as the long-nosed ape.

The evolution of the eye is not less interesting and instructive than that of the nose. Although this noblest of the sensory organs is one of the most elaborate and purposeful on account of its optic perfection and remarkable structure, it nevertheless develops, without preconceived design, from a simple process of the outer germinal layer. The fully-formed human eye is a round capsule, the



FIG. 313.



FIG. 314.

FIGS. 313 AND 314.—Upper part of the body of a human embryo, two-thirds of an inch long; of the sixth week. FIG. 313 from the left, FIG. 314 from the front. The organs of the nose and the upper lip from two lateral and originally separate halves can be clearly seen. Nose and upper lip are large in proportion to the rest of the face, and especially to the lower lip. (From Kollmann.)

The double nose has now acquired the characteristic form that man shares with the other mammals. Its further development is easy to follow; it consists of the formation of the inner and outer processes of the walls of the two cavities. The external nose is not formed until long after all these essential parts of the internal organ of smell. The first traces of it in the human embryo are found about the middle of the second month (Figs. 313-316). As can be seen in any human embryo during the first month, there is at first no trace of the external nose. It only develops afterwards from the foremost nasal part of the primitive

eye-ball (Fig. 317). This lies in the bony cavity of the skull, surrounded by protective fat and motor muscles. The greater part of it is taken up with a semi-fluid, transparent gelatinous substance, the *corpus vitreum*. The crystalline lens is fitted into the anterior surface of the ball (Fig. 317 *l*). It is a lenticular, bi-convex, transparent body, the most important of the refractive media in the eye. Of this group we have, besides the *corpus vitreum* and the lens, the watery fluid (*humor aqueus*) that is found in front of the lens (at the letter *m* in Fig. 317). These three transparent refractive media, by which the rays of light that

enter the eye are broken up and refocused, are enclosed in a solid round capsule, composed of several different coats, something like the concentric layers of an onion. The outermost and



FIG. 155.—Type of a human embryo, seven weeks old. (From *Development*.) Joining of the nasal process (p. 100) with the upper jaw process (p. 101) is visible, as corresponding.

thickest of these envelopes is the white *sclerotic coat* of the eye. It consists of tough white connective tissue. In front of the lens a circular, strongly-curved, transparent plate is fitted into the sclerotic, like the glass of a watch—the *cornea* (f.). At its outer surface the cornea is covered with a very thin layer of the epidermis, this is known as the *conjunctiva*. It goes from the cornea over the inner surface of the eye-lids, the upper and lower folds which we draw over the eye in closing it. At the inner corner of the eye we have a rudimentary organ in the shape of the relic of a third (inner) eye-lid, which is greatly developed, as "nictitating (winking) membrane," in the lower Vertebrates (p. 32). Underneath the upper eye-lid are the *lacrimal glands*, the product of which, the *lacrimal fluid*, keeps the outer surface of the eye smooth and clean.

Immediately under the sclerotic we find a very delicate, dark-red membrane, very rich in blood-vessels—the *choroid coat*—and inside this the *retina* (o), the expansion of the optic nerve (f). The latter is the second cerebral nerve. It proceeds from the optic thalami (the second cerebral

vesicle) to the eye, penetrates its outer envelope, and then spreads out like a net between the choroid and the corpus vitreum. Between the retina and the choroid there is a very delicate membrane, which is usually (but wrongly) associated with the latter. This is the black pigment-membrane (u). It consists of a single stratum of graceful, hexagonal, regularly-joined cells, full of granules of black colouring matter. This pigment membrane clothes, not only the inner surface of the choroid proper, but also the hind surface of its anterior muscular continuation, which covers the edge of the lens in front as a circular membrane, and arrests the rays of light at the sides. This is the well-known *iris* of the eye (f.), coloured differently in different individuals (blue, grey, brown, etc.); it forms the anterior border of the choroid. The circular opening that is left in the middle is the *pupil*, through which the rays of light penetrate into the eye. At the point where the lens leaves the anterior border of the choroid proper the latter is very thick, and forms a delicate crown of folds (g), which surrounds the edge of the lens with about seventy large and many smaller *rays* (*fenestrae ciliari*).

At a very early stage a couple of pear-shaped vesicles develop from the foremost part of the first cerebral vesicle in the embryo of man and the other Craniotes (Figs. 155 a, 297 a). These growths are the primary optic vesicles. They are at first directed outwards and forwards, but presently grow downward, so that, after the complete separation of the five cerebral vesicles, they lie at the base of the intermediate brain. The inner cavities of these pear-shaped vesicles, which soon attain a considerable size, are openly connected with the ventricle of the intermediate brain by their hollow stems. They are covered externally by the epidermis.

At the point where this comes into direct contact with the most curved part of the primary optic vesicle there is a thickening (f) and also a depression (o) of the hairy plate (Fig. 218, f). This pit, which we may call the *lens-pit*, is converted into a closed sac, the thick-



FIG. 156.—Part of a human embryo, eight weeks old. (From *Development*.)

walled lens-vesicle (2, 1), the thick edges of the pit joining together above it. In the same way in which the medullary tube separates from the outer germinal layer, we now see this lens-sac sever itself entirely from the horny plate (A), its source of origin. The hollow of the sac is afterwards filled with the cells of its thick walls, and thus we get the solid crystalline lens. This is, therefore, a purely epidermic structure. Together with the lens the small underlying piece of corneum-plate also separates from the skin.

As the lens separates from the corneous plate and grows inwards, it necessarily hollows out the contiguous primary optic vesicle (Fig. 318, 1-3). This is done in just the same way as the invagination of the blastula, which gives rise to the gastrula in the amphioxus (Fig. 38 C-F). In both cases the hollowing of the closed vesicle on one side goes so far that at last the inner, folded part touches the outer, not folded part, and the cavity disappears. As in the gastrula the first part is converted into the endoderm and the latter into the ectoderm, so in the invagination of the primary optic vesicle the retina (r) is formed from the first (inner) part, and the black pigment membrane (a) from the latter (outer, non-invaginated) part. The hollow part of the primary optic vesicle is converted into the optic nerve. The lens (1), which has so important a part in this process, acts at first directly on the invaginated part, or the retina (r). But they soon separate, a new structure, the corpus vitreum (v), growing between them. While the lenscular sac is being detached and is causing the invagination of the primary optic vesicle, another invagination is taking place from below; this proceeds from the superficial part of the skin-fibre layer—the corneum of the hood. Behind and under the lens a last-absorbed process from the cutis-plate (Fig. 319 g), hollows out the cap-shaped optic vesicle from below, and presses between the lens (1) and the retina (r). In this way the optic vesicle acquires the form of a hood.

Finally, a complete fibrous envelope, the fibrous capsule of the eye-bulb, is

formed about the secondary optic vesicle and its stem (the secondary optic nerve). It originates from the part of the head-plate which immediately encloses the eye. This fibrous envelope takes the form of a closed round vesicle, surrounding the whole of the ball and pushing between the lens and the horny plate at its outer side. The round wall of the capsule soon divides into two different membranes by surface-cleavage. The inner membrane becomes the choroid or vascular coat, and in front the ciliary corona and iris. The outer membrane is




FIG. 317.—The lampbrush eye in section. *a*, sclerotic coat; *b*, cornea; *c*, conjunctiva; *d*, vascular tunic of the eye; *e*, choroid coat; *f*, ciliary muscles; *g*, ciliary cilia; *h*, iris; *i*, optic nerve; *k*, anterior border of the cornea; *l*, crystalline lens; *m*, inner covering of the cornea (aqueous membrane); *n*, pigment membrane; *o*, retina; *p*, Part of canal; *q*, yellow spot of the retina. (From Huxley.)

converted into the white protective or sclerotic coat—in front, the transparent cornea. The eye is now formed in all its essential parts. The further development—the complicated differentiation and composition of the various parts—is a matter of detail.

The chief point in this remarkable evolution of the eye is the circumstance that the optic nerve, the retina, and the pigment membrane originate really from a part of the brain—an outgrowth of the intermediate brain—while the lens, the chief refractive body, develops from the outer skin. From the skin—the horny

plate—also arises the delicate conjunctiva, which afterwards covers the outer surface of the eyeball. The lacrimal glands are ramified growths from the conjunctiva (Fig. 286). All these important parts of



FIG. 286.—Eye of the shark embryo in longitudinal section (1) from an embryo 48 hours old, 2 from a somewhat older embryo, 3 from an embryo four days old. A, lens; B, iris; C, cornea; D, vitreous body; E, thickening of the horny plate at the point where the lens has severed itself; G, corpus vitreum; V, retina; P, pigment membrane. (From Huxley.)

the eye are products of the outer germinal layer. The remaining parts—the corpus vitreum (with the vascular capsule of the lens), the choroid (with the iris), and the sclerotic (with the cornea)—are formed from the middle germinal layer.

The outer protection of the eye, the eye-lid, are merely folds of the skin, which are formed in the third month of human embryonic life. In the fourth month the upper eye-lid reaches the lower, and the eye remains covered with them until birth. As a rule, they open wide shortly before birth (sometimes only after birth). Our craniote ancestors had a third eye-lid, the nictitating membrane, which was drawn over the eye from its inner angle. It is still found in many of the Selachii and Amniotes. In the apes and man it has degenerated, and there is now only a small relic of it at the inner corner of the eye, the lacrimonial fold, a useless rudimentary organ (cf. p. 32). The apes and man have also lost the Harderian gland that opened under the nictitating membrane; we find this in the rest of the mammals, and the birds, reptiles, and amphibia.

The peculiar embryonic development of the vertebrate eye does not enable us to draw any definite conclusions as to its obscure phylogeny; it is clearly convergent to a great extent, or obscured by the reduction and curtailment of its original features. It is probable that many of the earlier stages of its phylogeny have disappeared without leaving a trace.

It can only be said positively that the peculiar ontogeny of the complicated optic apparatus in man follows just the same laws as in all the other Vertebrates. Their eye is a part of the fore brain, which has grown forward towards the skin, not an original cutaneous sense-organ, as in the Invertebrates.

The vertebrate ear resembles the eye and nose in many important respects, but is different in others, in its development. The auscultory organ in the fully-developed man is like that of the other mammals, and especially the apes, in the main features. As in them, it consists of two chief parts—an apparatus for conducting sound (external and middle ear) and an apparatus for the sensation of sound (internal ear). The external ear opens in the skull at the side of the head (Fig. 320 a). From this point the external passage (b), about an inch in length, leads into the head. The inner end of it is closed by the tympanum, a vertical, but not quite upright, thin membrane of an oval shape (c). This tympanum separates the external passage from the tympanic cavity (d). This is a small cavity, filled with air, in the temporal bone; it is connected with the mouth by a special tube.



FIG. 320.—Horizontal transverse section of the ear of a human embryo, four weeks old (magnified one hundred times). (From Huxley.) 1, inner (the dark wall of which is as thick as the diameter of the central artery); 2, corpus vitreum (constituted by a stalk, g, with the corpus vitreum (pressing behind the lens) inside the corpus vitreum by means of the stem g); 3, retina (inner portion, invaginated layer of the primary optic vesicle); 4, pigment membrane (outer, thin, non-invested layer of same); 5, space between retina and pigment membrane (remainder of the cavity of the primary optic vesicle).

This tube is rather longer, but much narrower, than the outer passage, leads inwards obliquely from the anterior wall of the tympanic cavity, and opens in the throat below, behind the nasal

opening. It is called the Eustachian tube (*e*); it serves to equalize the pressure of the air within the tympanic cavity and the outer atmosphere that enters by the external passage. Both



FIG. 300.—The human ear. Left ear, seen from the front, external aspect. *a* shell of ear, *b* external passage, *c* tympanum, *d* tympanic cavity, *e* Eustachian tube, *f, g, h* the three bones of the ear (*f* hammer, *g* anvil, *h* stirrup), *i* cochlea, *j* the three semi-circular canals, *k* the sacculus, or cochlea, *l* the auditory nerve.

the Eustachian tube and the tympanic cavity are lined with a thin mucous coat, which is a direct continuation of the mucous lining of the throat. Inside the tympanic cavity there are three small bones which are known (from their shape) as the hammer, anvil, and stirrup (Fig. 300, *f, g, h*). The hammer (*f*) is the outermost, next to the tympanum. The anvil (*g*) fits between the other two, above and inside the hammer. The stirrup (*h*) lies inside the anvil, and touches with its base the outer wall of the internal ear, or auditory vesicle. All these parts of the external and middle ear belong to the apparatus for conducting sound. Their chief task is to convey the waves of sound through the thick wall of the head to the inner-lying auditory vesicle. They are not found at all in the fishes. In these the waves of sound are conveyed directly by the wall of the head to the auditory vesicle.

The internal apparatus for the sensation

of sound, which receives the waves of sound from the conducting apparatus, consists in man and all other mammals of a closed auditory vesicle filled with fluid and an auditory nerve, the ends of which expand over the wall of this vesicle. The vibrations of the sound-waves are conveyed by these media to the nerve-endings. In the labyrinthine water that fills the auditory vesicle there are small stones at the points of entry of the acoustic nerves, which are composed of groups of microscopic calcareous crystals (otoliths). The auditory organ of most of the Invertebrates has substantially the same composition. It usually consists of a closed vesicle, filled with fluid, and containing otoliths, with the acoustic nerve expanding on its wall. But, while the auditory vesicle is usually of a simple round or oval shape in the Invertebrates, it has in the Vertebrates a special and curious structure, the labyrinth. This thin-membraned labyrinth is enclosed in a bony capsule of the same shape, the osseous labyrinth (Fig. 321), and this lies in the middle of the petrous bone of the skull. The labyrinth is divided into two vesicles in all the Gnathostomes. The larger one is called the *utricle*, and has three arched appendages, called the "semi-circular canals" (*c, d, e*). The smaller vesicle is called the *sacculus*, and is connected with a peculiar appendage, with (in man and the higher mammals) a spiral form something like a snail's shell, and therefore called the *cochlea* (= snail, *h*). On the thin wall of this delicate labyrinth the acoustic nerve, which comes from the after-brain, spreads out in most elaborate fashion. It divides

into two main branches—a cochlear nerve (for the cochlea) and a vestibular nerve (for the rest of the labyrinth). The former seems to have more to do with the equality, the latter with the quantity, of the acoustic sensations. Through the cochlear nerves we learn the height and timbre, through the vestibular nerves the intensity, of tones.

The first structure of this highly elaborate organ is very simple in the embryo of man and all the other Craniotes; it is a



FIG. 321.—The bony labyrinth of the human ear. Left side. *a* utricle, *b* saccule, *c* upper canal of posterior canal, *d* outer canal, *e* oval window, *f* round window. (From Meyer.)

pit-like depression in the skin. At the back part of the head at both sides, near the after brain, a small thickening of the horny plate is formed at the upper end of the second gill-cleft (Fig. 322 *A B*). This sinks into a sort of pit, and covers from the epidermis, just as the lens of the eye does. In this way is formed at each side, directly under the horny plate of the back part of the head, a small vesicle filled with fluid, the primitive auricular vesicle, or the primary labyrinth. As it separates from its source, the horny plate, and presses inwards and backwards into the skull, it changes from round to pear-shaped (Figs. 322 *B C*, 323 *a*). The outer part of it is lengthened into a thin stem, which at first still opens outwards by a narrow canal. This is the labyrinthine appendage (Fig. 322 *B*). In the lower Vertebrates it develops into a special cavity filled with

fluid in the shape of simple pouch-like invaginations of the utricle (*cap* and *cap*). The edges join together in the middle part of each fold, and separate from the utricle, the two ends remaining in open connection with its cavity. All the Gnathostomes have these three canal-like man, whereas among the Cyclostomes the lampreys have only two and the hag-fishes only one. The very complex structure of the cochlea, one of the most elaborate and wonderful outcome of adaptation in the mammal body develops originally in very simple fashion as a flask-like projection from the succulus. As Hesse and Reizner have pointed out we find the successive ontogenetic stages of its growth represented permanently in the series of the higher Vertebrates. The cochlea is wanting even in the Monotremes, and is restricted to the rest of the mammals and man.




FIG. 322.—Development of the auditory labyrinth of the chick, in six successive stages (*A–F*). (Vertical transverse sections of the skull.) *A*, auditory vesicle; *B*, auditory vesicle; *C*, labyrinthine appendage; *D*, secondary cavity; *E*, posterior canal; *F*, external canal; *G*, papillary vein. (From Reizner.)

calcareous crystals, which remains open permanently in some of the primitive fishes, and opens outwards in the upper part of the skull. But in the mammals the labyrinthine appendage degenerates. In these it has only a phylogenetic interest as a rudimentary organ, with no actual physiological significance. The useless relic of it passes through the wall of the petrous bone in the shape of a narrow canal, and is called the vestibular aqueduct.

It is only the inner and lower bulbous part of the separated auricular vesicle that develops into the highly complex and differentiated structure that is afterwards known as the secondary labyrinth. This vesicle divides at an early stage into an upper and larger and a lower and smaller section. From the one we get the *utricle* with the semi-circular canals; from the other the *sacculus* and the *cochlea* (Fig. 320 *c*). The canals are

The auditory nerve, an eighth cranial nerve, expands with one branch in the cochlea, and with the other in the remaining parts of the labyrinth. This nerve is, as Gegenbaur has shown, the sensory dorsal branch of a cerebro-spinal nerve, the motor ventral branch of which acts for the muscles of the face (*nervus facialis*). It has therefore origi-

nated phylogenetically from an ordinary cutaneous nerve, and so is of quite different origin from the optic and olfactory nerves, which both represent direct outgrowths of the brain. In this respect the auditory organ is essentially different from the organs of sight and smell. The acoustic nerve is formed from ectodermic cells of the hind brain, and develops from the nervous structure that appears at its dorsal limit. On the other hand, all the membranous, cartilaginous, and osseous coverings of the labyrinth are formed from the mesodermic head-plates.

The apparatus for conducting sound which we find in the external and middle ear of mammals develops quite separately from the apparatus for the sensation of sound. It is both phylogenetically and ontogenetically an independent secondary formation, a later accession to

the primary internal ear. Nevertheless, its development is not less interesting, and is explained with the same ease by comparative anatomy. In all the fishes and in the lowest Vertebrates there is no



FIG. 341.—Primitive skull of the human embryo, four weeks old, ventral aspect, left half seen (cf. middle of p. 284). *a* the eye, *b* the ear, *c* the mouth, *d* the cerebellum, *e* the cerebellar vermis, *f* the cerebellar hemispheres, *g* the cerebellar peduncles, *h* the cerebellar vermis, *i* the cerebellar hemispheres, *j* the cerebellar peduncles, *k* the cerebellar vermis, *l* the cerebellar hemispheres, *m* the cerebellar peduncles, *n* the cerebellar vermis, *o* the cerebellar hemispheres, *p* the cerebellar peduncles, *q* the cerebellar vermis, *r* the cerebellar hemispheres, *s* the cerebellar peduncles, *t* the cerebellar vermis, *u* the cerebellar hemispheres, *v* the cerebellar peduncles, *w* the cerebellar vermis, *x* the cerebellar hemispheres, *y* the cerebellar peduncles, *z* the cerebellar vermis. (From Kollmer.)

special apparatus for conducting sound, no external or middle ear, they have only a labyrinth, an internal ear, which lies within the skull. They are without the tympanum and tympanic cavity, and all its appendages. From many observations made in the last few decades it seems that many of the fishes (if not all) cannot distinguish tones; their labyrinth seems to be chiefly (if not exclusively) an organ for the sense of space (or equilibrium). If it is destroyed, the fishes lose their balance and fall. In the opinion of recent physiologists this applies also to many of the Invertebrates (including the higher ancestors of the Vertebrates). The round vesicles which are considered to be their auscultory vesicles, and which contain an otolith, are supposed to be merely organs of the sense of space ("static vesicles or statocysts").

The middle ear makes its first appearance in the amphibian class, where we find a tympanum, tympanic cavity, and Eustachian tube; these animals, and all terrestrial Vertebrates, certainly have the faculty of hearing. All these essential parts of the middle ear originate from the first gill-cleft and its surrounding part; in the *Seichth* this remains throughout life an open squaring-hole, and first between the first and second gill-arch. In the embryo of the higher Vertebrates it closes up in the centre, and thus forma

the tympanic membrane. The outlying remainder of the first gill-cleft is the rudiment of the external meatus. From its inner part we get the tympanic cavity, and, further inward still, the Eustachian tube. Connected with this is the development of the three bones of the mammalian ear from the first two gill-arches; the hammer and anvil are formed from the first, the stirrup from the upper end of the second, gill-arch.

Finally, the shell (pinna or concha) and external meatus (passage to the tympanum) of the outer ear are developed in a very simple fashion from the skin that borders the external aperture of the first gill-cleft. The shell rises in the shape of a circular fold of the skin, in which cartilage and muscles are afterwards formed (Figs. 313 and 315). This organ is only found in the mammalian class. It is very rudimentary in the lowest section, the Monotremes. In the others it is found at very different stages of development, and sometimes of degeneration. It is degenerate in most of the aquatic mammals. The majority of them have lost it altogether—for instance, the walrus and whales and most of the seals. On the other hand, the pinna is



FIG. 342.—The rudimentary muscles of the ear in the human skull. *a* raising muscle (*M. attollens*), *b* drawing muscle (*M. attrahens*), *c* withdrawing muscle (*M. retrahens*), *d* large muscle of the helix (*M. helix major*), *e* small muscle of the helix (*M. helix minor*), *f* muscle of the angle of the ear (*M. angulus*), *g* antihelical muscle (*M. antihelical*). (From H. Meyer.)

well developed in the great majority of the Marsupials and Placentals; it occurs as and collects the waves of sound, and is equipped with a very elaborate muscular apparatus, by means of which the pinna

can be turned freely in any direction and its shape be altered. It is well known how readily domestic animals—horses, cows, dogs, hares, etc.—pivot their ears and move them in different directions. Most of the apes do the same, and our earlier ape ancestors were also able to do it. But our later simian ancestors, which we have in common with the anthropoid apes, abandoned the use of these muscles, and they gradually became rudimentary and useless. However, we possess them still (Fig. 324). In fact, some men can still move their ears a little backward and forward by means of the drawing and withdrawing muscles (*b* and *c*); with practice this faculty can be much improved. But no man can now lift up his ears by the raising muscle (*a*), or change the shape of them by the small inner muscles (*d, e, f, g*). These muscles were very useful to our ancestors, but are of no consequence to us. This applies to most of the anthropoid apes as well.

We also share with the higher anthropoid apes (gorilla, chimpanzee, and orang) the characteristic form of the human outer ear, especially the folded border, the helix and the lobe. The lower apes have pointed ears, without folded border or lobe, like the other mammals. But Darwin has shown that at the upper

part of the folded border there is in many men a small pointed process, which most of us do not possess. In some individuals this process is well developed. It can only be explained as the relic of the original point of the ear, which has been turned inwards in consequence of the curving of the edge. If we compare the pinna of man and the various apes in this respect, we find that they present a connected series of degenerate structures. In the common catarrhine ancestors of the anthropoids and man the degeneration set in with the folding together of the pinna. This brought about the helix of the ear, in which we find the significant angle which represents the relic of the absent point of the ear in our earliest simian ancestors. Here again, therefore, comparative anatomy enables us to trace with certainty the human ear to the similar, but more developed, organ of the lower mammals. At the same time, comparative physiology shows that it was a more or less useful implement in the latter, but it is quite useless in the anthropoids and man. The conducting of the sound has scarcely been affected by the loss of the pinna. We have also in this the explanation of the extraordinary variety in the shape and size of the shell of the ear in different men; in this it resembles other rudimentary organs.


CHAPTER XXVI.

EVOLUTION OF THE ORGANS OF MOVEMENT

THE peculiar structure of the locomotive apparatus is one of the features that are most distinctive of the vertebrate stem. The chief part of this apparatus is formed, as in all the higher animals, by the active organs of movement, the muscles; in consequence of their contractility they have the power to draw up and shorten themselves. This effects the movement of the various parts of the body, and thus the whole body is conveyed from place to place. But the

arrangement of these muscles and their relation to the solid skeleton are different in the Vertebrates from the Invertebrates.

In most of the lower animals, especially the Platyodes and Vermalis, we find that the muscles form a simple, thin layer of flesh immediately underneath the skin. This muscular layer is very closely connected with the skin itself; it is the same in the Mollusc stem. Even in the large division of the Articulata, the classes of crabs, spiders, myriapods, and



insects, we find a similar feature, with the difference that in this case the skin forms a solid armour—a rigid cutaneous skeleton made of chitin (and often also of carbonate of lime). This external chitine coat undergoes a very elaborate articulation both on the trunk and the limbs of the Arthropods, and in consequence the muscular system also, the contractile fibres of which are attached inside the chitine tubes, is highly articulated. The Vertebrates form a direct contrast to this. In them alone a solid internal skeleton is developed, of cartilage or bone, to which the muscles are attached. This bony skeleton is a complex lever apparatus, or *force* apparatus of movement. Its rigid parts, the arms of the levers, or the bones, are brought together by the actively mobile muscles, as if by drawing-rope. This admirable locomotorium, especially its skeleton, is a special feature of the Vertebrates, and has given the name to the group.

In order to get a clear idea of the chief features of the development of the human skeleton, we must first examine its composition in the adult frame (Fig. 325, the human skeleton seen from the right, Fig. 326, from view of the whole skeleton). As in other mammals, we distinguish first between the axial or dorsal skeleton and the skeleton of the limbs. The axial skeleton consists of the vertebral column (the skeleton of the trunk) and the skull (skeleton of the head); the latter is a peculiarly modified part of the former. As appendages of the vertebral column we have



FIG. 325.—The human vertebral column (showing upright form of the right side). (From H. Meyer.)

the ribs, and of the skull we have the hyoid bone, the lower jaw, and the other products of the gill-arches.

The skeleton of the limbs or extremities is composed of two groups of parts—the skeleton of the extremities proper and the auto-skeleton, which connects these with the vertebral column. The auto-skeleton of the arms (or fore legs) is the shoulder-girdle; the auto-skeleton of the legs (or hind legs) is the pelvic zone.

The vertebral column (Fig. 327) in man is composed of thirty-three to thirty-five ring-shaped bones in a continuous series (about each other, in man's upright position). These vertebrae are separated from each other by elastic ligaments, and at the same time connected by joints, so that the whole column forms a firm and solid, but flexible and elastic, axial skeleton, moving freely in all directions. The vertebrae differ in shape and connection at the various parts of the trunk, and we distinguish the following groups in the series, beginning at the top: seven cervical vertebrae, twelve dorsal vertebrae, five lumbar vertebrae, five sacral vertebrae, and four to six caudal vertebrae. The uppermost, or those next to the skull, are the cervical vertebrae (Fig. 327). They have a hole in each of the lateral processes. There are seven of these vertebrae

in man and almost all the other mammals, even the neck is as long as that of the camel or giraffe, or as short as that of the mole or hedgehog. This constant number, which has few exceptions (due to adaptation), is a strong proof of the common descent of the mammals; it can only be explained by faithful heredity from a common stem-form, a primitive mammal with seven cervical vertebrae. If each species had been created separately, it would have been better to have given the long-necked mammals more, and the short-necked animals less, cervical vertebrae. Next to these come the dorsal (or postoral)




FIG. 327.—A piece of the axial rod (vertebra) from a sheep's skull, showing the dorsal process. (From H. Meyer.)

the ribs, and of the skull we have the hyoid bone, the lower jaw, and the other products of the gill-arches.

vertebrae, which number twelve to thirteen (usually twelve) in man and most of the other mammals. Each dorsal vertebra (Fig. 165) has at the side, connected by joints, a couple of ribs, long bony arches



FIG. 34.—Three dorsal vertebrae, from a human embryo, eight weeks old, in lateral longitudinal section. *a*, cartilaginous vertebral body; *b*, intervertebral disk; *c*, chorda; *d*, ribs. (From Kolliker.)

that lie in and protect the wall of the chest. The twelve pairs of ribs, together with the connecting intercostal muscles and the sternum, which joins the ends of the right and left ribs in front, form the chest (*thorax*). In this strong and elastic frame are the lungs, and between them the heart. Next to the dorsal vertebra comes a short but stronger section of the column, formed of five large vertebrae. These are the lumbar vertebrae (Fig. 166); they have no ribs and no holes in the transverse processes. To these succeeds the sacral bone, which is fitted between the two halves of the pelvic girdle. The sacrum is formed of five vertebrae, completely blended together. Finally, we have at the end a small rudimentary caudal column, the *caevæ*. This consists of a varying number (usually four, more rarely three, or five or six) of small degenerated vertebrae, and is a useless rudimentary organ with no actual physiological significance. Morphologically, however, it is of great interest as an irrefragable proof of the descent of man and the anthropoids from long-tailed apes. On no other theory can we explain the existence of this rudimentary tail. In the earlier stages of development the tail of the human embryo protrudes considerably. It afterwards atrophies; but the role of the atrophied caudal vertebrae and of the rudimentary muscles that once moved it remains permanently. Sometimes, in fact, the external tail is preserved. The older anatomists say that the tail is usually one vertebra longer in the human female than in the male (or four against five). Steinbach says it is the reverse.

In the human vertebral column there are usually thirty-three vertebrae. It is interesting to find, however, that the

number often changes, one or two vertebrae dropping out or an additional one appearing. Often, also, a mobile rib is formed at the last cervical or the first lumbar vertebra, so that there are then thirteen dorsal vertebrae, besides six cervical and four lumbar. In this way the contiguous vertebrae of the various sections of the column may take each other's place.

In order to understand the embryology of the human vertebral column we must first carefully consider the shape and connection of the vertebrae. Each vertebra has, in general, the shape of a seal-ring (Figs. 164-166). The thicker portion, which is turned towards the ventral side, is called the body of the vertebra, and forms a short osseous disk; the thinner part forms a semi-circular arch, the *vertebral arch*, and is turned towards the back. The arches of the successive vertebrae are connected by thin intervertebral ligaments in such a way that the cavity they collectively enclose represents a long canal. In this vertebral canal we find the trunk part of the central nervous system, the spinal cord. Its head part, the brain, is enclosed by the skull, and the skull itself is merely the uppermost part of the vertebral column, distinctively modified. The base or ventral side of the vascular cranial capsule corresponds originally to a number of developed vertebral bodies; its vault or dorsal side to their combined upper vertebral arches.

While the solid, massive bodies of the vertebrae represent the real central axis of



FIG. 35.—A dorsal vertebra of the same order as in lateral-transverse section. *a*, cartilaginous vertebral body; *b*, chorda; *c*, transverse process; *d*, vertebral arch (upper arch); *e*, upper end of the rib (lower arch). (From Kolliker.)

the skeleton, the dorsal arches serve to protect the central marrow they enclose. But similar arches develop on the ventral side for the protection of the viscera in the breast and belly. These lower or

ventral vertebral arches, proceeding from the ventral side of the vertebral bodies, form, in many of the lower Vertebrates, a canal in which the large blood-vessels are enclosed on the lower surface of the



FIG. 271.—Inter-vertebral disc of a sea-horse (horse), transverse section. a rest of the chorda. (From Atthiler.)

vertebral column (aorta and caudal vein). In the higher Vertebrates the majority of these vertebral arches are lost or become rudimentary. But at the thoracic section of the column they develop into independent strong osseous arches, the ribs (*costae*). In reality the ribs are merely large and independent lower vertebral arches, which have lost their original connection with the vertebral bodies.

If we turn from this anatomic survey of the composition of the column to the question of its development, I may refer the reader to earlier pages with regard to the first and most important points (pp. 145-148). It will be remembered that in the human embryo and that of the other vertebrates we find at first, instead of the segmented column, only a simple unarticulated cartilaginous rod. This solid but flexible and elastic rod is the axial rod (or the *chorda dorsalis*). In the lowest Vertebrate, the Amphioxus, it retains this simple form throughout life, and permanently represents the whole internal skeleton (Fig. 210 f). In the Tunicates, also, the nearest Invertebrate relatives of the Vertebrates, we meet the same chorda—transitorily in the passing larva tail of the Ascidia, permanently in the Cephalopoda (Fig. 225 c). Undoubtedly both the Tunicates and Acrania have inherited the chorda from a common unsegmented stem-form; and these ancient, long-extinct ancestors of all the

chordonia are our hypothetical Prochordonia.

Long before there is any trace of the skull, limbs, etc., in the embryo of man or any of the higher Vertebrates—at the early stage in which the whole body is merely a sail-shaped embryonic shield—there appears in the middle line of the shield, directly under the medullary furrow, the simple chorda. (Cf. Figs. 131-135 ab). It follows the long axis of the body in the shape of a cylindrical axial rod of elastic but firm composition, equally pointed at both ends. In every case the chorda originates from the dorsal wall of the primitive gut; the cells that compose it (Fig. 328 f) belong to the ectoderm (Figs. 216-221). At an early stage the chorda develops a transparent structureless sheath, which is secreted from its cells (Fig. 328 a). This *chordallemma* is often called the "inner chorda-sheath," and must not be confused with the real external sheath, the mesoblastic perichorda.

But this unsegmented primary axial skeleton is soon replaced by the segmented secondary axial skeleton, which we know as the vertebral column. The provertebral plates (Fig. 124 f) differentiate from the innermost, median part of the visceral layer of the ectoderm-pouches at each side of the chorda. As they grow round the chorda and enclose it they form the skeleton plate or skeletogenetic layer—that is to say, the skeleton-forming stratum of cells, which provides the mobile foundation of the permanent vertebral column and skull (scleroblasts). In the head-half of the embryo the



FIG. 322.—Human skull.

skeletal plate remains a continuous, simple, undivided layer of tissue, and presently enlarges into a thin-walled capsule enclosing the brain, the primordial skull. In the trunk-half the provertebral




FIG. 224.—Rear of the skulls of nine Primates (Cathartidae), seen from above and reduced to a common size. 1. *Mediterraneus*, 2. *Samlapeui*, 3. *Pithecanthropus*, 4. *Gorilla*, 5. *Schimpanse*, 6. *Orang*, 7. *Hylobates*, 8. *Presbytis*, 9. *Cynocephalus*.

branchial arches of the Selachii prove that their skull originally consisted of a large number of (at least nine or ten) warts, and that the cerebral nerves had proceeded from the base of the brain entirely confirm this. These cerebral nerves are (with the exception of the first and second pair, the olfactory and optic nerves) merely modifications of spinal nerves, and are essentially similar to them in their peripheral expansion. The comparative anatomy of these cerebral nerves, their origin and their expansion, furnishes one of the strongest arguments for the new vertebral theory of the skull.

We have not space here to go into the details of Gegenbaur's theory of the

each side—the primitive upper jaw (or *palato-quadratum*, *o*) and the primitive lower jaw (*a*); *IV*, the hyaloid bone (*II*); finally, *V-X*, six branchial arches in the narrower sense (*III-VIII*). From the anatomic features of these nine to ten cranial ribs or "lower vertebral arches" and the cranial nerves that spread over them, it is clear that the apparently simple cartilaginous primitive skull of the Selachii was originally formed from so many (at least nine) somites or provertebrae. The blending of these primitive segments into a single capsule is, however, so ancient that, in virtue of the law of curtailed heredity, the original division seems to have dis-




FIG. 324.—Skeleton of the hand or foot of various mammals. *I* man, *II* dog, *III* pig, *IV* eagle, *V* tiger, *VI* horse. *a* thumb, *b* index, *c* middle, *d* ring, *e* little, *f* metacarpus, *g* trapezium, *h* trapezoid, *i* trapezoid, *j* trapezoid, *k* trapezoid, *l* trapezoid, *m* trapezoid, *n* trapezoid, *o* trapezoid, *p* trapezoid, *q* trapezoid, *r* trapezoid, *s* trapezoid, *t* trapezoid, *u* trapezoid, *v* trapezoid, *w* trapezoid, *x* trapezoid, *y* trapezoid, *z* trapezoid. (*From Gegenbaur*)

skull, I must be content to refer the reader to the great work I have mentioned, in which it is thoroughly established from the empirico-philosophical point of view. He has also given a comprehensive and up-to-date treatment of the subject in his *Comparative Anatomy of the Vertebrates* (1868). Gegenbaur indicates an original "cranial ribs," or "lower arches of the cranial vertebrae," at each side of the head of the Selachii (Fig. 324), the following pairs of arches: *I* and *II*, two tip cartilages, the anterior (*a*) of which is composed of an upper piece only, the posterior (*b*) from an upper and lower piece; *III*, the maxillary arches, also consisting of two pieces on

appeared; in the embryonic development it is very difficult to detect it in isolated traces, and in some respects quite impossible. It is claimed that several (three to six) traces of provertebrae have been discovered in the anterior (pre-chordal) part of the Selachii-skull: this would bring up the number of cranial somites to twelve or sixteen, or even more.

In the primitive skull of man (Fig. 325) and the higher Vertebrates, which has been evolved from that of the Selachii, five consecutive sections are discoverable at a certain early period of development, and one might be induced to trace them to five primitive vertebrae; but these sections are due entirely to adaptation to




FIG. 343.

FIG. 344.

FIG. 345.

FIGS. 343-45.—ARM AND HAND OF THREE BAT-SPECIES. FIG. 343 *Chiropterus (Antrozous) niger*. FIG. 344 *Vesperugo (Hesperugo) arcticus*. FIG. 345 *Eptesicus (Hesperugo) nocturnus*. (From FAUL and FINE, 1928.)

the five primitive cerebral vesicles, and correspond, like these, to a large number of metamers. That we have in the primitive skull of the mammals a greatly modified and transformed organ, and not at all a primitive formation, is clear from the circumstance that its original soft membranous form only assumes the cartilaginous character for the most part at the base and the sides, and remains membranous at the roof. At this part the bones of the subsequent osseous skull develop as external coverings over the membranous structure, without an intermediate cartilaginous stage, as there is at the base of the skull. Thus a large part of the cranial bones develop originally as covering bones from the corium, and only secondarily come into close touch with the primitive skull (Fig. 333). We have previously seen how this very rudimentary beginning of the skull in man is formed ontogenetically from the "head-plates," and thus the fore end of the chorda is enclosed in the base of the skull. (Cf. Fig. 143 and pp. 138, 144, and 149.)

The phylogeny of the skull has made great progress during the last three decades through the joint attainments of comparative anatomy, ontogeny, and paleontology. By the judicious and comprehensive application of the phylogenetic method (in the sense of Gegenbaur) we have found the key to the great and important problems that arise from the thorough comparative study of the skull. Another school of research, (the school of what is called "exact craniology" (in the sense of Virchow), has, meantime, made fruitless efforts to obtain this result. We may gratefully acknowledge all that this descriptive school has done in the way of accurately describing the various forms and measurements of the human skull, as compared with those of other mammals. But the vast empirical material that it has accumulated in its extensive literature is more dead and sterile erudition until it is vivified and illumined by phylogenetic speculation.

Virchow confined himself to the most careful analysis of large numbers of human skulls and those of anthropoid mammals. He saw only the differences between them, and sought to express these in figures.

Without adducing a single solid reason, or offering any alternative explanation, he rejected evolution as an unproved hypothesis. He played a most unfortunate

part in the controversy as to the significance of the fossil human skulls of Spy and Neanderthal, and the comparison of these with the skull of the *Pithecanthropus* (Fig. 383). All the interesting features of these skulls that clearly indicated the transition from the anthropoid to the man were declared by Virchow to be chance pathological variations. He said that the roof of the skull of *Pithecanthropus* (Fig. 335, 3) must have belonged to an ape, because so pronounced an *orbital stricture* (the horizontal constriction between the outer edge of the eye-orbit and the temples) is not found in any human being. Immediately afterwards Nehring showed in the skull of a Brazilian Indian (Fig. 335, 2), found in the Sambaquis of Santos, that this stricture can be even deeper in man than in many of the apes. It is very



FIG. 344.—Transverse section of a fish's skull (from the trunk). (From Johannes Müller.) A, upper (dorsal) lateral muscles; C, D, lower (ventral) lateral muscles; B, vertical fishes; E, section of incomplete coronal suture (in A at basilar base of the inter-orbital ligaments (from the side).

instructive in this connection to compare the roofs of the skulls (seen from above) of different primates. I have, therefore, arranged nine such skulls in Fig. 335, and reduced them to a common size.

We turn now to the branchial arches, which were regarded even by the earliest natural philosophers as "head-ribs." (Cf. Figs. 167-170). Of the four original gill-arches of the mammals the first lies between the primitive mouth and the first gill-cleft. From the base of this arch is formed the upper-jaw process, which joins with the inner and outer nasal processes on each side, in the manner we have previously explained, and forms the chief parts of the skeleton of the upper jaw (palate bone, pterygoid bone, etc.) (Cf. p. 284.) The remainder of the first branchial arch, which is now called, by

way of contrast, the M_1 forms from its base two (hammer and anvil), are converted into a long 1 that is known, after 1 "Möckel's cartilage," or of the

from the cellular ring or accessory bone, the permanent bony lower jaw. From the first part or base of the second branchial arch we get, in the mammals, the third ossicle of the ear, the stapes; and from the succeeding parts we get (in this order) the muscle of the ear, the styloid process of the

third branchial arch is only cartilaginous at the foremost part, and here the body of the hyoid bone and its larger horn are formed at each side by the junction of its two halves. The fourth branchial arch is only found transiently in the mammalian embryo as a rudimentary organ, and does not develop special parts, and there is no trace in the embryo of the higher Vertebrates of the posterior branchial arches (fifth and sixth pair), which are permanent in the Selachii. They have been lost long ago. Moreover, the four gill-clefts of the human embryo are only interesting as rudimentary organs, and they soon close up and disappear. The first alone (between the first and second branchial arches) has any permanent significance; from it are developed the tympanic cavity and the Eustachian tube. (Cf. Figs. 169, 320.)

It was Carl Gegenbaur again who solved the difficult problem of tracing the skeleton of the limbs of the Vertebrates to a common type. Few parts of the vertebrate body have undergone such infinitely varied modifications in regard to size, shape, and adaptation of structure as the limbs or extremities; yet we are in a position to reduce them all to the same hereditary standard. We may generally distinguish three groups among the Vertebrates in relation to the formation of their limbs. The lowest and earliest Vertebrates, the Acrania and Cyclostomes, had, like their invertebrate ancestors, no pairs of limbs, as we see in the Amphioxus and the Cyclostomes to-day (Figs. 210, 247). The second group is formed of the two classes of the true fishes and the Dipneusts; here there are always two pairs of limbs at first, in the shape of many-toed fins—one pair

of breast-fins or fore legs, and one pair of belly-fins or hind legs (Figs. 248-259). The third group comprises the four higher classes of Vertebrates—the amphibians, reptiles, birds, and mammals; in these quadrupeds there are at first the same two pairs of limbs, but in the shape of five-toed feet. Frequently more than five toes, and sometimes the foot is wholly atrophied (Figs. 260-263). But the original stem-form of the group had five toes or fingers before and behind (Figs. 263-265).

The true primitive form of the pairs of limbs they were found in primitive fishes of the Silurian period, is preserved for us in the Australian dipneust, the remarkable *Ceratodus* (Fig. 257). Both the breast-fin and the belly-fin are flat oval paddles, in which we find a bimerial cartilaginous skeleton (Fig. 336). This consists, firstly, of a much segmented fin-rod or "stern" (A, B), which runs through the fin from base to tip; and secondly of a double row of thin articulated fin-radii (r, r), which are attached to both sides of the fin-rod, like the feathers of a feathered leaf. This primitive fin, which Gegenbaur first recognised, is attached to the vertebral column by a single zone in the shape of a cartilage.

It has probably originated from the b

W^hile Gegenbaur's primitive fin more or less preserved in the fossilized remains of the earliest Selachii (Fig. 258), Ganoids (Fig. 253), and Dipneusts (Fig. 256). It is also found in modified form in some of the actual sharks and pilce. But in the majority of the Selachii it has already degenerated to the extent that the radii on one side of the fin-rod have been partly or entirely lost, and are retained only on the other (Fig. 337). Thus we get the uniserial fin, which has been led from the Selachii to the rest of the fishes (Fig. 338).

Gegenbaur has shown how the five-toed leg of the Amphibia, that has been

¹ While Gegenbaur derives 1 from the anterior segmental branch, 1 and 1 they have been derived from a pair of originally continuous lateral fin rays.

² The fin of the four highest of the group was developed at the same time as the lower fin of the lower (lower or higher) fin, and was in the fifth fin 1 ; it was formerly believed to be at the inner (basal or dorsal) side, and was at the first




FIG. 347.

FIG. 348.

FIG. 347—Human skeleton. (Cf. Fig. 346.)

FIG. 348—Skeleton of the giant gorilla. (Cf. Fig. 346.)

In the dipneust ancestors of the Amphibia the radii gradually atrophy, and are lost, for the most part, on the other side of the fin-rod as well (the lighter cartilages in Fig. 338). Only the four lowest radii (shaded in the illustration) are preserved; and these are the four inner toes of the foot (first to fourth). The fifth or fifth toe is developed from the lower end of the fin-rod. From the middle and upper part of the fin-rod was developed the long stem of the limb—the important radius and ulna (Fig. 339 r and u) and humerus (h) of the higher Vertebrates.

In this way the five-toed foot of the Amphibia, which we first meet in the Carboniferous *Stegcephala* (Fig. 260), and which was inherited from them by the reptiles on one side and the mammals on the other, was formed by gradual degeneration and differentiation from the many-toed fish-fin (Fig. 341). The reduction of the radii to four was accompanied by a further differentiation of the fin-rod, its transverse segmentation into upper and lower halves, and the formation of the zone of the limb, which is composed originally of three limbs before and behind in the higher Vertebrates. The simple arch of the original shoulder-zone divides on each side into an upper (dorsal) piece, the shoulder-blade (*scapula*), and a lower (ventral) piece, the anterior part of the latter forms the primitive clavicle (*procoracoidium*), and the posterior part the *coracoidium*. In the same way the simple arch of the pelvic zone breaks up into an upper (dorsal) piece, the iliac-bone (*os ilium*), and a lower (ventral) piece; the anterior part of the latter forms the pubic bone (*os pubis*), and the posterior the ischial bone (*os ischii*).

There is also a complete agreement between the fore and hind limb in the stem or shaft. The first section of the stem is supported by a single strong bone—the humerus in the fore, the femur in the hind limb. The second section contains two bones: in front the radius (r) and ulna (u), behind the tibia and fibula. (Cf. the skeletons in Figs. 260, 265, 270, 278-283, and 338.) The succeeding numerous small bones of the wrist (*carpus*) and ankle (*tarsus*) are also similarly arranged in the fore and hind extremities, and so are the five bones of the middle-hand (*metacarpus*) and middle-foot (*metatarsus*). Finally, it is the same with the toes themselves, which have a similar characteristic composition

from a series of bony pieces before and behind. We find a complete parallel in all the parts of the fore leg and the hind leg.

When we thus learn from comparative anatomy that the skeleton of the human limbs is composed of just the same bones, put together in the same way, as the skeleton in the four higher classes of Vertebrates, we may at once infer a common descent of them from a single stem-form. This stem-form was the earliest amphibian that had five toes on each foot. It is particularly the outer parts of the limbs that have been modified by adaptation to different conditions. We need only recall the immense variations they offer within the mammal class. We have the slender legs of the deer and the strong springing legs of the kangaroo, the climbing feet of the sloth and the digging feet of the mole, the fins of the whale and the wings of the bat. It will readily be granted that these organs of locomotion differ as much in regard to size, shape, and special function as can be conceived. Nevertheless, the bony skeleton is substantially the same in every case. In the different limbs we always find the same characteristic bones in essentially the same rigidly hereditary connection, this is as splendid a proof of the theory of evolution as comparative anatomy can discover in any organ of the body. It is true that the skeleton of the limbs of the various mammals undergoes many distortions and degenerations besides the special adaptations (Fig. 342). Thus we find the first finger or the thumb atrophied in the fore-foot (or hand) of the dog (II). It has entirely disappeared in the pig (III) and tapir (V). In the ruminants (such as the ox, IV) the second and fifth toes are also atrophied, and only the third and fourth are well developed (VI, 3). Nevertheless, all these different fore-feet, as well as the hand of the ape (Fig. 340) and of man (Fig. 341), were originally developed from a common pentadactyle stem-form. This is proved by the rudiments of the degenerated toes, and by the similarity of the arrangement of the wrist-bones in all the pentanomes (Fig. 342 a-f).

If we candidly compare the bony skeleton of the human arm and hand with that of the nearest anthropoid apes, we find an almost perfect identity. This is especially true of the chimpanzee. In regard to the arrangement of the various

parts, the lowest living races of men (the Veddahs of Ceylon, Fig. 344) are midway between the chimpanzees (Fig. 343) and the European (Fig. 345). More considerable are the differences in structure and the proportions of the various parts between the different genera of anthropoid apes (Figs. 276-282); and still greater is the morphological distance between these and the lowest apes (the *Cynopithecus*). Here, again, impartial and thorough anatomic comparison confirms the accuracy of Huxley's pithcanonema principle (p. 177).

The complete unity of structure which is thus revealed by the comparative anatomy of the limbs is fully confirmed by their embryology. However different the extremities of the four-footed Craniotes may be in their adult state, they all develop from the same rudimentary structure. In every case the first trace of the limb in the embryo is a very simple protuberance that grows out of the side of the hypocoema. These simple structures develop directly into fins in the fishes and Dipneusts by differentiation of their cells. In the higher classes of Vertebrates each of the four takes the shape in its further growth of a leaf with a stalk, the inner half becoming narrower and thicker and the outer half broader and thinner. The inner half (the stalk of the leaf) then divides into two sections—the upper and lower parts of the limb. Afterwards four shallow indentations are formed at the free edge of the leaf, and gradually deepen, these are the intervals between the five toes (Fig. 274). The toes soon make their appearance. But at first all five toes, both of fore and hind feet, are connected by a thin membrane like a "swimming-web"; they remind us of the original chapping of the foot as a puddling tin. The further development of the limbs from this rudimentary structure takes place in the same way in all the Vertebrates according to the laws of heredity.

The embryonic development of the muscles, or active organs of locomotion, is not less interesting than that of the skeleton, or passive organs. But the comparative anatomy and ontogeny of the muscular system are much more difficult and inaccessible, and consequently have hitherto been less studied. We can therefore only draw some general phylogenetic conclusions therefrom.

It is incontestable that the musculature of the Vertebrates has been evolved from

that of lower Invertebrates; and among these we have to consider especially the unarticulated Vermalia. They have a simple cutaneous muscular layer, developing from the mesoderm. This was afterwards replaced by a pair of internal lateral muscles, that developed from the middle wall of the coelom-pouches; we still find the first rudiments of the muscles arising from the muscle-plate of these in the embryos of all the Vertebrates (cf. Figs. 124, 158-60, 232-4 *mp*). In the unarticulated stem-forms of the Chordonia, which we have called the Prochordonia, the two coelom-pouches, and therefore also the muscle-plates of their walls, were not yet segmented. A great advance was made in the articulation of them, as we have followed it step by step in the *Amphioxus* (Figs. 124, 158). This segmentation of the muscles was the momentous historical process with which vertebration, and the development of the vertebrate stem, began. The articulation of the skeleton came after this segmentation of the muscular system, and the two entered into very close correlation.

The epimerites or dorsal coelom-pouches of the Acrania, Cyclostomes, and Selachii (Fig. 161 A) first develop from their inner or median wall (from the cell-layer that lies directly on the skeletal plate [*sk*] and the medullary tube [*nr*]) a strong muscle-plate (*mp*). By dorsal growth (*w*) it also reaches the external wall of the coelom-pouches, and proceeds from the dorsal to the ventral wall. From these segmental muscle-plates, which are chiefly concerned in the segmentation of the Vertebrates, proceed the lateral muscles of the stem, as we find in the simplest form in the *Amphioxus* (Fig. 220). By the formation of a horizontal frontal septum they divide on each side into an upper and lower series of myotomes, dorsal and ventral lateral muscles. This is seen with typical regularity in the transverse section of the tail of a fish (Fig. 346). From these earlier lateral muscles of the trunk develop the greater part of the subsequent muscles of the trunk, and also the much later "muscular buds" of the limbs.*

* The ontogeny of the muscles is mostly recapitulative. The greater part of the motion of the head for the ventral muscles) being originally at the hypocoema of the vertebrate organism, and develop from the wall of the hypocoema or ectoderm coelom-pouches. These also appear originally in the primary muscles of the limbs, as shown in the following phylogenetically to the hypocoema. (cf. Chapter XIV.)

CHAPTER XXVII.

THE EVOLUTION OF THE ALIMENTARY SYSTEM

The chief of the vegetal organs of the human frame, to the evolution of which we now turn our attention, is the alimentary canal. The gut is the oldest of all the organs of the metazoic body, and it leads us back to the earliest age of the formation of organs—to the first section of the Laurentian period. As we have already seen, the result of the first division of labour among the homogeneous cells of the earliest multicellular animal body was the formation of an alimentary cavity. The first duty and first need of every organism is self-preservation. This is met by the functions of the nutrition and the covering of the body. When, therefore, in the primitive globular *Blastera* the homogeneous cells began to effect a division of labour, they had first to meet this twofold need. One half were converted into alimentary cells and enclosed a digestive cavity, the gut. The other half became covering cells, and formed an envelope round the alimentary tube and the whole body. Thus arose the primary germinal layers—the inner,

modifications, has everywhere essentially the same paingenic structure (Figs. 30-35). The elaborate alimentary canal of the higher animals develops ontogenetically from the same simple primitive gut of the *gastrula*.

This gastrula theory is now accepted by nearly all zoologists. It was first supported and partly modified by Professor Ray-Lankester; he proposed three years afterwards (in his essay on the development of the Molluscs, 1875) to give the name of *archenteron* to the primitive gut and *blastopore* to the primitive mouth.

Before we follow the development of the human alimentary canal in detail, it is necessary to say a word about the general features of its composition in the fully-developed man. The mature alimentary canal in man is constructed in all its main features like that of all the higher animals, and particularly resembles that of the Catarrhines, the narrow-nosed apes of the Old World. The entrance into it, the mouth, is armed with thirty-two teeth, fixed in rows in the upper and

outer, covering, or animal layer. (Cf. pp. 214-17.)

When we try to construct an animal frame of the simplest conceivable type, that has some such primitive alimentary canal and the two primary layers constituting its wall, we inevitably come to the very remarkable embryonic form of the *gastrula*, which we have found with extraordinary persistence throughout the whole range of animals, with the exception of the unicellulars—in the Sponges, Cnidaria, Platyodes, Vermalia, Molluscs, Articulata, Echinoderms, Tunicates, and Vertebrates. In all these stems the *gastrula* recurs in the same very simple form. It is certainly a remarkable fact that the *gastrula* is found in various animals as a larva-stage in their individual development, and that this *gastrula*, though much disguised by ontogenetic

tion is exactly the same as that of the Catarrhines, and differs from that of all other animals (p. 257). Above the mouth-cavity is the double nasal cavity; they are separated by the palate-wall. But we saw that this separation is not there from the first, and that originally there is a common mouth-nasal cavity in the embryo; and this is only divided afterwards by the hard palate into two—the nasal cavity above and that of the mouth below (Fig. 313).

At the back the cavity of the mouth is half closed by the vertical curtain that we call the soft palate, in the middle of which is the uvula. A glance into a mirror with the mouth wide open will show its shape. The uvula is interesting because, besides man, it is only found in the ape. At each side of the soft palate are the tonsils. Through the curved opening that we find

underneath the soft palate we penetrate into the gullet or pharynx behind the mouth-cavity. Into this opens on either side a narrow canal (the Eustachian tube), through which there is direct communication with the tympanic cavity of the ear (Fig. 320 c). The pharynx is continued in a long, narrow tube, the oesophagus (*es*). By this the food passes into the stomach when masticated and swallowed. Into the gullet also opens, right above, the trachea (*tr*), that leads to the lungs. The entrance to it is covered by the

the canal; it opens only when the pulpy food passes from the stomach into the intestine. In man and the higher Vertebrates the stomach itself is the chief organ of digestion, and is especially occupied with the solution of the food; this is not the case in many of the lower Vertebrates, which have no stomach, and discharge its function by a part of the gut farther on. The muscular wall of the stomach is comparatively thick; it has externally strong muscles that accomplish the digestive movements and internally a form

The cartilaginous epiglottis is found only in the mammals, and has developed from the fourth branchial arch of the fishes and amphibies. The lungs are found, in man and all the mammals, to the right and left in the pectoral cavity, with the heart between them. At the upper end of the trachea there is, under the epiglottis, a specially differentiated part, strengthened by a cartilaginous skeleton, the larynx. This important organ of human speech also develops from a part of the alimentary canal. In front of the larynx is the thyroid gland, which sometimes enlarges and forms goitre.

The oesophagus descends into the pectoral cavity along the vertebral column, behind the lungs and the heart, pierces the diaphragm, and enters the visceral cavity. The diaphragm is a membranous-muscular partition that completely separates the thoracic from the abdominal cavity in all the animals (and these alone). This partition is not found in the beginning; there is at first a common breast-belly cavity, the coeloma or pleuro-peritoneal cavity. The diaphragm is formed later on as a muscular horizontal partition between the thoracic and abdominal cavities. It then completely separates the two cavities, and is only pierced by several organs that pass from the one to the other. One of the chief of these organs is the oesophagus. After this has passed through the diaphragm, it expands into the gastric sac in which digestion takes place.

adult man (Fig. 340) is a long, somewhat expanding on the left into a mass of the stomach (*st*), but narrowing on the right, and passing at the pylorus (*p*) into the small intestine. At that point there is a valve, the pyloric valve (*v*), between the two sections of

glands, which secrete the gastric juice.

Next to the stomach comes the longest section of the alimentary canal, the middle gut or small intestine. Its chief function is to absorb the peptonised fluid



FIG. 340.—Human stomach and duodenum, longitudinal section. a, cardiac end of oesophagus; b, fundus (breadth of the left side); c, pylorus; d, pyloric valve; e, pyloric cavity; f, duodenum; g, gall-bladder; h, pancreatic duct. (From Meyer.)

mass of food, or the chyle, and it is subdivided into several sections, of which the (next to the stomach) is called the duodenum (Fig. 340 f, g, h). It is a short, horseshoe-shaped loop of the gut. The glands of the alimentary canal open into it—the liver, the chief digestive gland, that secretes the gall, and the pancreas, which secretes the pancreatic juice. The two glands pour their secretions, the bile and pancreatic juice, close together into the duodenum (*i*). The opening of the gall-duct is of particular phylogenetic importance, as it is the same in all the Vertebrates, and indicates the principal point of the hepato- or trunk-gut (Gegenbaur). The liver, phylogenetically older than the stomach, is a large gland, rich in blood, in the adult man, immediately under the diaphragm on the left

side, and separated by it from the lungs. The pancreas lies a little further back and more to the left. The remaining part of the small intestine is so long that it has to coil itself in many folds in order to find room in the narrow space of the abdominal cavity. It is divided into the jejunum above and the ileum below. In the last section of it is the part of the small intestine at which in the embryo the yolk-sac opens into the gut. This long and thin intestine then passes into the large intestine, from which it is cut off by a special valve. Immediately behind this "Bauhin-valve" the first part of the large intestine forms a wide, pouch-like structure, the caecum. The atrophied end of the caecum is the famous rudimentary organ, the




FIG. 330.—Head of a hermaphrodite, one-fourth of its length. (From *Metaceras*.) The deep mouth-pit (B) is separated by the head-gut (C) from the pharynx (D). The oesophagus (E) is derived from the mouth (A). The stomach (F) is derived from the mouth (A). The intestine (G) is derived from the mouth (A).

vermiform appendix. The large intestine (caecum) consists of three parts—an ascending part on the right, a transverse middle part, and a descending part on the left. The latter finally passes through an S-shaped bend into the last section of the alimentary canal, the rectum, which opens behind by the anus. Both the large and small intestines are equipped with numbers of small glands, which secrete mucous and other fluids.

For the greater part of its length the alimentary canal is attached to the inner dorsal surface of the abdominal cavity, or to the lower surface of the vertebral column. The fixing is accomplished by means of the thin membranous plate that we call the mesentery.

Although the fully-formed alimentary canal is thus a very elaborate organ, and although in detail it has a quantity of complex structure, it is not entirely

newly evolved from the very simple form of the primitive. We find in gastrea-ancestors, and that every gastrula brings before us to-day. We have already pointed out (Chapter IX.) how the epigastrula of the mammals (Fig. 67) can be reduced to the original type of the bell-gastrula, which is now preserved by the amphioxus alone (Fig. 33). Like the latter, the human gastrula and that of all other mammals must be regarded as the ontogenetic reproduction of the phylogenetic form that we call the Gastraea, in which the whole body is nothing but a double-walled gastric sac.

We already know from embryology the manner in which the gut develops in the embryo of man and the other mammals. From the gastrula is first formed the spherical embryonic vesicle filled with fluid (*gastrula*, Fig. 106). In the dorsal wall of this the sole-shaped embryonic shield is developed, and on the under-side of this a shallow groove appears in the middle line, the first trace of the later, secondary alimentary tube. The gut-groove becomes deeper and deeper, and its edges bend towards each other, and finally form a tube.

As we have seen, this simple cylindrical gut-tube is at first completely closed before and behind in man and in the Vertebrates generally (Fig. 148); the permanent openings of the alimentary canal, the mouth and anus, are only formed later on, and from the outer skin. A mouth-pit appears in the skin in front (Fig. 330 A), and this grows towards the blind fore-end of the cavity of the head-gut (A), and at length breaks into it. In the same way a shallow anus-pit is formed in the skin behind, which grows deeper and deeper, advances towards the blind hinder end of the pelvic gut, and at last connects with it. There is at first, both before and behind, a thin partition between the external cutaneous pit and the blind end of the gut—the throat-membrane in front and the anus-membrane behind; these disappear when the connection takes place.

Directly in front of the anus-opening the alimentary develops from the hind gut; this is the important embryonic structure

that forms into the placenta in One Placental (including man). In this more advanced form the human alimentary canal (and that of all the other mammals) is a slightly bent, cylindrical tube, with an opening at each end, and two appendages growing from its lower wall: the anterior one is the umbilical vesicle or yolk-sac, and the posterior the allantois or urinary sac (Fig. 195).

The thin wall of this simple alimentary tube and its ventral appendages is found, on microscopic examination, to consist of two strata of cells. The inner stratum, lining the entire cavity, consists of larger and darker cells, and is the gut-gland layer. The outer stratum consists of smaller and lighter cells, and is the gut-fibre layer. The only exception is in the cavities of the mouth and anus, because these originate from the skin. The inner coat of the mouth-cavity is not provided by the gut-gland layer, but by the skin-epithelium layer; and its muscular substratum is provided, not by the gut-fibre, but the skin-fibre, layer. It is the same with the wall of the small anus-cavity.

If it is asked how these constituent layers of the primitive gut-wall are related to the various tissues and organs that we find afterwards in the fully-developed system, the answer is very simple. It can be put in a single sentence. The epithelium of the gut—that is to say, the internal soft stratum of cells that lines the cavity of the alimentary canal and all its appendages, and is immediately occupied with the processes of nutrition—is formed solely from the gut-gland layer, all other tissues and organs that belong to the alimentary canal and its appendages originate from the gut-fibre layer. From the latter is also developed the whole of the outer envelope of the gut and its appendages; the fibrous connective tissue and the smooth muscles that compose its muscular layer, the cartilages that support it (such as the cartilages of the larynx and the trachea), the blood-vessels and lymph-vessels that absorb the nutritive fluid from the intestines—in a word, all that there is in the alimentary system besides the epithelium of the gut. From the same layer we also get the whole of the mesentery, with all the organs embedded in it—the heart, the large blood-vessels of the body, etc.

Let us now leave this original structure of the mammal gut for a moment, in

order to compare it with the alimentary canal of the lower Vertebrates, and of those Invertebrates that we have recognised as man's ancestors. We find, first of all, in the lowest Metazoa, the Gastrozooids, that the gut remains permanently in the very simple form in which we find it transiently in the paligenetic gastrula of the other animals; it is thus in the Gastrozooids (*Pennatulidæ*), the Physoczooids (*Ephyra*), the simplest Sponges (*Olynthus*), the freshwater Polyps (*Hydra*), and the aculeate embryos of many other Coelentera (Figs.



FIG. 195.—Scales or antagonistic teeth of a shark (*Chondrichthys* *acronotus*). A three-pronged tooth runs obliquely on each of the quadrangular bony plates that lie in the corners. (From Gegenbaur.)

213-218). Even in the simplest forms of the Platyodes, the Rhabdocoela (Fig. 240), the gut is still a simple straight tube, lined with the ectoderm; but with the important difference that in this case its single opening, the primitive mouth (*m*), has formed a muscular gullet (*es*) by invagination of the skin.

We have the same simple form in the gut of the lowest Vermalia (*Gastrotrocha*, Fig. 242, *Nematodes*, *Sagitta*, etc.). But in these a second important opening of the gut has been formed at the opposite end to the mouth, the anus (Fig. 243 *a*).

We see a great advance in the structure of the ventral gut in the remarkable *Balanoglossus* (Fig. 245), the sole survivor of the Enteropneust class. Here we have the first appearance of the division of the alimentary tube into two sections that characterize the Chordonia. The fore half, the head-gut (*cephalogaster*), becomes the organ of respiration (branchial gut, Fig. 245 *b*); the hind half, the trunk-gut (*truncogaster*), alone acts as digestive organ (hepatic gut, *d*). respiratory branchial gut, the posterior the digestive hepatic gut. In both it develops palliologically from the primitive gut of the gastrula, and in both the hinder end of the medullary tube covers the primitive mouth to such an extent that the remarkable medullary intestinal duct is formed, the passing communication between the neural and intestinal tubes (*canalis neuropneumaticus*, Figs. 83, 85 *nr*). In the vicinity of the closed primitive mouth, possibly in its place, the later




FIG. 245.—Gut of a lampbrush embryo, one-fourth of an inch long, magnified fifteen times. (From Hux)

The differentiation of these two parts of the gut in the Enteropneust is just the same in all the Vertebrates.

It is particularly interesting and instructive in this connection to compare the Enteropneusts with the Ascidia and the Amphioxus (Figs. 220, 210)—the remarkable animals that form the connecting link between the Invertebrates and the Vertebrates. In both forms the gut is of substantially the same construction; the anterior section forms the

anus is developed. In the same way the mouth is a fresh formation in the Amphioxus and the Ascidia.

and that the Craniotes generally. The secondary formation of the mouth in the Chordonia is probably connected with the development of the gill-slits which are formed in the gut-wall immediately behind the mouth. In this way the anterior section of the gut is converted into a respiratory organ. I have already pointed out that this modification is distinctive of the

Vertebrates and Tunicates. The phylogenetic appearance of the gill-clefts indicates the commencement of a new epoch in the stem-history of the Vertebrates.

In the further ontogenetic development of the alimentary canal in the human embryo the appearance of the gill-clefts is the most important process. At a very early stage the gullet-wall joins with the external body-wall in the head of the human embryo, and this is followed by the formation of four clefts, which lead directly into the gullet from without, on the right and left sides of the neck, behind the mouth. These are the gill or gullet clefts, and the partitions that separate them are the gill or gullet-arches (Fig. 271). These are most interesting embryonic structures. They show us that all the higher Vertebrates reproduce in their earlier stages, in harmony with the biogenetic law, the process that had so important a part in the rise of the whole Chordanus-stem. This process was the differentiation of the gut into two sections—an anterior respiratory section, the branchial gut, that was restricted to breathing, and a posterior digestive section, the hepatic gut. As we find this highly characteristic differentiation of the gut into two different sections in all the Vertebrates and all the Tunicates, we may conclude that it was also found in their common ancestors, the Prochordonia—especially as even the Enteropterygians have it. (Cf. pp. 119, 151, 227, and Figs. 210, 220, 245.) It is entirely wanting in all the other Invertebrates.

There is at first only one pair of gill-clefts in the Amphioxus, as in the Ascidia and Enteropneusta, and the Cephalopoda (Fig. 225) have only one pair throughout life. But the number presently increases in the former. In the Crustaceans, however, it decreases still further. The Cyclostomes have six to eight pairs (Fig. 247); some of the Selachii six or seven pairs, most of the fishes only four or five pairs. In the embryo of man, and the higher Vertebrates generally, where they make an appearance at an early stage, only three or four pairs are developed. In the fishes they remain throughout life, and form an exit for the water taken in at the mouth (Figs. 249-251). But they are partly lost in the amphibia, and entirely in the higher Vertebrates. In these nothing is left but a relic of the first gill-cleft. This is

formed into a part of the organ of hearing; from it are developed the external meatus, the tympanic cavity, and the Eustachian tube. We have already considered these remarkable structures, and need only point here to the interesting fact that our middle and external ear is a modified inheritance from the fishes. The branchial arches also, which separate the clefts, develop into very different parts. In the fishes they remain gill-arches, supporting the respiratory gill-leaves. It is the same with the lowest amphibia, but in the higher amphibia they undergo various



FIG. 225.

FIG. 226.

FIG. 225.—Gut of a dog-embryo (shown in Fig. 226 from *Reichert*), seen from the ventral side. a, gill-arches (four pairs); b, rudiments of pharynx and larynx; c, lumen of stomach; d, wall of the open intestine (into which the middle gut opens with a blind aperture); e, rectum.

FIG. 226.—The same gut seen from the dorsal side. a, lungs; b, stomach; c, larynx; d, intestine.

modifications, and in the three higher classes of Vertebrates (including man) the hyoid bone and the ossicles of the ear develop from them. (Cf. p. 291.)

From the first gill-arch, from the inner surface of which the muscular tongue proceeds, we get the first structure of the maxillary skeleton—the upper and lower jaws, which surround the mouth and support the teeth. These important parts are wholly wanting in the two lowest classes of Vertebrates, the Acrania and Cyclostoma. They appear first in the earliest Selachii (Figs. 248-251), and have been transmitted from this stem-group of the Gnathostomes to the higher

Vertebrates. Hence the original formation of the skeleton of the mouth can be traced to these primitive fishes, from which we have inherited it. The teeth are developed from the skin that clothes

it resembles the gut of the earliest Vermalis (*Gastrotrocha*). It then divides into two sections, a fore or buccal gut and a hind or hepatic gut, like the alimentary canal of the *Balanoglossum*,




FIG. 350.—Medial section of the head of a *Petromyzon-larva*. (From Gegenbaur.) *a* hypobranchial groove (above it is the gill slit as in the ventral openings of the seven gill-slits); *b* gill slit; *c* mouth; *d* sensory tentacle; *e* lateral fold of pharynx.

the jaws. As the whole mouth cavity originates from the outer integument (Fig. 350), the teeth also must come from it. As a fact, this is found to be the case on microscopic examination of the development and finer structure of the teeth. The scales of the fishes, especially of the shark type (Fig. 351), are in the same position as their teeth in this respect (Fig. 352). The osseous matter of the tooth (dentine) develops from the corium; its enamel covering is a secretion of the epidermis that covers the corium. It is the same with the cutaneous teeth or placoid scales of the *Selachii*. At first the whole of the mouth was armed with these cutaneous teeth in the *Selachii* and in the earliest amphibians. Afterwards the formation of them was restricted to the edges of the jaws.

Hence our human teeth are, in relation to their original source, modified fish-scales. For the same reason we must regard the salivary glands, which open into the mouth, as epidermic glands, as they are formed, not from the glandular layer of the gut like the rest of the alimentary glands, but from the epidermis, from the horny plate of the outer germinal layer. Naturally, in harmony with this evolution of the mouth, the salivary glands belong genetically to one series with the sudoriferous, sebaceous, and mammary glands.

Thus the human alimentary canal is as simple as the primitive gut of the gastrula in its original structure. Later

it resembles the gut of the earliest Vermalis (*Gastrotrocha*). It then divides into two sections, a fore or buccal gut and a hind or hepatic gut, like the alimentary canal of the *Balanoglossum*, the *Ascidia*, and the *Amphioxus*. The formation of the jaws and the branchial arches changes it into a real fish-gut (*Selachii*). But the branchial gut, the one reminiscence of our fish-ancestors, is afterwards atrophied as such. The parts of it that remain are converted into entirely different structures.

But, although the anterior section of our alimentary canal thus entirely loses its original character of branchial gut, it retains the physiological character of respiratory gut. We are now astonished to find that the permanent respiratory organ of the higher Vertebrates, the air-breathing lung, is developed from this first part of the alimentary canal. Our lungs, tracheae, and larynx are formed from the ventral wall of the branchial gut. The whole of the respiratory apparatus, which occupies the greater part of the pectoral cavity in the adult man, is at first merely a small pair of vesicles or sacs, which grow out of the floor of the head-



FIG. 351.—Transverse section of the head of a *Petromyzon-larva*. (From Gegenbaur.) Beneath the pharynx (*d*) we see the hypobranchial groove, above it the chorda and neural tube. *A*, *B*, *C* stages of development.

gut immediately behind the gills (Fig. 354 *c*, 147 *f*). These vesicles are found in all the Vertebrates except the two lowest classes, the *Acromia* and *Cyclostomes*. In the lower Vertebrates they do not develop

193), but into a large air-filled bladder, which occupies a good deal of the body-cavity and has a quite different purpose. It serves, not for breathing but to effect swimming movements up and down, and so is a sort of hydrostatic apparatus—the floating bladder of fishes (*Neohydrus*, p. 233). However, the human lungs, and those of all air-breathing Vertebrates, develop from the same simple vesicular appendage of the head-gut that becomes the floating bladder in the fishes.

At first this bladder has no respiratory function, but merely acts as hydrostatic apparatus for the purpose of increasing or lessening the specific gravity of the body. The fishes, which have a fully-developed floating bladder, can press it together, and thus condense the air it contains. The air also escapes sometimes from the alimentary canal, through an air-duct that connects the floating bladder with the pharynx, and is ejected by the mouth. This lessens the size of the bladder, and so the fish becomes heavier and sinks. When it wishes to rise again, the bladder is expanded by relaxing the pressure. In many of the Crocodontory, the wall of the bladder is covered with bony plates as in the *Trinacrus Undatus* (Fig. 254).

This hydrostatic apparatus begins the Dipneusts to change into a respiratory organ; the blood-vessels in the bladder now no longer mere air themselves, but also take air through the air-duct. This process reaches its full development in the Amphibia. In these the floating bladder has turned into lungs, and the air-passages into a trachea. The lungs of the Amphibia have been transmitted to the three higher classes of Vertebrates. In the lowest Amphibia the lungs on either side are still very simple transparent sacs with thin walls, as in the common Salamander, the Triton. It still entirely resembles the floating bladder of the fishes. It is true that the Amphibia have two lungs, right and left. But the floating bladder is also double in many of the fishes (such as the early Ganoids), and divides into right and left halves. On the other hand, the lung is single in *Ceratodus* (Fig. 257).

In the human embryo and that of all the other Amniotes the lungs develop from the hind part of the ventral wall of the head-gut (Fig. 199). Immediately behind the single structure of the thyroid

gland a median groove, the rudiment of a trachea, is detached from the gullet. From its hinder end a couple of vesicles develop—the simple tubular rudiments of the right and left lungs. They afterwards increase considerably in size, fill the greater part of the thoracic cavity, and take the heart between them. In the frogs we find that the simple sac has developed into a spongy body of peculiar froth-like tissue. The originally short connection of the pulmonary sacs with the head-gut extends into a long, thin tube. This is the wind-pipe (trachea); it opens into the gullet above, and divides below into two branches which go to the two lungs. In the wall of the trachea



FIG. 257.—Thoracic and abdominal viscera of a human embryo of twelve weeks, natural size. (From KÄNDLER.) The head is omitted. Ventral and mesal walls are removed. The greater part of the body-cavity is taken up with the lungs, from the middle part of which the coronary and the venous appendages protrude. Above the diaphragm, in the middle, is the second heart; to the right and left of it are the two small lungs.

circular cartilages develop, and these keep it open. At its upper end, underneath the pharyngeal opening, the larynx is formed—the organ of voice and speech. The larynx is found at various stages of development in the Amphibia, and continues in the same position to trace the progressive growth of this important organ from the rudimentary structure of the lower Amphibia up to the elaborate and delicate vocal apparatus in the larynx of man and of the birds.

We refer here to an interesting rudimentary organ of the respiratory gut, the thyroid gland, the large gland in front of the larynx, that lies below the "Adam's

apple," and is often especially developed in the male sex. It has a certain function—not yet fully understood—in the nutrition of the body, and arises in the embryo by constriction from the lower wall of the pharynx. In many mining districts the thyroid gland is peculiarly liable to morbid enlargement, and then forms goitre, a growth that hangs at the front of the neck. But it is much more interesting phylogenetically. As Wilhelm Müller, of Jena, has shown, this rudimentary organ is the last relic of the hypobranchial groove, which we considered in a previous chapter, and which runs in the middle line of the gill-plate in the Ascidia and Amphioxus, and conveys food to the stomach. (Cf p. 184, Fig. 246). We still find it in its original character in the larvae of the Cyclostomes (Figs. 355 and 356).

The second section of the alimentary canal, the trunk or hepatic gut, undergoes not less important modifications among our vertebrate ancestors than the first section. In tracing the further development of this digestive part of the gut, we find that most complex and elaborate organs originate from a very rudimentary original structure. For clearness we may divide the digestive gut into three sections: the fore gut (with oesophagus and stomach), the middle gut (duodenum, with liver, pancreas, jejunum, and ileum), and the hind gut (colon and rectum). Here again we find vesicular growths or appendages of the originally simple gut developing into a variety of organs. Two of these embryonic structures, the yolk-sac and allantois, are already known to us. The two large glands that open into the duodenum, the liver and pancreas, are growths from the middle and most important part of the trunk-gut.

Immediately behind the vesicular rudiments of the lungs comes the section of the alimentary canal that forms the stomach (Figs. 353 d, 354 b). This sac-shaped organ, which is chiefly responsible for the solution and digestion of the food, has not in the lower Vertebrates the great physiological importance and the complex character that it has in the higher. In the Acrania and Cyclostomes and the earlier fishes we can scarcely distinguish a real stomach; it is represented merely by the short piece from the branchial to the hepatic gut. In some of the other fishes also the stomach is only a very simple spindle-shaped enlargement at the beginning of the digestive section

of the gut, running straight from front to back in the median plane of the body, underneath the vertebral column. In the mammals its first structure is just as rudimentary as it is permanently in the preceding. But its various parts soon begin to develop. As the left side of the spindle-shaped sac grows much more quickly than the right, and as it turns considerably on its axis at the same time, it soon comes to be obliquely. The upper end is more to the left, and the lower end more to the right. The foremost end draws up into the longer and narrower canal of the oesophagus. Underneath this on the left the blind sac (fundus) of the stomach bulges out, and thus the later form gradually develops (Figs. 349, 384 c). The original longitudinal axis becomes oblique, sinking below to the left and rising to the right, and approaches nearer and nearer to a transverse position. In the outer layer of the stomach-wall the powerful muscles that accomplish the digestive movements develop from the gut-fibre layer. In the inner layer a number of small glandular tubes are formed from the gut-gland layer; these are the peptic glands that secrete the gastric juice. At the lower end of the gastric sac is developed the valve that separates it from the duodenum (the pylorus, Fig. 349 d).

Underneath the stomach there now develops the disproportionately long stretch of the small intestine. The development of this section is very simple, and consists essentially in an extremely rapid and considerable growth lengthways. It is at first very short, quite straight, and simple. But immediately behind the stomach we find at an early stage a horseshoe-shaped bend and loop of the gut, in connection with the severance of the alimentary canal from the yolk-sac and the development of the first mesentery. The thin delicate membrane that fastens this loop to the ventral side of the vertebral column, and fills the inner bend of the horseshoe formation, is the first rudiment of the mesentery (Fig. 347 g). We find at an early stage a considerable growth of the small intestine; it is thus forced to coil itself in a number of loops. The various sections that we have to distinguish in it are differentiated in a very simple way—the duodenum (next to the stomach), the succeeding long jejunum, and the last section of the small intestine, the ileum.

From the duodenum are developed the two large glands that we have already mentioned—the liver and pancreas. The liver appears first in the shape of two small sacs, that are found to the right and left immediately behind the stomach (Figs. 353 f, 354 c). In many of the lower Vertebrates they remain separate for a long time (in the *Myxinoidei* throughout life), or are only imperfectly joined. In the higher Vertebrates they soon blend more or less completely to form a single large organ. The growth of the liver is very brisk at first. In the human embryo it grows so much in the second month of development that in the third it occupies by far the greater part of the body-cavity (Fig. 357). At first the two halves develop equally; afterwards the left falls far behind the right. In consequence of the unsymmetrical development and turning of the stomach and other abdominal viscera, the whole liver is now pushed to the right side. Although the liver does not afterwards grow so disproportionately, it is comparatively larger in the embryo at the end of pregnancy than in the adult. Its weight relatively to that of the whole body is 1 : 35 in the adult, and 1 : 18 in the embryo. Hence it is very important physiologically during embryonic life, it is chiefly concerned in the formation of blood, not so much in the secretion of bile.

Immediately behind the liver a second large visceral gland develops from the duodenum, the pancreas or sweetbread. It is wanting in most of the lowest classes of Vertebrates, and is first found in the fishes. This organ is also an outgrowth from the gut.

The last section of the alimentary canal, the large intestine, is at first in the embryo a very simple, short, and straight tube, which opens behind by the anus. It remains thus throughout life in the lower Vertebrates. But it grows considerably in the mammals, coils into various folds, and divides into two sections, the first and longer of which is the colon, and the second the rectum. At the beginning of the colon there is a valve (*valve Bauhinii*) that separates it from the small intestine. Immediately behind this there is a sac-like growth, which enlarges into the caecum (Fig. 357 v). In the plant-eating mammals this is very large, but it

is very small or completely atrophied in the flesh-eaters. In man, and most of the apes, only the first portion of the caecum is wide; the blind end-part of it is very narrow, and seems later to be merely a useless appendage of the former. This "verruiform appendage" is very interesting as a rudimentary organ. The only significance of it in man is that not infrequently a cherry-stone or some other hard and indigestible matter penetrates into its narrow cavity, and by setting up inflammation and suppuration causes the death of otherwise sound men. Teleology has great difficulty in giving a rational explanation of, and attributing to a beneficent Providence, thus dreaded appendicitis. In our plant-eating ancestors this rudimentary organ was much larger and had a useful function.

Finally, we have important appendages of the alimentary tube in the bladder and urethra, which belong to the alimentary system. These urinary organs, acting as reservoir and duct for the urine excreted by the kidneys, originate from the innermost part of the allantoic pedicle. In the Dipneusts and Amphibia, in which the allantoic sac first makes its appearance, it remains within the body-cavity, and functions entirely as bladder. But in all the Amniotes it grows far outside of the body-cavity of the embryo, and forms the large embryonic "primitive bladder," from which the placenta develops in the higher mammals. This is lost at birth. But the long stalk or pedicle of the allantois remains, and forms with its upper part the middle vesico-umbilical ligament, a rudimentary organ that goes in the shape of a solid string from the venter of the bladder to the navel. The lowest part of the allantoic pedicle (or the "urachus") remains hollow, and forms the bladder. At first this opens into the last section of the gut in man as in the lower Vertebrates, thus there is a real cloaca, which takes off both urine and excrements. But among the mammals this cloaca is only permanent in the Monotremes, as it is in all the birds, reptiles, and amphibia. In all the other mammals (marsupials and placentals) a transverse partition is afterwards formed, and this separates the urogenital aperture in front from the anus-opening behind. (Cf. p. 249 and Chapter XXIX.)

CHAPTER XXVIII.

EVOLUTION OF THE VASCULAR SYSTEM

THE use that we have hitherto made of our biogenetic law will give the reader an idea how far we may trust its guidance in phylogenetic investigation. This differs considerably in the various systems of organs; the reason is that heredity and variability have a very different range in these systems. While some of them faithfully preserve the original palingeneic development inherited from earlier animal ancestors, others show little trace of this rigid heredity; they are rather disposed to follow new and divergent ontogenetic lines of development in consequence of adaptation. The organs of the first kind represent the *conservative* element in the multicellular state of the human frame, while the latter represent the *progressive* element. The course of historic development is a result of the correlation of the two tendencies, and they must be carefully distinguished.

There is perhaps no other system of organs in the human body in which this is more necessary than in that of which we are now going to consider the obscure development—the vascular system, or apparatus of circulation. If we were to draw our conclusions as to the original features in our earlier animal ancestors solely from the phenomena of the development of this system in the embryo of man and the other higher Vertebrates, we should be wholly misled. By a number of important embryonic adaptations, the chief of which is the formation of an extensive food-yell, the original course of the development of the vascular system has been so much falsified and curtailed in higher Vertebrates that I

nothing now remains in their embryology of some of the principal phylogenetic features. We should be quite unable to explain these if comparative anatomy and ontogeny did not come to our assistance,

an and all the apparatus of

cavities filled with juices or cell-containing fluids. These "vessels" (*vascula*) play an important part in the nutrition of the body. They partly conduct the nutritive red blood to the various parts of the body (blood-vessels); partly absorb from the gut the white chyle formed in digestion (chyle-vessels); and partly collect the used-up juices and convey them away from the tissues (lymphatic vessels). With the latter are connected the large cavities of the body, especially the body-cavity, or coeloma. The lymphatic vessels conduct both the colourless lymph and the white chyle into the venous part of the circulation. The lymphatic glands act as producers of new blood-cells, and with them is associated the spleen. The centre of movement for the circulation of the fluids is the heart, a strong muscular sac, which contracts regularly and is equipped with valves like a pump. This constant and steady circulation of the blood makes possible the complex metabolism of the higher animals.

But, however important the vascular system may be to the more advanced and larger and highly-differentiated animals, it is not at all an indispensable element of animal life as is commonly supposed. The older science of medicine regarded the blood as the real source of life. Even in the still prevalent confused notions of heredity the blood plays the chief part. People speak generally of full blood, half blood, etc., and imagine that the hereditary transmission of certain characters "lies in the blood." The incorrectness of these ideas is clearly seen from the fact that in the act of generation the blood of the parents is not directly transmitted to the offspring, nor does the embryo possess blood in its early stages. We have already seen that not only the differentiation of the four secondary germinal layers, but also the first structures of the principal organs in the embryo of all the Vertebrates, takes place long before there is any

trace of the vascular system—the heart and the blood. In accordance with this ontogenetic fact, we must regard the vascular system as one of the latest organs from the phylogenetic point of view, just as we have found the alimentary canal to be one of the ear-

liest. The vascular system is much later than the alimentary.

The important nutritive fluid that circulates as blood and lymph in the elaborate canals of our vascular system is not a clear, simple fluid, but a very complex chemical juice with millions of

The red colour of the blood is caused by the great accumulation of the former, the others circulate among them in a colourless state. When the colourless cells increase at the expense of the red cells (or chlorocysts).

The lymph-cells (*leucocytes*), commonly called the "white corpuscles" of the blood, are phylogenetically older and more widely distributed in the animal world than the red. The great majority of the invertebrates that have acquired an independent vascular system have only colourless lymph-cells in the circulation.



FIG. 358



FIG. 359

FIG. 358.—Red blood-cells of various Vertebrates (equally magnified). 1 of man 1 tenth, 3 down, 4, proteus 5 water-snail (3 times), 6 frog, 7 rabbit (3 times), 8 kangaroo (3 times), 9 salmon (3 times), 10 salmon (3 times). (From Wagner)

A capillary from the mammary vascular system.

cells floating in it. These blood-cells are just as important in the complicated life of the higher animal body as the circulation of money is to the commerce of a civilized community. Just as the citizens meet their needs most conveniently by means of a financial circulation, so the various tissue-cells, the microscopic citizens of the multicellular human body, have their food conveyed to them best by the circulating cells in the blood. These blood cells (*hemocytes*) are of two kinds in man and all the other Craniotes—red cells (*erythrocytes* or *erythrocytes*) and colourless or lymph cells (*leucocytes*).

fluid. There is an exception in the Nemertines (Fig. 358) and some groups of Annelids. When we examine the colourless blood of a cray-fish or a snail (Fig. 358) under a high power of the microscope, we find in each drop numbers of mobile leucocytes, which behave just like independent Amoebae (Fig. 17). Like these uncellular Protozoa, the colourless blood-cells creep slowly about, their unshapely plasma-body constantly changing its form, and stretching out finger-like processes first in one direction, then another. Like the Amoebae, they take particles into their cell-body. On account

of this feature these amoeboid plastids are called "eating cells" (*phagocytes*), and on account of their motions "travelling cells" (*leucocytes*). It has been shown by the discoveries of the last few decades that these leucocytes are of the greatest physiological and pathological consequence to the organism. They can absorb either solid or dissolved particles from the wall of the gut, and convey them to the blood in the chyle; they can absorb and remove unusable matter from the tissues. When they pass in large quantities through the fine pores of the capillaries and accumulate at irritated spots, they cause inflammation. They can consume and destroy bacteria, the dreaded vehicles of infectious diseases; but they can also transport these injurious Monera to fresh regions, and so extend the sphere

of globs) is regularly distributed in the pores of their protoplasm. The red cells of most of the Vertebrates are elliptical flat disks, and enclose a nucleus of the same shape; they differ a good deal in size (Fig. 358). The mammals are distinguished from the other Vertebrates by the circular form of their biconcave red cells and by the absence of a nucleus (Fig. 1); only a few genera still have the elliptic form inherited from the reptiles (Fig. 2). In the embryos of the mammals the red cells have a nucleus and the power of increasing by cleavage (Fig. 10).

The origin of the blood-cells and vessels in the embryo, and their relation to the germinal layers and tissues, are among the most difficult problems of ontogeny—those obscure questions on which the most divergent opinions are still advanced




FIG. 360.—Transverse section of the trunk of a chick-embryo, forty-five hours old. (From *Reimer*.) A ectoderm (outer layer), P epidermis, HZ protonal duct, PP embryonic cavity, Se somatopleuric layer, Sp gut-tube layer, S blood-vessels in interior, an primitive aorta, containing red blood-cells.

of infection. It is probable that the sensitive and travelling leucocytes of our invertebrate ancestors have powerfully co-operated for millions of years in the phylogenesis of the advancing animal organization.

The red blood-cells have a much more restricted sphere of distribution and activity. But they also are very important in connection with certain functions of the craniote-organism, especially the exchange of gases or respiration. The cells of the dark red, carbonised or venous blood, which have absorbed carbonic acid from the animal tissues, give this off in the respiratory organs; they receive instead of it fresh oxygen, and thus bring about the bright red colour that distinguishes oxydised or arterial blood. The red colouring matter of the blood (*hemo-*

by the most competent scientists. In general, it is certain that the greater part of the cells that compose the vessels and their contents come from the mesoderm—in fact, from the gut-tube layer; it was on this account that Boer gave the name of "vascular layer" to this visceral layer of the coeloma. But other important observers say that a part of these cells come from other germinal layers, especially from the gut-gland layer. It seems to be true that blood-cells may be formed from the cells of the endoderm before the development of the mesoderm. If we examine sections of chickens, the earliest and most familiar subjects of embryology, we find at an early stage the "primitive aorta" we have already described (Fig. 360 so) in the ventral angle between the epinotum (*Pe*) and hypocoelum (*Sp*). The

thin wall of these first vessels of the primitive embryo consists of flat cells (*endothelium* or *vascular epithelium*); the fluid within already contains numbers of red blood-cells; both have been developed from the gut-fibre layer. It is the same with the vessels of the germinal area (Fig. 36 v), which lie on the entodermic membrane of the yolk-sac (*e*). These features are seen still more clearly in the transverse section of the duck-embryo in Fig. 152 (p. 141). In this we see clearly how a number of stellate cells proceed from the "vascular layer" and spread in all directions in the "primary body-cavity" — *r e*, in the spaces between the germinal layers. A part of these travelling cells come together and line the wall of the larger spaces, and thus form the first vessels; others enter into the cavity, live in the fluid that fills it, and multiply by cleavage—the first blood-cells.

But, besides these mesodermic cells of the "vascular layer" proper, other travelling cells, of which the origin and purpose are still obscure, take part in the formation of blood in the meroblastic Vertebrates (especially fishes). The chief of these are those that Rucker has most aptly designated "merocytes." These "eating yolk-cells" are found in large numbers in the food-yolk of the *Salpichthys*, especially in the yolk-wall—the border zone of the germinal disk in which the embryonic vascular net is first developed. The nuclei of the merocytes become ten times as large as the ordinary cell-nucleus, and are distinguished by their strong capacity for taking colour, or their special richness in chromatin. Their protoplasmic body resembles the stellate cells of osseous tissue (astrocytes), and behaves just like a rhizopod (such as *Gromia*); it sends out numbers of stellate processes all round, which ramify and stretch into the surrounding food-yolk. These variable and very mobile processes, the pseudopodia of the merocytes, serve both for locomotion and for getting food; as in the real rhizopoda, they surround the solid particles of food (granules and plates of yolk), and accumulate round their nucleus (the food they have received and digested). Hence we may regard them both as eating-cells

(*phagocytes*) and travelling-cells (*planocytes*). Their lively nucleus divides quickly and often repeatedly, so that a number of new nuclei are formed in a short time; as each fresh nucleus surrounds itself with a mantle of protoplasm, it provides a new cell for the construction of the embryo. Their origin is still much disputed.

Half of the twelve stems of the animal world have no blood-vessels. They make their first appearance in the Vermalia. Their earliest source is the primary body-cavity, the simple space between the two primary germinal layers, which is either a relic of the segmentation-cavity, or is a subsequent formation. Amoeboid planocytes




FIG. 36.—Merocytes of a shark-embryo. Amoeboid-like yolk-cells underneath the embryonic cavity (*B*). (From Rucker.) *a* = leucocyte-like cells; *b* nuclei of the merocytes, which wander about in the yolk and eat small yolk-plates (*d*); *c* smaller, more superficial, lighter nuclei; *e* a deeper nucleus, in the act of cleavage; *f* chromatin-filled border-nucleus, freed from the surrounding yolk in order to show the numerous pseudopodia of the protoplasmic cell-body.

cytes, which migrate from the entoderm and reach this fluid-filled primary cavity, live and multiply there, and form the first colourless blood-cells. We find the vascular system in this very simple form to-day in the Bryozoa, Rotatoria, Nemertoda, and other lower Vermalia.

The first step in the improvement of this primitive vascular system is the formation of larger canals or blood-conducting tubes. The spaces filled with blood, the relics of the primary body-cavity, receive a special wall. "Blood-vessels" of this kind (in the narrower sense) are found among the higher forms in various forms, sometimes very simple, at other times very complex. The forms

that was probably the incipient structure of the elaborate vascular system of the Vertebrates (and of the Articulates) is found in two primordial principal vessels—a dorsal vessel in the middle line of the dorsal wall of the gut, and a ventral vessel that

middle line of li
ventral v

dorsal vessel is evolved the aorta (or principal artery), from the ventral vessel the principal or subintestinal vein. The two vessels are connected in front and behind by a loop that runs round the gut. The blood contained in the two tubes is propelled by their peristaltic contractions.

The earliest Vermalin in which we first find this independent vascular

the Nemertina (Fig. 244). As a rule, they have three parallel longitudinal vessels connected by loops, a single dorsal vessel above the of lat.

right and left. re of the Nemertina the blood is already coloured, and the red colouring matter is real hæmoglobin, connected with elliptical discoid cells, as in the Vertebrates. The further evolution of this rudimentary vascular system can be gathered from the class of the Annelids in which

the Vascular system of Annelids (See next p. for peristaltic action of dorsal vessel, & ventral vessel, & transverse connections of two (enlarged in shape of heart). The below indicates the direction of the flow of blood. (From Gegenbaur.)

we find it at various stages of development. First, a number of transverse connections are formed between the dorsal and ventral vessels, which pass round the gut ring-wise (Fig. 362). Other vessels grow into the body-wall and ramify in order to convey blood to it. In addition to the two large vessels of the middle plane there are often two lateral vessels, one to the right and one to the left; as, for instance, in the leech. There are four of these parallel longitudinal vessels in the Enteropneusta (*Balanoglossus*, Fig. 245). In these important Vermalin the foremost section of the gut

has already been converted into a gill-plate, and the vascular arches that rise in the wall of this from the dorsal vessel have become branchial vessels.

We have a further important advance in the Tunicates, which we have recognised as the nearest blood-relatives of our early vertebrate ancestors.

central heart—i.e., a central organ of circulation, driving the blood into the vessels by the regular contractions of its muscular wall. It is of a very rudimentary character, a spindle-shaped tube, passing at both ends into a principal vessel (Fig. 221). By its original position behind the gill-

s (sometimes 2, sometimes less, forward), the heart is clearly that it has been formed by local enlargement of a section of the tubular vessel. We have already noticed remarkable alternation of the direction of the blood stream, the heart driving

190). This is very instructive, because most of the worms (even the Enteropneusts) the blood in the dorsal vessel flows from back to front, but in the vertebrates in the opposite direction. As a Axillia-heart alternates steadily from its direction to the other, it shows us permanently, in a sense, the phylogenetic position from the earlier forward direction of the dorsal current (in the worms) the new backward direction (in the vertebrates).




FIG. 363.—Head of a fish-embryo, with rudimentary vascular system, from the left. *ac* Carotid duct (structure of the anterior and posterior reversed vessel, or carotid stem (enlarged end of C) of the gill-chains (arterial arches between), *ad* aorta, *v* ventral artery, *v* ventral vet. (From Gegenbaur.)

vessels that proceed from either end of the tubular heart acquired a fixed function.

The foremost section of the ventral vessel beneath always conveys blood from the heart, and so acts as an artery; the hind section of the same vessel brings the blood from the body to the heart, and so becomes a vein. In view of their relation to the two sections of the gut, we may call the latter the intestinal vein and the former the branchial artery. The blood contained in both vessels, and also in the heart, is venous or carbonised blood—*s.e.*, rich in carbonic acid; on the other hand, the blood that is pumped into the gills into the dorsal vessel is provided with fresh oxygen—arterial or oxygenated blood. The finest branches of the arteries

blood gathers in a ventral vessel under the gut (intestinal vein), and goes back to the gills. A number of branchial vascular arches, which effect respiration and rise in the wall of the branchial gut from belly to back, absorb oxygen from the water. They get rid of carbonic acid; they combine with the dorsal vessel. As the evolution of the ventral vessel, which also opens the heart in the Craniotes, has developed in the Acoel into a simple tubular heart, we may regard the evolution of this in the Amphioxus as a result of degeneration, a return in this case to an earlier form of the vascular system, *s.*




FIG. 354.

FIG. 356.

rules of the Craniotes (Fig. 355) in their original disposition—*a* arterial cross-
ary (foremost continuation of the roots of the aorta). (From Kolliker.)

FIG. 354.—The five arterial arches of the birds; the lighter parts of the structure disappear, only the shaded parts remain. Letters as in Fig. 356. *a* subclavian artery, *b* pulmonary artery, *c* branches of same, *d* outer aorta, *e* inner aorta. (From Kolliker.)

FIG. 356.—The five arterial arches of mammals; letters as in Fig. 354. *a* ventral artery, *b* Botall's duct (open in the embryo, closed afterwards). (From Kolliker.)

and veins pass into each other in the tissues by means of a network of very fine, ventral, hair-like vessels, or capillaries (Fig. 359).

When we turn from the Tunicates to the closely-related Amphioxus we are astonished at first to find an apparent retrogression in the formation of the vascular system. As we have seen, the Amphioxus has no real heart; its colourless blood is driven along in its vascular system by the principal vessel itself, which contracts regularly in its whole length (cf. Fig. 310). A dorsal vessel that lies above the gut (aorta) receives the arterial blood from the gills and drives it into the body. Returning from here,

find it in many of the Craniotes we assume that the arterial system really belongs to our ancestral Craniotes and not to the Tunicates, but inherited the one-chambered heart of the Prochordonia, and transmitted it directly to the earliest Craniotes (cf. the Ideal Primitive Vertebrate, *Protophyllus*, Figs. 93-102).

The further phylogenetic evolution of the vascular system is revealed to us by the comparative anatomy of the Craniotes. At the lowest stage of this group, in the Cyclostomes, we find for the first time the differentiation of the vasculature into two sections: a system of blood-vessels proper, which convey the red blood about the body, and a system of lymphatic vessels,

receiving the venous blood from the body. The left auricle receives the arterial blood from the pulmonary veins. The two auricles have a common opening into the single ventricle,



FIG. 371




FIG. 372

FIG. 371.—Heart of a rabbit, anterior view. From behind, a pulmonary vein, a branch of the heart, a branch of ventricle, a branch of the heart, a branch of the heart, a branch of the heart. (From Rieckhoff)

FIG. 372.—Heart of the same embryo (FIG. 371) from the front. a, ventricle, b, auricle, c, arterial bulb, d, left ventricle, e, right ventricle, f, arterial bulb. (From Rieckhoff)

where the two kinds of blood

are mixed. From the last arterial arches the pulmonary arteries arise (FIG. 365 g). These force a part of the mixed blood into the lungs, the other part of it going through the aorta into the body.

From the Dipneusts upwards we now trace a progressive development of the vascular system, which ends finally with the loss of branchial respiration and a complete separation of the two halves of the circulation. In the Amphibia the partition between the two auricles is complete. In their earlier stages, as tadpoles (FIG. 261), they have still the branchial respiration and the circulation of the fishes, and their heart contains venous blood alone. Afterwards the lungs and pulmonary vessels are developed, and henceforth the ventricle of the heart contains mixed blood. In the reptiles the ventricle and its arterial cone begin to divide into two halves by a longitudinal partition, and this partition becomes complete in the higher reptiles and birds on the one hand, and the stem-forms of the mammals on the other. Henceforth, the right half of the heart contains only venous, and the left half only arterial, blood, as we find in all birds and mammals. The right auricle receives its carbonised or venous blood from the veins of the body, and the right ventricle drives it

through the pulmonary arteries into the lungs. From here the blood returns, as oxygenated or arterial blood, through the pulmonary veins to the left auricle, and is forced by the left ventricle into the arteries of the body. Between the pulmonary arteries and veins is the capillary system of the small or pulmonary circulation. Between the body-arteries and veins is the capillary system of the large or body-circulation. It is only in the two highest classes of Vertebrates—the birds and mammals—that we find a complete division of the circulations. Moreover, this complete separation has been developed quite independently in the two classes, as the dissimilar formation of the aorta shows of itself. In the birds the right half of the fourth arterial arch has become the permanent arch (FIG. 365). In the mammals this has been developed from the left half of the same fourth arch (FIG. 366).

If we compare the fully-developed arterial system of the various classes of animals, we find a good deal of variety, yet it always proceeds from the same fundamental type. Its development



FIG. 373



FIG. 374

FIG. 373.—Heart and head of a dog, anterior view. a, fore brain, b, middle brain, c, hind brain, d, primitive upper jaw, e, gill-arch, f, left auricle, g, left ventricle, h, right ventricle. (From Rieckhoff)

FIG. 374.—Heart of the same embryo, from behind. a, aorta, b, c, d, e, f, g, h, i, j, k, l, m, n, o, p, q, r, s, t, u, v, w, x, y, z. (From Rieckhoff)

just the same in man as in the other mammals; in particular, the modification of the six pairs of arterial arches is the same in both (FIGS. 367-370). At first there is only a single pair of arches, which

lie on the inner surface of the first pair of gill-arches. Behind this there then develop a second and third pair of arches (lying on the inner side of the second and third gill-arches, Fig. 367). Finally, we



FIG. 375.—Heart of a human embryo, four weeks old, *a*, front view, *b*, back view, *c*, upper half of the atrium removed, *d*, left auricle, *e*, right auricle, *f*, left ventricle, *g*, right ventricle, *h*, arterial bulb, *i*, superior vena cava, *j*, inferior vena cava, *k*, rudiment of the interventricular wall. (From Kolliker.)

FIG. 376.—Heart of a human embryo, six weeks old, front view, *a*, right ventricle, *b*, left ventricle, *c*, furrow between ventricles, *d*, arterial bulb, *e*, furrow on its surface, to right and left are the two large sinuses. (From Kolliker.)

FIG. 377.—Heart of a human embryo, eight weeks old, back view, *a*, left auricle, *b*, right auricle, *c*, left ventricle, *d*, right ventricle, *e*, right superior vena cava, *f*, inferior vena cava. (From Kolliker.)

get a fourth, fifth, and sixth pair. Of the six primitive arterial arches of the Annelids three soon pass away (the first, second, and fifth), of the remaining three, the third gives the carotids, the fourth the aorta, and the sixth (number 5 in Figs. 364 and 368) the pulmonary arteries.

The human heart also develops in just the same way as that of the other mammals (Fig. 378). We have already seen the first rudiments of its embryology, which in the main corresponds to its phylogeny (Figs. 202, 203). We saw that the palæogenetic form of the heart is a spindle-shaped thickening of the gut-tube layer in the ventral wall of the head-gut. The structure is then hollowed out, forms a simple tube, detached from its place of origin, and henceforth lies freely in the coelomic cavity. Presently the tube bends into the shape of an S, and turns spirally on an imaginary axis in such a way that

the hind part comes to lie on the dorsal surface of the fore part. The united vitelline veins open into the posterior end. From the anterior end spring the aortic arches.

This first structure of the human heart, enclosing a very simple cavity, corresponds to the tunicate-heart, and is a reproduction of that of the Prochordonia, but it now divides into two, and subsequently into three, compartments; this reminds us for a time of the heart of the Cyclostomes and fishes. The spiral turning and bending of the heart increases, and at the same time two transverse constrictions appear, dividing it externally into three sections (Figs. 371, 372). The foremost section, which is turned towards the ventral side, and from which the aortic arches rise, reproduces the auricular half of the Selachii. The middle section is a simple ventricle, and the hindmost, the section turned towards the dorsal side, into which the vitelline veins insinuate, is a simple auricle (or *sinus*). The latter forms, like the simple atrium of the fish-heart, a pair of lateral dilatations, the




FIG. 378.—Heart of the adult man, fully developed, front view, natural position, *a*, right auricle (underneath at the right ventricle), *b*, left auricle (underneath at the left ventricle), *c*, superior vena cava, *d*, pulmonary veins, *e*, pulmonary artery, *f*, Botalli's duct, *g*, aorta. (From Kolliker.)

auricles (Fig. 371 *b*); and the constriction between the atrium and ventricle is called the auricular canal (Fig. 371 *an*). The heart of the human embryo is now a complete fish-heart.

In perfect harmony with its phylogeny, the embryonic development of the human heart shows a gradual transition from the fish-heart, through the amphibian and reptile, to the mammal form. The most important point in the transition is the formation of a longitudinal

afterwards c

of the heart into right (venous) and left (arterial) halves (cf. Figs. 373-378). The separate right and left half, each of which absorbs squodling auricle, into the right auricle open the body-veins (upper and lower vena cava, Figs. 375 c, 377 c); the left auricle receives the pulmonary veins. In superficial ventricular

The heart of all the Vertebrates belongs originally to the hypostoma of the head, and we accordingly find it in the embryo of man and all the other Amniotes right in front on the under-side of the head; just as in the fishes it remains permanently in front of the gullet. It afterwards

unlike, with the a

segment of the breast, and at last reaches the breast, between the two lungs. At first it is symmetrically in the middle of the body, so that its long axis corresponds

tion. But in the apes the axis begins to be oblique, and the apex of the heart to move towards the left side. The dis-




FIG. 376.—Transverse section of the back of the head of a chick-embryo, forty hours old. (From Kollmer.) m, mouth; ap, apical cavity (head-part); h, heart; aa, aorta; sp, spinal process; d, dorsal cavity (head-part); ap, apical cavity (head-part); aa, aorta; sp, spinal process; d, dorsal cavity (head-part). The two endothelial cardiac tubes are still separated by the octogenetic septum (s) of the Amnion, g, vessels.

furrow is soon seen in the ventricle (Fig. 376 c). This is the external sign of the internal partition by which the ventricle is divided into two—a right venous and left arterial ventricle. Finally a longitudinal partition is formed in the third section of the primitive fish-like heart, the arterial bulb, externally indicated by a longitudinal furrow (Fig. 376 d). The cavity of the bulb is divided into two lateral halves, the pulmonary-artery bulb, that opens into the right ventricle, and the aorta-bulb, that opens into the left ventricle. When all the partitions are complete, the small (pulmonary) circulation is distinguished from the large (body) circulation; the motive centre of the former is the right half, and that of the latter the left half, of the heart.

placement is greatest in the anthropoid apes—chimpanzee, gorilla, and orang—which resemble man in this.

As the heart of all Vertebrates is originally, in the light of phylogeny, only a local enlargement of the middle principal vein, it is in perfect accord with the biogenetic law that its first structure in the embryo is a simple spindle-shaped tube in the ventral wall of the head-gut. A thin membrane, standing vertically in the middle plane,

of the head-gut with the lower head-wall. As the tube extends and detaches from the gut-wall, it divides the mesocardium into an upper (dorsal) and lower (ventral) plate (usually called the *mesocardium anterior* and *posterior* in man, Fig. 379 abg). The

mesocardium divides two lateral cavities, Remak's "neck-cavities" (Fig. 379 A). These cavities afterwards join and form the simple pericardial cavity, and are therefore called by Kolliker the "primitive pericardial cavities."

The double cervical cavity of the Amniotes is very interesting, both from the anatomical and the evolutionary point of view; it corresponds to a part of the hypocoelomites of the head of the lower Vertebrates—that part of the ventral coelom-pouch...

Wilhe's "visceral cavities" below. Each of the cavities still communicates freely behind with the two coelom-pouches of




FIG. 38.—Frontal section of a human embryo, one-twelfth of an inch long in the neck, magnified forty times. "separated by Wilhelm's. Seen from ventral side, and mesothoracic, surrounded by the bronchial processes, ad, lobes of lungs, ao, middle part of ventricle, ad, left lateral part of meso, ae, aorta, d, diaphragm, vc, superior vena cava, va, umbilical vein, cp, vitelline space, lg, liver, lg, hepatic duct.

the trunk; and, just as these afterwards coalesce into a simple body-cavity (the ventral mesentery disappearing), we find the same thing happening in the head. This simple primary pericardial cavity has been well called by Gegenbaur the "head-coelom," and by Hertwig the "pericardial breast-cavity." As it now encloses the heart, it may also be called *cardio-coel*.

The cardio-coel, or head-coelom, is often disproportionately large in the Amniotes, the simple cardiac tube growing considerably and lying in several folds. This causes the ventral wall of the amniote embryo between the head and the meso-

to be pushed outwards as in rupture (cf. Fig. 380 A). A transverse fold of the ventral wall, which receives all the vein-trunks that open into the heart, grows up from below between the pericardium and the stomach, and forms a transverse partition, which is the first structure of the primary diaphragm (Fig. 380 d). This important muscular partition, which completely separates the thoracic and abdominal cavities in the mammals alone, is still very imperfect here; the two cavities still

als. These canals, which belong to dorsal part of the head-coelom, and which we may call briefly *pleural ducts*, receive the two pulmonary sacs, which develop from the hind end of the ventral wall of the head-gut; they thus become the two pleural cavities.

The diaphragm makes its first appearance in the class of the Amphibia (in the salamanders) as an insignificant muscular transverse fold of the ventral wall, which arises from the fore end of the transverse abdominal muscle, and grows between the pericardium and the liver. In the reptiles (turtles and crocodiles) a later dorsal part is joined to this earlier ventral part of the rudimentary diaphragm, a pair of subvertebral muscles rising from the vertebral column and being added as "columns" to the transverse partition. But it was probably in the Permian sauro-mammals that the two originally separate parts were united, and the diaphragm became a complete partition between the thoracic and abdominal cavities in the mammals; as it considerably enlarges the chest-cavity when it contracts, it becomes an important respiratory muscle. The ontogeny of the diaphragm in man and the other mammals reproduces this phylogenetic process to-day, in accordance with the biogenetic law, in all the mammals the diaphragm is formed by the secondary conjunction of the two originally separate structures, the earlier ventral part and the later dorsal part.

Sometimes the blending of the two diaphragmatic structures, and consequently the severance of the one pleural duct from the abdominal cavity, is not completed in man. This leads to a diaphragmatic rupture (*hernia diaphragmatica*). The two cavities then remain in communication by an open pleural duct, and loops of the intestine may penetrate by this "rupture opening" into the chest-cavity. This is one of three

fatal mis-growths that show the great part that blind chance has in organic development.

Thus the thoracic cavity of the mammals, with its important contents, the heart and

specia. In its earliest form the heart is double, as is

shown, in all the Anniotes, and the simple spindle-shaped cardiac tube, which

took its name from a point, is only formed at a later stage, when the two lateral tubes move backwards, touch each other, and at last combine in the middle line. In man, as in the rabbit, the two embryonic hearts are still far apart at the stage when there are already eight primitive segments (Fig. 134 A). So also the two out-pouches of the head in which they lie




FIG. 131.—Transverse section of the head of a rabbit-embryo, twenty-four hours old. Underneath the medullary tube the two primitive

lungs, belongs originally to the head-part of the vertebrate body, and its inclusion in the trunk is secondary. This instructive and very interesting fact is entirely proved by the concordant evidence of comparative anatomy and ontogeny. The lungs are outgrowths of the head-gut; the heart develops from its inner wall. The pleural sacs that enclose the lungs are dorsal parts of the head-coelom, originating from the pleuroducts; the pericardium in which the heart afterwards lies is also double originally, being formed from ventral halves of the head-coelom, which only combine at a later stage.

When the lung of the air-breathing Vertebrates issues from the head-cavity and enters the trunk-cavity, it follows the example of the floating bladder of the fishes, which also originates from the pharyngeal wall in the shape of a small pouch-like out-growth, but soon grows so large that, in order to find room, it has to pass far behind into the trunk-cavity. To put it more precisely, the lung of the quadrupeds retains this hereditary growth-process of the fishes; for the hydrostatic floating bladder of the latter is the air-filled organ from which the air-breathing organ of the

There is an interesting ontogenetic phenomenon in the formation of the heart

bryo develops and detaches from the embryonic vesicle that the separate lateral structures join together, and finally combine in the middle line. As the median partition between the right and left cardiac cavities disappears, the two cervical cavities freely communicate (Fig. 381), and form, on the ventral side of the annulate head, a horseshoe-shaped arch, the points of which advance backwards into the pleuroducts or pleural cavities, and from there into the two peritoneal sacs of the trunk. But even after the conjunction of the cervical cavities (Fig. 381) the two cardiac tubes remain separate



FIG. 132.—Transverse section of the anterior region of the mouse embryo (before the pharynx). In the cervical cavity (a), the heart (b) is still connected by a constriction (c) with the gut-tube layer (d). e, primitive layer, or pericardial space. f, embryonic cavity in the horn of the annulate field. (From Arnold.)

at first; and even after they have united a delicate partition in the middle of the simple endothelial tube (Figs. 379 & 381 A) indicates the original separation. This ontogenetic "primary cardiac sep-

turn" presently disappears, and has no relation to the subsequent permanent partition between the halves of the heart, which, as a heritage from the reptiles, has a great phylogenetic importance.

Thorough opponents of the biogenetic law have laid great stress on these and similar ontogenetic phenomena, and endeavoured to urge them as striking disproofs of the law. As in every other instance, careful, discriminating, comparative-morphological examination converts these supposed disproofs of evolution into strong arguments in its favour. In his excellent work, *On the Structure of the Heart in the Amphibia* (1886), Carl Rabl has shown how easily these curious ontogenetic facts can be explained by the secondary adaptation of the embryonic

structure to the great extension of the food-yolk.

The embryology of all the other parts of the vascular system also gives us abundant and valuable data for the purposes of phylogeny. But as one needs a thorough knowledge of the intricate structure of the whole vascular system in man and the other Vertebrates in order to follow this with profit, we cannot go into it further here. Moreover, many important features in the ontogeny of the vascular system are still very obscure and controverted. The characters of the embryonic circulation of the Amniotes, which we have previously considered (Chapter XV.), are late acquisitions and entirely ontogenetic. (Cf. pp. 170-171, Figs. 198-202.)

CHAPTER XXIX.

EVOLUTION OF THE SEXUAL ORGANS

If we measure the importance of the systems of organs in the animal frame according to the richness and variety of their phenomena and the physiological interest that this implies, we must regard as one of the principal and most interesting systems the one which we are now going to examine—the system of the reproductive organs. Just as nutrition is the first and most urgent condition for the self-maintenance of the individual organism, so reproduction alone secures the maintenance of the species—or, rather, the maintenance of the long series of generations which the totality of the organic stem represents in their genealogical connection. No individual organism has the prerogative of immortality. To each is allotted only a brief span of

million-year course of the

of life.

Hence, reproduction and the correlative phenomenon, heredity, have long been regarded, together with nutrition, as the most important and fundamental functions of living things, and it has been attempted to distinguish them from "lifeless bodies" on this very score. As a

matter of fact, this division is not so profound and thorough as it seems to be, and is generally supposed to be. If we examine carefully the nature of the reproductive process, we soon see that it can be reduced to a general property that is found in inorganic as well as organic bodies—growth. Reproduction is a nutrition and growth of the organism beyond the individual limit, which raises a part of it into the whole. This is most clearly seen when we study it in the simplest and lowest organisms, especially the Monera (Figs. 226-228) and the unicellular Amœbe (Fig. 27). There the simple individual is a single plantlet. As soon as it has reached a certain limit of size by continuous feeding and normal growth, it cannot pass it, but divides, by simple cleavage, into two equal halves.

Each of these half

independent life, and grow

until it reaches the limit of growth, and divides. In each of these acts of self-cleavage two new centres of attraction are formed for the particles of bodies, the foundations of the two new-formed individuals. There is no such thing as immortality even in these unicellulars.

The individual as such is annihilated in the act of cleavage (cf. p. 48).

In many other Protzoa reproduction takes place not by cleavage, but by budding (gemmation). In this case the growth that determines reproduction is not total (as in segmentation), but partial. Hence in gemmation also we may oppose the head growth-product, that becomes a new individual in the bud, as a child-organism to the parent-organism from which it is formed. The latter is older and larger than the former. In cleavage the two products are equal in age and morphological value. Next to gemmation we have, as other forms of asexual reproduction, the forming of embryonic buds and the forming of embryonic cells. But the latter leads us at once to sexual generation, the distinctive feature of which is the separation of the sexes. I have dealt fully with these various types of reproduction in my *History of Creation* (chap. viii.) and my *Wonders of Life* (chap. xi.).

The earliest ancestors of man and the higher animals had no faculty of sexual reproduction, but multiplied solely by asexual means—cleavage, gemmation, or the formation of embryonic buds or cells, as many Protzoa still do. The differentiation of the sexes came at a later stage. We see this most plainly in the Protists, in which the union of two individuals precedes the continuous cleavage of the unicellular organism (transitory conjugation and permanent copulation of the Infusoria). We may say that in this case the growth (the condition of reproduction) is attained by the coalescence of two full-grown cells into a single, disproportionately large individual. At the same time, the mixture of the two plastids causes a rejuvenation of the plasma. At first the copulating cells are quite homogeneous, but natural selection soon brings about a certain contrast between them—larger female cells (*macrospores*) and smaller male cells (*microspores*). It must be a great advantage in the struggle for life for the new individual to have inherited different qualities from the two cellular parents. The further advance of this contrast between the generating cells led to sexual differentiation. One cell became the female ovum (*macrogamete*), and the other the male sperm-cell (*microgamete*).

The simplest forms of sexual reproduction among the living Metazoa are seen in the Coelenterata (p. 232), the lower

sponges, the common fresh-water polyp (*Hydra*), and other Coelenterata of the lowest rank. *Prophyrenia* (Fig. 234), *Olynthus* (Fig. 238), *Hydra*, etc., have very simple tubular bodies, the thin wall of which consists (as in the original gastrula) only of the two primary germinal layers. As soon as the body reaches sexual maturity, a number of the cells in its wall become female ova, and others male sperm-cells: the former become very large, as they accumulate a considerable quantity of yolk-granules in their protoplasm (Fig. 235 c); the latter are very small on account of their repeated cleavage, and change into mobile cone-shaped spermatozoa (Fig. 20). Both kinds of cells detach from their source of origin, the primary germinal layers, fall either into the surrounding water or into the cavity of the gut, and unite there by fusing together. This is the momentous process of fecundation, which we have examined in the seventh Chapter (cf. Figs. 23-29).

From these simplest forms of sexual propagation, as we can observe them to-day in the lowest Zoophytes, the Coelenterata, Sponges, and Polyps, we gather most important data. In the first place, we learn that, properly speaking, nothing is required for sexual reproduction except the fusion or coalescence of two different cells—a female ovum and male sperm-cell. All other features, and all the very complex phenomena that accompany the sexual act in the higher animals, are of a subordinate and secondary character, and are later additions to this simple, primary process of copulation and fecundation. But if we bear in mind how extremely important a part this relation of the two sexes plays in the whole of organic nature, in the life of plants, of animals, and of man, how the mutual attraction of the sexes, love, is the mainspring of the most remarkable processes—in fact, one of the chief mechanical causes of the highest development of life—we cannot too greatly emphasise this tracing of love to its source, the attractive force of two erotic cells.

Throughout the whole of living nature the greatest effects proceed from this very small cause. Consider the part that the flowers, the sexual organs of the flowering plants, play in nature; or the exuberance of wonderful phenomena that sexual selection produces in animal life; or the

mother . . . influence of love in the life of man. In every case the fusion of two cells is the sole original motive power; in every case this invisible process profoundly affects the development of the most varied structures. We may say, indeed, that no other organic . . . be compared to it for a more prehensiveness and intensity of . . .

. . . not the Semitic myth of Adam and Eve, the old Greek legend of Paris and Helena, and so many other famous traditions, only the poetic expression of the vast influence that love selection have exercised over of history ever since the differ

other pass

agitate the heart

tripped in their joint influence by this senso-inflaming and mind-benumbing Eros. On the one hand, we look to love with gratitude as the source of the greatest artistic achievements—the noblest creations of poetry, plastic art, and music; we see in it the chief factor in the moral advances of humanity, the foundation of family life, and therefore of social advance. On the other hand, we dread it as the devouring flame that brings destruction on so many, and has caused more misery, vice, and crime than all the other evils of human life put together. So wonderful is love and so momentous its influence on the life of the soul, or on the different functions of the medullary tube, that here more than anywhere else the "supernatural" result seems to mock any attempt at natural explanation. Yet comparative evolution leads us clearly and indubitably to the first source of love affinity of two different erotic cells, the sperm-cell and ovum.

The lowest Metazoa throw light on this very simple origin of the intricate phenomena of reproduction, and they also teach us that the earliest sexual form was hermaphroditism, and that the separation of the sexes (by division of labour) is a secondary and later phenomenon. . . . aphroditism predominant . . .

most varied groups of the low each sexually-mature individual, each person, contains female and male sexual cells, and is therefore able to fertilize itself and reproduce. Thus we find our

and sperm-cells in the same individual, not only in the lowest Zoophytes (Gastreaads, Sponges, and many Polyps), but also in many worms (leeches and earth-worms), many of the sn garden and vineyard unicates, and animals. All r

the Gastreaads up the Prochordonia, were hermaphroditic—possibly even the earliest Acrania. have an instructive proof of this i remarkable . . .

genera of fishes are still hermaphrodites, and that it is occasionally found in the higher Vertebrates of all classes (as stavians). We may conclude from this that gonochorism (separation of the sexes) was a later stage in our development. At first, male and female individuals differ only in the possession of one or other kind of gonads; in other respects they were identical, as we still find in the Amphioxus and the Cyclostomes. Afterwards, accessory organs (ducts, etc.) are associated with the primary sexual glands; and much later again sexual selection has given rise to the secondary sexual characters—those differences between the sexes which do not affect the sexual organs themselves, but other parts of the body (such as the man's beard or the woman's breast).

THE third important fact that we learn from the lower Zoophytes relates to the earliest origin of the two kinds of sexual cells. As in the Gastreaads (the lowest sponges and hydroids), in which we find the first beginnings of sexual differentiation, the whole body consists merely of

cooded from the cells of these primary layers, either the inner or outer, or from both. This simple fact is extremely important, because the first trace of the ova as well as the spermatozoa is found in the middle germinal layer or mesoderm in the higher animals, especially the Vertebrates.

development from . . . section with the secondary formation of the (mesoderm).

If we trace the phylogeny of the sexual organs in our earliest Metazoa ancestors,

of the lowest Coelentera (*Chidaria*, *Platodaria*) exhibit it to us, we find that the first step in advance is the localisation or concentration of the two kinds of sexual

altered in the epithelium
the Sponges and

for

detached &

vary germinal layers, and become free
the Cnidaria and
Platodes we find these associated in
groups which we call sexual glands
(gonads). We can now for the first
time speak of sexual organs in the mor-
phological sense. The female germina-
tive glands, which in this simplest form
are merely groups of homogeneous cells,
are the ovaries (Fig. 241 c). The male
germinative glands, which also in their
first form consist of a cluster of sperm-
cells, are the testicles (Fig. 241 A). In
the medusae, which descend, both ontog-
enetically and phylogenetically, from the
more simply organized Polype, we find
these simple sexual glands sometimes as

that appear at the edge of the primitive
mouth (right and left), as a rule during
gastrulation or immediately afterwards—
the important promesoblasts, or "polar
cells of the mesoderm," or "primitive
cells of the middle germinal layer" (p.
194). In the real Enterocœla, in which
the mesoderm appears from the first in
the shape of a couple of coelom-pouches,
these are very probably the original
gonads (p. 194). This is seen very clearly
in the arrow-worm (*Sagitta*) in the
gastrula of *Sagitta* (Fig. 383 A) we find
at an early stage a couple of entodermic
cells of an unusual size (*g*) at the base
of the primitive gut (*st*). These primi-
tive sexual cells (*progonadas*) are sym-
metrically placed to the right and left of
the middle plane, like the two promeso-
blasts of the bilateral gastrula of the
Amphioxus (Fig. 38 p. p. 66). A little



383.—*Xenopus* of *Sagitta*, in three earlier stages of development. (From Herberich) A gastrula, *st* with open primitive mouth. B the same with primitive mouth closed, no permanent gut. C primitive gonads (entodermic primitive mesodermic cells), no coelom-pouches, *p* parietal layer, *m* mesodermic layer of meso, *st* permanent gut (stomach), *st* mouth-pit (stomodæum).

gastric pouches, sometimes as outgrowths
of the radial canals that projected from the
stomach. Particularly interesting in con-
nection with the question of the first
origin of the gonads are the lowest forms
of the Platodes, the *Cryptocœla* that have
of late been separated as a special class
(*Platodæa*) from the Turbellaria proper
(Fig. 239). In these very primitive
Platodes the two pairs of sexual glands
are merely two pairs of rows of differenti-
ated cells in the entodermic wall of the
primitive gut—two median ovaries (*o*)
within, and two lateral spermaries (*s*)
without. The mature sexual cells are
ejected by the posterior outlets; the female
(*f*) lies in front of the male (*m*).

In the great majority of the Bilateria
or Coelomaria it is the mesoderm from
which the gonads develop. Probably the
first traces of them are the two large cells

outwards from them the two coelom
pouches (*B, c*) are developed out of the
primitive gut, and each progenitor
divides into a male and a female sexual
cell (*B, g*). The two male cells (at first
rather the larger) lie close together within,
and are the parent-cells of the testicles
(*protermanes*). The two female cells
lie outwards from these, and are the
parent-cells of the ovary (*protermanes*).
Afterwards, when the coelom-pouches
have detached from the permanent gut
(*C, d*) and the primitive mouth (*A, st*)
is closed, the female cells advance towards
the mouth (*C, st*), and the male towards
the rear. The foremost pair of ovaries
are then separated by a transverse parti-
tion from the hind pair. Thus the first
structures of the sexual glands of the
Sagitta are a couple of hermaphroditic
entodermic cells; each of these divides

into a male and a female cell; and these four cells are the parent-cells of the four sexual glands. Probably the two pro-mesoblasts of the *Amphioxus-gastrula* (Fig. 98) are also hermaphroditic primitive sexual cells in the same sense, inherited by this earliest vertebrate from its ancient bilateral gastræan ancestors.



FIG. 94.—A Part of the Kidneys of *Mallophaga*. A proximal duct (nephridium). B distal part of primitive urinary ducts (nephridium), renal or Malpighian capsule. C portion of duct, highly convoluted. D renal papilla with the glomerulus of afferent artery. E efferent artery. From Johannes Müller (*Mallophaga*).

The sexually-mature *Amphioxus* is not hermaphroditic, as its nearest invertebrate relatives, the Tunicates, are, and as the long-extinct pre-Silurian Primitive Vertebrate (*Protophyllus*, Figs. 98-102) probably was. The actual lancelet has gonochoristic structures of a very interesting kind. As we saw in the anatomy of the *Amphioxus*, we find the ovaries of the female and the spermaries of the male in the shape of twenty to thirty pairs of elliptical or roundish four-cornered sacs,

which lie on either side of the gut on the parietal surface of the respiratory pore (Fig. 119 g). According to the important discovery of Ruckert (1888), the sexual glands of the earliest fishes, the *Selachii*, are similarly arranged. They only unite afterwards to form a pair of simple gonads. These have been transmitted by heredity to all the rest of the Craniotes. In every case they lie originally on each side of the mesentery, underneath the chorda, at the bottom of the body-cavity. The first traces of them are found in the coelom-epithelium, at the spot where the skin-fibre layer and gut-fibre layer meet in the middle of the mesenteric plate (Fig. 93 mp). At this point we observe at an early stage in all craniote embryos a small string-like cluster of cells, which we may call, with Waldeyer, the "germ epithelium," or (in harmony with the other plate-shaped rudimentary organs) the *sexual plate* (Fig. 173 g). This germinal or sexual plate is found in the fifth week in the human embryo, in the shape of a couple of long whitish streaks, on the inner side of the primitive kidneys (Figs. 183 f). The cells of this sexual plate are distinguished by their cylindrical form and chemical composition from the rest of the coelom-cells; they have a different purpose from the flat cells which line the rest of the body-cavity. As the germ epithelium of the sexual plate becomes thicker, and supporting tissue grows into it from the mesoderm, it becomes a rudimentary sexual gland. This ventral gonad then develops into the ovary in the female Craniotes, and the testicles in the male.

In the formation of the gonidia or erotic sexual cells and their conjunction at fecundation we have the sole essential features of sexual reproduction; but in the great majority of animals we find other organs taking part in it. The chief of these secondary sexual organs are the gonoducts, which serve to convey the mature sexual cells out of the body, and the copulative organs, which bring the fecundating male sperm into touch with the ovum-bearing female. The latter organs are, as a rule, only found in the higher animals, and are much less widely distributed than the gonoducts. But these also are secondary formations, and are wanting in many animals of the lower groups.

In the lower animals the mature sexual cells are generally ejected directly from

the body. Sometimes they pass out immediately through the skin (Hydra and many hydroids); sometimes they fall into the gastric cavity, and are evacuated by the mouth (gastropods, sponges, many medusae, and corals); sometimes they pass into the body-cavity, and are ejected by a special pore (*porus genitalis*) in the ventral wall. The latter procedure is found in many of the worms, and also in the lowest Vertebrates. Amphioxus has the peculiar feature that the mature sexual products fall first into the mantle-cavity; from there they are either evacuated by the respiratory pore, or else they pass through the gill-slits into the branchial gut, and so out by the mouth (p. 183). In the Cyclostomes they fall into the body-cavity, and are ejected by a genital pore in its wall; so also in some of the fishes. From these we gather the fea-

to convey the sexual products, and this had originally a totally different function—namely, the system of urinary organs. These organs have primarily the sole duty of removing unusable matter from the body in a fluid form. Their liquid excretory product, the urine, is eliminated directly through the skin or through the last section of the gut. It is only at a later stage that the tubular urinary passages also convey the sexual products from the body. In this way they become "inducts." This remarkable secondary conjunction of the urinary and sexual organs into a common urogenital system is very characteristic of the Gnathostomes, the six higher Vertebrates. It is wanting in the lower classes. In order to appreciate it fully, we must give a comparative glance at the structure of the u



1. Transverse section of the embryonic shield of a shark, forty-two hours old. (From "The Medullary Tube, or Notochord, a Horny Plate (skin-venter layer), skin apophyses, the stomodaeum (stomodaeum layer), the stomodaeum layer (stomodaeum layer of the hypodermis), the gonoduct layer (ventral layer of the hypodermis).")

tures of our earlier ancestors in this respect. On the other hand, in all the higher and most of the lower Vertebrates (and most of the higher Invertebrates) we find in both sexes special tubular passages of the sexual gland, which are called "gonoducts." In the female they conduct the ova from the ovary, and so are called "oviducts," or "Fallopian tubes." In the male they convey the spermatozoa from the testicles, and are called "spermatoducts," or *vas deferens*.

The original and genetic relation of these two kinds of ducts is just the same in man as in the rest of the higher Vertebrates, and quite different from what we find in most of the Invertebrates. In the latter, as a rule, the gonoducts develop directly from the embryonic glands or from the outer skin; but in the Vertebrates an independent organic system is employed

The renal or urinary system is one of the oldest and most important systems of organs in the differentiated animal body, as I have pointed out on several previous occasions (cf. Chapter XVII.). We find it not only in the higher stems, but also very generally distributed in the earlier group of the Vermalia. Here we meet it in the lowest worms, the Rotatoria (Gastrotrocha, Fig. 242), and in the instructive stem of the Platyhelminthes. It consists of a pair of simple or branching canals, which are lined with one layer of cells, absorb unusable juices from the tissue, and eject them by an outlet in the outer skin (Fig. 240 am). Not only the free-living Turbellaria, but also the parasitic Suctorid, and even the still more degenerate tapeworms, which have lost their alimentary canal in consequence of their parasitic life, are equipped with these renal canals

or nephridia. In the first embryonic structure they are merely a pair of simple cutaneous glands, or depressions in the ectoderm. They are generally described as excretory organs in the worms, but



FIG. 34.—Bodimentary primitive kidneys of a dog-embryo. The head end of the embryonic body is seen from the ventral side and marked with the visceral layer of the yolk-sac, which is torn away and folded down in front in order to show the nephridium with the primitive urinary canal (a). A primitive nephridium, a special canal of entrance into the pronephros — (b). (From Dr. B. S. P. S. S. S.)

formerly often as "water vessels." They may be conceived as largely-developed tubular cutaneous glands, formed by invagination of the cutaneous layer. According to another view, they

nephridium has an inner opening (with cilia) into the body-cavity and an outer one on the epidermis.

In these lowest, unsegmented worms, and in the unsegmented Molluscs, there is only one pair of renal canals. They are more numerous in the higher Articulata. In the Annelids, the body of which is composed of a large number of joints, there is a pair of these pronephridia in each segment (hence they are called segmental canals or organs). Even here they are still simple tubes; on account of their coiled or looped form they are often called

"looped canals." In most of the Annelids, and many of the Vermalis, we can distinguish three sections in the nephridium—an outer muscular duct, a glandular middle part, and an inner part that opens by a ciliated funnel into the body-cavity. This opening is furnished with whirling cilia, and can, therefore, take up the juices to be excreted directly from the body-cavity and convey them from the body. But in these worms the sexual cells, which develop in very primitive form on the inner surface of the body-cavity, also fall into it when mature, and are sucked up by the funnel-shaped inner ciliated openings of the renal canals, and hence into the urine. Thus the urine

materially different from these segmental canals of the Annelids. The peculiar development of it and its relations to the sexual organs are among the most difficult problems in the morphology of our stem. If we examine briefly the vertebrate renal system from the phylogenetic point of view, it is confirmed by cent

we may
n three
it: (1)
says or
eye (pro-
(2) primi-
dle kid-
neurons-

kidneys / met-
These
of

kidneys are not fundamentally and completely distinct, as earlier students (such as Semper) wrongly supposed; they represent three different generations of one and the same excretory apparatus; they correspond to three phylogenetic stages,




FIG. 35.—Primitive kidneys of a human embryo. a, the urinary canal of the primitive kidneys, as Wolffian duct of hypoblast and of the same (Müllerian's hyalid), as Müllerian duct of epiblast and of (Falkenberg's hyalid) and (sexual glands) (F. S. S. S.)

and succeed each other in the stern-history of the Vertebrates in such wise that the more advanced generation develops further behind in the body, and replaces the older and less



FIG. 95.—Pig-embryo, three-fifths of an inch long magnified ten times, seen from the ventral side. *a* fore leg; *b* head leg; *c* ventral wall; *d* notochord; *e* pronephros or primitive kidney; *f* primitive kidneys, as their inner part (pronephros).

advanced generation that preceded it in time and space. The *fore kidneys*, first accurately described by Wilhelm Müller in 1875 in the Cyclostomes and Ichthyofauna, form the sole excretory organ of the Acrania (Amphioxus); they continue in the Cyclostomes and some of the fishes, but are found only in slight traces and for a time in the embryos of the six other classes of Vertebrates. The *primitive kidneys* are first found in the Cyclostomes, behind the fore kidneys; they have been transmitted from the Selachii to all the Gnathostomes. In the *Amphioxus* they act permanently as urinary glands; in the *Amphioxus* their anterior part ("germinal kidneys") changes into organs of the sexual apparatus, while the third generation develops from the end of their posterior part ("urinal kidneys")—the characteristic after or permanent kidneys of the three higher classes of Vertebrates. The order in which the three renal systems succeed each other in the embryo of man and the higher Vertebrates corresponds to their phylogenetic

the history of our stem, and, consequently, in the natural classification of the Vertebrates.

As in the morphology of any other system of organs, so in the case of the urinary and sexual organs the *Amphioxus* is the real typical primitive Vertebrate; it affords the key to the mysteries of the structure of man and the higher Vertebrates. The kidneys of the *Amphioxus*—first discovered by Boveri in 1890—are typical "fore kidneys," composed of a double row of short segmental canals (Fig. 217 *x*). The inner aperture of these




FIG. 96.—Human embryo of the fifth week, two-fifths of an inch long, magnified ten times, seen from the ventral side (the anterior ventral wall, *a*, is removed, the body-cavity, *c*, exposed). *d* gut (not off); *e* forebrain; *f* heart; *g* liver; *h* fore-gut; *i* pulmonary sac; *j* primitive kidneys; *k* neural region; *l* pharynx (notal protuberance); *m* tail. (From Kallmann.)

pronephridia opens into the mesodermic body-cavity (the middle part of the coeloma, *B*); the external aperture into the ectodermic mantle or peribranchial cavity (*C*). Their position, their

structure, and their relation to the branchial vessel make it clear that these segmental protonephridia correspond to the rudimentary fore kidneys of the Crustacea. The mantle-cavity into which



FIG. 329.

FIG. 330.

FIG. 331.

secondary renal

Fig. 329 of a nemertean (larva) shows a primitive kidney, a ventral gland (rudiment of testis and ovary). The primary nephroduct (a) in Fig. 330 divides (cf. Figs. 329 and 330) into the two secondary nephroducts—the anterior (a1) and posterior (a2) ducts, joined together behind in the genital coel (c), forming the posterior kidneys. (From Gegenbaur.)

they open seems to correspond to the prorenal duct of the latter.

The next higher Vertebrates, the Cyclostomes, yield some very interesting data. Both orders of this class, the hags and lampreys, have still the fore kidneys inherited from the Acrania—the former permanently, the latter in their earlier stages. Behind these the primitive kidneys soon develop, and in a very characteristic form. The remarkable structure of the mesonephros of the Cyclostomes, discovered by Johannes Müller, explains the intricate formation of the kidneys in the higher Vertebrates. We find in the hag-fishes (*Bdellostoma*) a long tube, the prorenal duct (*nephroductus*, Fig. 384 a). This opens with its anterior end into the coeloma by a ciliated aperture, and externally with its posterior end by an outlet in the skin. Inside it opens a large number of small transverse canals ("segmental or primitive urinary canals," b). Each of these terminates blindly in a vesicular capsule (c), and this encloses a coil of blood-vessel (*glomerulus*, an arterial network, Fig. 384 B, c). Afferent branches of arterioles conduct arterial blood into the coiled branches of the glomerulus (d), and efferent arterial branches con-

duct it away from the set (e). The primitive renal canals (*metanephridia*) are distinguished by their out-formation from their predecessors.

In the Selachii also we find a longitudinal row of segmental canals on each side, which open outwards into the primitive renal ducts (*metanephros*, p. 149). The segmental canals (a pair in each segment of the middle part of the body) open internally by a ciliated funnel into the body-cavity. From the posterior



FIG. 383.

FIG. 384.

FIG. 383, 384.—Urinary and sexual organs of a lamprey (water excretory or Triton). Fig. 383 of a female, 384 of a male. a primitive kidney, ovary, uterine duct, and testis duct, both

water-take along a
a below testis, b below
in Digestive)

group of these organs a compact primitive kidney is formed, the anterior group taking part in the construction of the sexual organ.

In the same simple form that remains

throughout life in the Myxaloides and partly in the Selachii we find the primitive kidney first developing in the embryo of man and the higher Craniotes (Figs. 386, 387). Of the two parts that compose the comb-shaped primitive kidney the longitudinal channel, or nephroduct, is always the first to appear; afterwards the transverse "canals," the excreting nephridia, are formed in the mesoderm; and after this again the Malpighian capsules with their arterial coils are associated with these as curious outgrowths. The primitive renal duct, which appears first, is found in all craniote embryos at the early stage in which the differentiation of the medullary tube takes place in the ectoderm, the severance of the chorda from the visceral layer in the endoderm, and the first trace of the coelom-pouches arises between the limiting layers (Fig. 385). The nephroduct (*ang*) is seen on each side, directly under the horny plate, in the shape of a long, thin, thread-like string of cells. It presently hollows out and becomes a canal, running straight from front to back, and clearly showing in the transverse section of the embryo its original position... the space between horny plate (*A*), primitive segments (*an*), and lateral plates (*apl*). As the originally very short urinary canals lengthen and multiply, each of the two primitive kidneys assumes the form of a half-feathered leaf (Fig. 387). The lines of the leaf are represented by the urinary canals (*u*), and the rib by the outlying nephroduct (*x*). At the inner edge of the primitive kidneys the rudiment of the ventral sexual gland (*g*) can now be seen as a body of some size. The hindmost end of the nephroduct opens right behind into the last section of the rectum, thus making a cloaca of it. However, this opening of the nephroducts into the intestine must be regarded as a secondary formation. Originally they open, as the Cyclostomes clearly show, quite independently of the gut, in the external skin of the abdomen.

In the Myxaloides the primitive kidneys remain this simple comb-shaped structure,

and a part of it is preserved in the Selachii; but in all the other Craniotes it is only found for a short time in the embryo, as an ontogenetic reproduction of the earlier phylogenetic structure. In these the primitive kidney soon assumes the form (by the rapid growth, lengthening, increase, and serpentine of the urinary canals) of a large compact gland, of a long, oval or spindle-shaped character, which passes through the greater part of the embryonic body-cavity (Figs. 183 *w*, 184 *w*, 388 *w*). It lies near the middle line, directly under the primitive vertebral column, and reaches from the cardiac



human embryo, three inches in length (measuring of the snout with), magnified fifteen times. *A*, horny plate; *an*, primitive segments; *apl*, lateral plates; *u*, urinary canals; *x*, outlying nephroduct; *g*, ventral sexual gland; *ang*, nephroduct. (From Kollmann.)

region to the cloaca. The right and left kidneys are parallel to each other, quite close together, and only separated by the mesentery—the thin narrow layer that attaches the middle gut to the under surface of the vertebral column. The passage of each primitive kidney, the nephroduct, runs towards the back on the lower and outer side of the gland, and opens in the cloaca, close to the starting-point of the allantois; it afterwards opens into the allantois itself.

The primitive or primordial kidneys of the equiote embryos were formerly called the "Wolffian bodies," and sometimes "Olsen's bodies." They act for a time as

kidneys, absorbing unusable juices from the embryonic body and conducting them to the cloaca—afterwards to the allantois. There the primitive urine accumulates, and thus the allantois acts as bladder or urinary sac in the embryos of man and the other Amniotes. It has, however, no genetic connection with the primitive kidneys, but is a pouch-like growth from the anterior wall of the rectum (Fig. 147 N). Thus it is a product of the visceral layer, whereas the primitive kidneys are a product of the middle layer. Phylo-

ventral wall. It is true that many of the fishes also have a "bladder," but this is merely a local enlargement of the lower section of the nephroducts, and so totally different in origin and composition from the real bladder. The two structures can be compared from the physiological point of view, and so are *analogous*, as they have the same function; but not from the morphological point of view, and are therefore not *homologous*. The false bladder of the fishes is a mesodermic product of the nephroducts; the true

bladder of the Dipneusta, Amphibia, and Amniotes is an endodermic blind sac of the rectum.

In all the Acamnia (the lower ammonious iotes, Cyclostomes, Fishes, Dipneusta and Amphibia) the urinary organs remain at a lower stage of development to this extent, that the primitive kidneys (*protonephros*) last per-

.....
sist. This is only so as a passing phase of the early embryonic life in the three higher classes of Vertebrates, the Amniotes. In the permanent or after or (really *ter* neys (re- neys) th tion of

their appearance. They represent the gear

..... of the
brans kidneys. The permanent kidneys do not arise (as was long independent glands from the alimentary tube, but from the last section of the primitive kidneys and the nephroduct. Here a simple tube, the secondary renal duct, develops, near the point of its entry into the cloaca; and this tube grows considerably forward. With its blind upper or anterior end is connected a glandular renal growth, that owes its origin to a differentiation of the last part of the primitive kidneys. This rudiment of the



FIG. 397.

FIG. 398.

FIG. 399.

FIG. 397-399.—Urinary and sexual organs of an embryo. FIG. 397, female embryo one and a half inches long. FIG. 398, male embryo, one and a half inches long. FIG. 399, female embryo two and a half inches long. a, primitive kidney, see Wolffian duct, in Müllerian duct, at upper end of same (top of a), a lower and b, other part of same (rudiment of uterus), c, genital cord, d, testis (A lower and B upper testicular ligament), e, ovary, f, lower ovarian ligament, g, inguinal ligament of primitive kidney, h, dophergastic ligament of primitive kidney, an accessory kidney, n permanent kidney, under them the h-shaped the rectum, v bladder, a umbilical artery. (From Kähler.)

genetically we must suppose that the allantois originated as a pouch-like growth from the cloaca-wall in consequence of the expansion caused by the urine accumulated in it and secreted by the kidneys. It is originally a blind sac of the rectum. The real bladder of the vertebrate certainly made its first appearance among the Dipneusta (in Lepidodactylus), and has been transmitted from hibia, and from these to the embryo of the latter it protrudes far out of the not yet closed

permanent kidneys consists of coiled urinary canals with Malpighian capsules and vascular coils (without ciliated funnels), of the same structure as the segmental mesonephridia of the primitive kidneys. The further growth of these metanephridia gives rise to the compact permanent kidneys, which have the familiar bean-shape in man and most of the higher mammals, but consist of a number of separate folds in the lower mammals, birds, and reptiles. As the permanent kidneys grow rapidly and advance forward, their passage, the ureter, detaches altogether from its birth-place, the posterior end of the nephroduct; it passes to the posterior surface of the allantois. At first in the oldest Amniotes this ureter opens into the cloaca together with the last section of the nephroduct, but afterwards separately from this, and finally into the permanent bladder apart from the rectum altogether. The bladder originates from the hindmost and lowest part of the allantoic pedicle (*urachus*), which enlarges in spindle shape before the entry into the cloaca. The anterior or upper part of the pedicle, which runs to the navel in the ventral wall of the embryo, atrophies subsequently, and only a useless string-like relic of it is left as a rudimentary organ; that is the single vesico-umbilical ligament. To the right and left of it in the adult man are a couple of other rudimentary organs, the lateral vesico-umbilical ligaments. These are the degenerate string-like relics of the earlier umbilical arteries.

Though in man and all the other Amniotes the primitive kidneys are thus early replaced by the permanent kidneys, and these alone then act as urinary organs, all the parts of the former are by no means lost. The nephroducts become very important physiologically by being converted into the passages of the sexual glands. In all the Gnathostomes—or all the Vertebrates from the fishes up to man—a second similar canal develops beside the nephroduct at an early stage of embryonic evolution. The latter is usually called the Mullerian duct, after its discoverer, Johannes Muller, while the former is called the Wolffian duct. The origin of the Mullerian duct is still obscure; comparative anatomy and ontogeny seem to indicate that it originates by differentiation from the Wolffian duct. Perhaps it would be best to say: "The

original primary nephroduct divides by differentiation (or longitudinal cleavage) into two secondary nephroducts, the Wolffian and the Mullerian ducts." The latter (Fig. 387 m) lies just on the inner side of the former (Fig. 387 w). Both open behind into the cloaca.

However uncertain the origin of the nephroduct and its two products, the Mullerian and the Wolffian ducts, may be, its later development is clear enough. In all the Gnathostomes the Wolffian duct is converted into the spermatiduct, and the Mullerian duct into the oviduct. Only one of them is retained in each sex; the other either disappears altogether, or only leaves relics in the shape of rudimentary organs. In the male sex,




FIG. 387.—Female sexual organs of a *Haptophrya* (*Crustacean*, Fig. 386). *ov.* oviducts, *bu.* bladder, *cl.* cloaca. *st.* is the outlet of the two oviducts, and between them the bladder (*bu.*) of cloaca. (From Gegenbaur.)

in which the two Wolffian ducts become the spermatiducts, we often find traces of the Mullerian ducts, which I have called "Rathke's canals" (Fig. 394 c). In the female sex, in which the two Mullerian ducts form the oviducts, there are relics of the Wolffian ducts, which are called "the ducts of Gaertner."

We obtain the most interesting information with regard to this remarkable evolution of the nephroducts and their association with the sexual glands from the Amphibia (Figs. 390-395). The first structure of the nephroduct and its differentiation into Mullerian and Wolffian ducts are just the same in both sexes in the Amphibia, as in the mammal embryos (Figs. 392, 396). In the female Amphibia

very short, and the lower common part is longer. Finally, in the ape and in man the blending of the two halves is complete, and there is only the one simple,

duct, and is found only in the ape and of the Mulleria.




FIG. 403.




FIG. 404.



FIGS. 403-404.—Origin of human ova in the female ovary. FIG. 403. Vertical section of the ovary of a newborn female infant. *a* ovarian epithelium, *b* rudimentary strand of ova, *c* young ova in the epithelium, *d* long strand of ova with follicle-formation (Follicle's index), *e* group of young follicles, *f* isolated young follicle, *g* blood-vessels in connective tissue (stroma) of the ovary. In the strand the young ova are distinguished by their considerable size from the surrounding follicle-cells. (From Naidoff.)


FIG. 404.—Two young Graafian follicles, isolated. In *a* the follicle-cells still form a single and in *b* double stratum around the young ova; in *c* they are beginning to form the outer stratum or the zona pellucida (*a*).

FIGS. 405 and 406.—Two older Graafian follicles, in which fluid is beginning to accumulate under the unilaterally thickened epithelial mass of the follicle-cells (FIG. 405 with little, and with much, follicle-steroid). *a* the young ova, with embryonic vesicle and spot, *b* cytoplasm or zona pellucida, *c* theca prolium, formed of the wall of the follicle, which surrounds the ovum, *d* follicle-liquid (antral fluid), gathered under the ovum (*b*), *e* connective-tissue stratum capsule of the Graafian follicle (theca interna).

pear-shaped uterine pouch, into which the oviducts open on each side. This simple uterus is a late evolutionary pro-

ducts at their lower ends. Here again they form a single genital cord (Fig. 397*g*), and this opens similarly into the

original urogenital sinus, which develops from the lowest section of the bladder (*v*). But while in the male mammal the Wolffian ducts develop into the permanent sperm ducts, there are only rudimentary relics left of the Müllerian ducts. The most notable of these is the "male womb" (*uterus masculinus*), which originates from the lowest fused part of the ducts, and corresponds to the female uterus. It is a small, flask-shaped vesicle without any physiological significance, which opens into the ureter between the two sperm ducts and the prostate folds (*vesicula prostatica*).



—A type human Graafian follicle. *a* the mature ovum, *b* the ovum, *c* the sperm duct, *d* the duct of the bladder, *e* the ureter, *f* the vas deferens.

The internal sexual organs of the mammal undergo very distinctive changes of position. At first the germinal glands of both sexes lie deep inside the ventral cavity, at the inner edge of the primitive kidneys (Figs. 385 *g*, 392 *h*), attached to the vertebral column by a short mesentery (*mesonephros* in the male, *mesonephros* in the female). But this primary arrangement is retained permanently only in the Monotremes (and the lower Vertebrates). In all other mammals (both Marsupial and Placental) they leave their cradle and travel more or less (for behind), following the direction of the ligament that goes from the primitive

kidneys to the inguinal region of the ventral wall. This is the inguinal ligament of the primitive kidneys, known in the male as the Hunterian ligament (Fig. 400 *g*), and in the female as the "round maternal ligament" (Fig. 401 *h*). In woman the ovaries travel more or less towards the small pelvis, or enter into it altogether. In the male the testicles pass out of the ventral cavity, and penetrate by the inguinal canal into a sac-shaped fold of the outer skin. When the right and left folds ("sexual swellings") join together they form the scrotum. The various mammae bring before us the successive stages of this displacement. In the elephant and the whale the testicles descend very little, and remain underneath the kidneys. In many of the rodents and carnivorans they enter the inguinal canal. In most of the higher mammals they pass through this into the scrotum. As a rule, the inguinal canal closes up. When the male opens it may periodically into the scrotum withdraw into the ventral cavity again. In the case of the rat (*Rattus norvegicus*) of the

piants, rodents, bats, etc.).

The structure of the external sexual organs, the copulative organs, that convey the fecundating sperm from the male to the female organism in the act of copulation, is also peculiar to the mammals. There are no organs of this character in most of the other Vertebrates. In those that live in water (such as the Acrania and Cyclostomes, and most of the fishes) the ova and sperm-cells are simply ejected into the water, where their conjunction and fertilization are left to chance. But in many of the fishes and amphibia, which are viviparous, there is a direct conveyance of the male sperm into the female body; and this is the case with all the Amniotes (reptiles, birds, and mammals). In these the urinary and sexual always open originally into the last of the rectum, which thus forms a cloaca

(p. 349). Among the mammals this arrangement is permanent only in the Monotremes, which take their name from it (Fig. 359 *cf.*). In all the other mammals a frontal partition is developed in the cloaca (in the human embryo about the beginning of the third month), and this divides it into two cavities. The anterior cavity receives the urogenital canal, and is the sole outlet of the urine and the sexual products; the hind or anus-cavity passes the excrements only.

Even before this partition has been formed in the Marsupials and Placentals, we see the first trace of the

external sexual organs. First a conical protuberance rises at the anterior border of the cloaca-outlet—the sexual prominence (*phallus*, Fig. 402 *A, c, B, c*). At the tip it is swollen in the shape of a club ("acorn" *glans*). On its under side there is a furrow, the sexual groove (*sulcus genitalis*, *f*), and on each side of this a fold of skin, the "sexual pad" (*terus genitalis*, *b, l*). The sexual protuberance or phallus is the chief organ of the sexual sense (p. 282); the sexual nerves spread on it, and these are the principal organs of the specific sexual sensation. As *erectile bodies* (*corpora cavernosa*) are developed in the male phallus by peculiar modifications of the blood-vessels, it becomes capable of erecting periodically as a strong accession of blood, becoming stiff, so

as to penetrate into the female vagina and thus effect copulation. In the male the phallus becomes the penis; in the female it becomes the much smaller clitoris; this is only found to be very large in certain apes (*Ateles*). A prepuce ("fore-skin") is developed in both sexes as a protecting fold on the anterior surface of the phallus.

The external sexual member (*phallus*) is found at various stages of development within the mammal class, both in regard to size and shape, and the differentiation and structure of its various parts; this applies especially to the terminal part of

the phallus, the glans, both the larger *glans penis* of the male and the smaller *glans clitoridis* of the female. The part of the cloaca from the upper wall of which it forms belongs to the *proctoderm*, the ectodermic invagination of the rectum (p. 312); hence its epithelial covering can develop the same horny growths as the corneous layer of the epidermis. Thus the glans, which is quite smooth in man and the higher apes, is covered with spines in many of the lower apes and in the cat, and in many of the rodents with hairs (marmoset) or scales (guinea-pig) or




FIG. 402.—The human clitoris after being treated with the Gracilis reflex, surrounded by the clitoral cells of the ductus *hygroscopus* (in two radiating rows), a cutaneous (from *phallus*) with radial, surface canals, *b* *cyrtoma* (protoplasm of the cuticular, deeper walls, higher wrinkles), *c* *ductus* of the ovum (embryonic vessels). (From *Neger*, magnified 200 times.) (*cf.* Figs. 1 and 2, pp. 36 and 45.)

solid horny warts (beaver). Many of the Ungulates have a free conical projection on the glans, and in many of the Ruminants this "phallus-tentacle" grows into a long cone, bent hook-wise at the base (as in the goat, antelope, gazelle, etc.). The different forms of the phallus are connected with variations in the structure and distribution of the sensory corpuscles—*i.e.*, the real organs of the sexual sense, which develop in certain papillae of the corium of the phallus, and have been evolved from ordinary tactile corpuscles of the corium by exotic adaptation (p. 282).

The formation of the *corpora cavernosa*, which cause the stiffness of the phallus and its capability of penetrating the vagina, by certain special structures of their spongy vascular spaces, also shows a good deal of variety within the vertebrate stem. This stiffness is increased in many orders of mammals (especially the carnassials and rodents) by the ossification of a part of the fibrous body (*corpus spongiosum*). This penis-bone (or *bone*) is very large in the badger and dog, and bent like a hook in the marten; it is also very large in some of the lower apes, and protrudes far out into the glans. It is wanting in most of the anthropoid apes; it seems to have been lost in their case (and in man) by atrophy.

The sexual groove on the under side of the phallus receives in the male the mouth of the urogenital canal, and is changed into a continuation of this, becoming a closed canal by the juncture of its parallel edges, the male urethra. In the female this only takes place in a few cases (some of the lemurs, rodents, and moles), as a rule, the groove remains open, and the borders of this "vestibule of the vagina" develop into the smaller labia (*nymphe*). The large labia of the female develop from the sexual pads (*fovee genitales*), the two parallel folds of the skin that are found on each side of the genital groove. They join together in the male, and form the closed scrotum. These striking differences between the two sexes cannot yet be detected in the human embryo of the ninth week. We begin to trace them in the tenth week of development, and they are accentuated in proportion as the difference of the sexes develops.

Sometimes the normal juncture of the two sexual pads in the male fails to take place, and the sexual groove may also remain open (*hypospadias*). In these cases the external male genitals resemble the female, and they are often wrongly regarded as cases of hermaphroditism. Other malformations of various kinds are not infrequently found in the human external sexual organs, and some of them have a great morphological interest. The reversal of *hypospadias*, in which the penis is split open below, is seen in *epispadias*, in which the urethra is open above. In this case the urogenital canal opens above at the dorsal root of the penis; in the former case down below. These and similar obstructions interfere

with a man's generative power, and thus prejudicially affect his whole development. They clearly prove that our history is not guided by a "kind Providence," but left to the play of blind chance.

We must carefully distinguish the rarer cases of real hermaphroditism from the preceding. This is only found when the essential organs of reproduction, the genital glands of both kinds, are united in one individual. In these cases either an ovary is developed on the right and a testicle on the left (or *vice versa*); or else there are testicles and ovaries on both sides, some more and others less developed. As hermaphroditism was probably the original arrangement in all the Vertebrates, and the division of the sexes only followed by later differentiation of this, these curious cases offer no theoretical difficulty. But they are rarely found in man and the higher mammals. On the other hand, we constantly find the original hermaphroditism in some of the lower Vertebrates, such as the *Myxoscolus*, many fishes of the perch-type (*perchus*), and some of the Amphibia (ringed snake, toad). In these cases the male often has a rudimentary ovary at the fore end of the testicle; and the female sometimes has a rudimentary, inactive testicle. In the carp also and some other fishes this is found occasionally. We have already seen how traces of the earlier hermaphroditism can be traced in the passages of the Amphibia.

Man has faithfully preserved the main features of his stem-history in the ontogeny of his urinary and sexual organs. We can follow their development step by step in the human embryo in the same advancing gradation that is presented to us by the comparison of the urogenital organs in the Acrania, Cyclostomes, Fishes, Amphibia, Reptiles, and then (within the mammal series) in the Monotremes, Marsupials, and the various Placentalia. All the peculiarities of urogenital structure that distinguish the mammals from the rest of the Vertebrates are found in man; and in all special structural features he resembles the apes, particularly the anthropoid apes. In proof of this fact that the special features of the mammals have been inherited by man, I will, in conclusion, point out the identical way in which the ova are formed in the ovary. In all the mammals the mature ova are contained in special capsules, which are known as the Graafian

follicles, after their discoverer, Roger de Graaf (1677). They were formerly supposed to be the ova themselves; but Baer discovered the ova within the follicles (p. 16). Each follicle (Fig. 407) consists of a round fibrous capsule (*d*), which contains fluid and is lined with several strata of cells (*c*). The layer is thickened like a knob at one point (*b*); this ovum-capsule encloses the ovum proper (*a*). The mammal ovary is originally a very simple oval body (Fig. 387 *g*), formed only of connective tissue and blood-vessels, covered with a layer of cells, the ovarian epithelium or the female germ epithelium. From this germ epithelium strings of cells grow out into the connective tissue or "stroma" of the ovary (Fig. 403 *b*). Some of the cells of these strings (or Pfliiger's tubos) grow larger and become ova (primitive ova, *c*), but the great majority remain small, and form a protective and nutritive stratum of cells round each ovum—the "follicle-epithelium" (*e*).

The follicle-epithelium of the mammal has at first one stratum (Fig. 404 *i*), but afterwards several (*g*). It is true that in all the other Vertebrates the ova are enclosed in a membrane, or "follicle," that consists of smaller cells. But it is only in the mammals that fluid accumulates between the growing follicle-cells, and distends the follicle into a large round capsule, on the inside wall of which the ovum lies, at one side (Figs. 405, 406). There again, as in the whole of his morphology, man proves indubitably his descent from the mammals.

In the lower Vertebrates the formation of ova in the germ-epithelium of the ovary continues throughout life; but in the higher it is restricted to the earlier stages, or even to the period of embryonic develop-

ment. In man it seems to cease in the first year; in the second year we find no new-formed ova or remains of ova (Pfliiger's tubos). However, the number of ova in the two ovaries is very large in the young girl; there are calculated to be 72,000 in the sexually-mature maiden. In the production of the ova men resemble most of the anthropoid apes.

Generally speaking, the natural history of the human sexual organs is one of those parts of anthropology that furnish the most convincing proofs of the animal origin of the human race. Any man who is acquainted with the facts and impartially weighs them will conclude from them alone that we have been evolved from the lower Vertebrates. The larger and the detailed structure, the action, and the embryological development of the sexual organs are just the same in man as in the apes. This applies equally to the male and the female, the internal and the external organs. The differences we find in this respect between man and the anthropoid apes are much slighter than the differences between the various species of apes. But all the apes have certainly a common origin, and have been evolved from a long-extinct early-Tertiary stem-form, which we must trace to a branch of the Lemnure. If we had this unknown pithacoid stem-form before us, we should certainly put it in the order of the true apes in the primate system; but within this order we cannot, for the anatomic and ontogenetic reasons we have seen, separate man from the group of the anthropoid apes. Here again, therefore, on the ground of the jithocometra-principle, comparative anatomy and ontogeny teach with full confidence the descent of man from the ape.

CHAPTER XXX.

RESULTS OF ANTHROPOGENY

Now that we have traversed the wonder-region of human embryology and are familiar with the principal parts of it, it will be well to look back on the way we have come, and forward to the further path to truth to which it has led us. We started from the simplest facts of ontogeny, or the development of the individual—from observations that we can repeat and verify by microscopic and anatomic study at any moment. The first and most important of these facts is that every man, like every other animal, begins his existence as a simple cell. This round ovum has the same characteristic form and origin as the ovum of any other mammal. From it is developed in the same manner in all the Placentalia, by repeated cleavage, a multicellular blastula. This is converted into a gastrula, and this in turn into a blastocystis (or embryonic vesicle). The two strata of cells that compose its wall are the primary germinal layers, the skin-layer (ectoderm), and gut-layer (entoderm). This two-layered embryonic form is the ontogenetic reproduction of the extremely important phylogenetic stem-form of all the Metazoa, which we have called the *Gastraea*. As the human embryo passes through the gastrula-form, like that of all the other Metazoa, we can trace its phylogenetic origin to the *Gastraea*.

As we continued to follow the embryonic development of the two-layered structure, we saw that first a third, or middle layer (mesoderm), appears between the two primary layers; when this divides into two, we have the four secondary germinal layers. These have just the same composition and genetic significance in man as in all the other Vertebrates. From the skin-sense layer are developed the epidermis, the central nervous system, and the chief part of the sense-organs. The skin-fibre layer forms the corium and the motor organs—the skeleton and the muscular system. From the gut-fibre layer are developed the vascular system, the muscular wall of the gut, and the sexual glands. Finally, the gut-stand

layer only forms the epithelium, or the inner cellular stratum of the mucous membrane of the alimentary canal and glands (lungs, liver, etc.).

The manner in which these different systems of organs arise from the secondary germinal layers is essentially the same from the start in man as in all the other Vertebrates. We saw, in studying the embryonic development of each organ, that the human embryo follows the special lines of differentiation and construction that are only found otherwise in the Vertebrates. Within the limits of this vast stem we have followed, step by step, the development both of the body as a whole and of its various parts. This higher development follows in the human embryo the form that is peculiar to the mammals. Finally, we saw that, even within the limits of this class, the various phylogenetic stages that we distinguish in a natural classification of the mammals correspond to the ontogenetic stages that the human embryo passes through in the course of its evolution. We were thus in a position to determine precisely the position of man in this class, and so to establish his relationship to the different orders of mammals.

The line of argument we followed in this explanation of the ontogenetic facts was simply a consistent application of the biogenetic law. In this we have throughout taken strict account of the distinction between pallogenic and cenogenetic phenomena. Pallogenesis (or "synoptic development") alone enables us to draw conclusions from the observed embryonic form to the stem-form preserved by heredity. Such inference becomes more or less precarious when there has been cenogenesis, or disturbance of development, owing to fresh adaptations. We cannot understand embryonic development unless we appreciate this very important distinction. Here we stand at the very limit that separates the older and the new science or philosophy of nature. The whole of the results of recent morphological research counsel us irresistibly

to recognise the biogenetic law and its far-reaching consequences. These are, it is true, irreconcilable with the legends and doctrines of former days, that have been impressed on us by religious education. But without the *biogenetic law*, without the distinction between *palaogenesis* and *ontogenesis*, and without the theory of *evolution* on which we base it, it is quite impossible to understand the facts of organic development; without them we cannot cast the faintest gleam of explanation over this marvellous field of phenomena. But when we recognise the causal correlation of ontogeny and phylogeny expressed in this law, the wonderful facts of embryology are susceptible of a very simple explanation; they are found to be the necessary mechanical effects of the evolution of the stem, determined by the laws of heredity and adaptation. The correlative action of these laws under the universal influence of the struggle for existence, or—as we may say in a word, with Darwin—"natural selection," is entirely adequate to explain the whole process of embryology in the light of phylogeny. It is the chief merit of Darwin that he explained by his theory of selection the correlation of the laws of heredity and adaptation that Lamarck had recognised, and pointed out the true way to reach a causal interpretation of evolution.

The phenomenon that it is most imperative to recognise in this connection is the inheritance of functional variations. Jean Lamarck was the first to appreciate its fundamental importance in 1809, and we may therefore justly give the name of Lamarckism to the theory of descent he based on it. Hence the radical opponents of the latter have very properly directed their attacks chiefly against the former. One of the most distinguished and most narrow-minded of these opponents, Wilhelm Kilm, affirms very positively that "characteristics acquired in the life of the individual are not inherited."

The inheritance of acquired characters is denied, not only by thorough opponents of evolution, but even by scientists who admit it and have contributed a good deal to its establishment, especially Weismann, Galton, Ray Lankester, etc. Since 1884, the chief opponent has been August Weismann, who has rendered the greatest service in the development of Darwin's theory of selection. In his work on *The Continuity of the Germ-*

Plasm, and in his recent excellent *Lectures on the Theory of Descent* (1902), he has with great success advanced the opinion that "only those characters can be transmitted to subsequent generations that were contained in rudimentary form in the embryo." However, this germ-plasm theory, with its attempt to explain heredity, is merely a "provisional molecular hypothesis"; it is one of those metaphysical speculations that attribute the evolutionary phenomena exclusively to internal causes, and regard the influence of the environment as insignificant. Herbert Spencer, Theodor Eimer, Lester Ward, Hering, and Zehnder have pointed out the untenable consequences of this position. I have given my view of it in the tenth edition of the *History of Creation* (pp. 191, 203). I hold, with Lamarck and Darwin, that the hereditary transmission of acquired characters is one of the most important phenomena in biology, and is proved by thousands of morphological and physiological experiences. It is an indispensable foundation of the theory of evolution.

Of the many and weighty arguments for the truth of this conception of evolution I will for the moment merely point to the invaluable evidence of dysmelody, the science of rudimentary organs. We cannot insist too often or too strongly on the great morphological significance of these remarkable organs, which are completely useless from the physiological point of view. We find some of these useless parts, inherited from our lower vertebrate ancestors, in every system of organs in man and the higher Vertebrates. Thus we find at once on the skin a scanty and rudimentary coat of hair, only fully developed on the head, under the shoulders, and at a few other parts of the body. The short hairs on the greater part of the body are quite useless and devoid of physiological value; they are the last relic of the thicker hairy coat of our simian ancestors. The sensory apparatus prevents a series of most remarkable rudimentary organs. We have seen that the whole of the shell of the external ear, with its cartilages, muscles, and skin, is in man a useless appendage, and has not the physiological importance that was formerly ascribed to it. It is the degenerate remainder of the pointed, freely moving, and more advanced mammal ear, the muscles of which we still have, but cannot work them. We found at the

inner corner of our eye a small, curious, semi-lunar fold that is of no use whatever to us, and is only interesting as the last relic of the nictitating membrane, the third, inner eye-lid that had a distinct physiological purpose in the ancient sharks, and still has in many of the Amniotes.

The motor apparatus, in both the skeleton and muscular systems, provides a number of interesting dysteleological arguments. I need only recall the projecting tail of the human embryo, with its rudimentary caudal vertebrae and muscles; this is totally useless in man, but very interesting as the degenerate relic of the long tail of our simian ancestors. From these we have also inherited various bony processes and muscles, which were very useful to them in climbing trees, but are useless to us. At various points of the skin we have cutaneous muscles which we never use—remnants of a strongly-developed cutaneous muscle in our lower mammal ancestors. This "cutaneous carnosus" had the function of contracting and creasing the skin to chase away the flies, as we see every day in the horse. Another relic in us of this large cutaneous muscle is the frontal muscle, by which we knit our forehead and raise our eye-brows, but there is another considerable relic of it, the large cutaneous muscle in the neck (*platysma myoides*), over which we have no voluntary control.

Not only in the systems of animal organs, but also in the vegetative apparatus, we find a number of rudimentary organs, many of which we have already noticed. In the alimentary apparatus there are the thymus-gland and the thyroid gland, the seat of goitre and the relic of a ciliated groove that the Tunicates and Acrasias still have in the gill-pump; there is also the vermiform appendix to the caecum. In the vascular system we have a number of useless cords which represent relics of atrophied vessels that were once active as blood-canals—the *ductus Botalli* between the pulmonary artery and the aorta, the *ductus venosus Arantii* between the portal vein and the vena cava, and many others. The many rudimentary organs in the urinary and sexual apparatus are particularly interesting. These are generally developed in one sex and rudimentary in the other. Thus the spermatheca is formed from the Wolffian ducts in the male, whereas in the female we have merely rudimentary

traces of them in Gartner's canals. On the other hand, in the female the oviducts and womb are developed from the Mullerian ducts, while in the male only the lowest cords of them remain as the "male womb" (*foresculla prosterhus*). Again, the male has in his nipples and mammary glands the rudiments of organs that are usually active only in the female.

A careful anatomical study of the human frame would disclose to us numbers of other rudimentary organs, and these can only be explained on the theory of evolution. Robert Wiedersheim has collected a large number of them in his work on *The Human Frame as a Witness to its Past*. They are some of the weightiest proofs of the truth of the mechanical conception and the strongest disproofs of the teleological view. If, as the latter demands, man or any other organism had been designed and fitted for his life-purposes from the start and brought into being by a creative act, the existence of these rudimentary organs would be an insoluble enigma; it would be impossible to understand why the Creator had put this useless burden on his creatures to walk a path that is in itself by no means easy. But the theory of evolution gives the simplest possible explanation of them. It says: The rudimentary organs are parts of the body that have fallen into disuse in the course of centuries, they had definite functions in our animal ancestors, but have lost their physiological significance. On account of fresh adaptations they have become superfluous, but are transmitted from generation to generation by heredity, and gradually atrophy.

We have inherited not only these rudimentary parts, but all the organs of our body, from the mammals—proximately from the apes. The human body does not contain a single organ that has not been inherited from the apes. In fact, with the aid of our biogenetic law we can trace the origin of our various systems of organs much further, down to the lowest stages of our ancestry. We can say, for instance, that we have inherited the oldest organs of the body, the external skin and the internal coat of the alimentary system, from the Coelenterates; the nervous and muscular systems from the Platyhelminths; the vascular system, the body-cavity, and the blood from the Vermes; the chorda and the bronchial gut from the Prochordonia;

the articulation of the body from the Acrania; the primitive skull and the higher sense-organs from the Cyclostomes; the limbs and jaws from the Selachii; the five-toed foot from the Amphibia; the palate from the Reptiles; the hairy coat, the mammary glands, and the external sexual organs from the Mammals. When we formulated "the law of the ontogenetic connection of systematically related forms," and determined the relative age of organs, we saw how it was possible to draw phylogenetic conclusions from the ontogenetic succession of systems of organs.

With the aid of this important law and of comparative anatomy we were also enabled to determine "man's place in nature," or, as we put it, assign to man his position in the classification of the animal kingdom. In recent zoological classification the animal world is divided into twelve stems or phyla, and these are broadly sub-divided into about sixty classes, and these classes into at least 300 orders. In his whole organisation man is most certainly, in the first place, a member of one of these stems, the vertebrate stem; secondly, a member of one particular class in this stem, the Mammals, and thirdly, of one particular order, the order of Primates. He has all the characteristics that distinguish the Vertebrates from the other eleven animal stems, the Mammals from the other sixty classes, and the Primates from the 300 other orders of the animal kingdom. We may turn and twist as we like, but we cannot get over this fact of anatomy and classification. Of late years this fact has given rise to a good deal of discussion, and especially of controversy as to the particular anatomic relationship of man to the apes. The most curious opinions have been advanced on this "ape-question," or "pithecolith-theory." It is as well, therefore, to go into it once more and distinguish the essential from the unessential. (Cf. above, pp. 262-5.)

We start from the undisputed fact that man is in any case—whether we accept or reject his special blood-relationship to the apes—a true mammal; in fact, a placental mammal. This fundamental fact can be proved so easily at any moment from comparative anatomy that it has been universally admitted since the separation of the Placentals from the lower mammals (Marsupials and Monotremes). But for every consistent sub-

scriber to the theory of evolution it must follow at once that man descends from a common stem-form with all the other Placentals, the stem-ancestor of the Placentals, just as we must admit a common mesozoic ancestor of all the mammals. This is, however, to settle decisively the great and burning question of man's place in nature, whether or no we go on to admit a nearer or more distant relationship to the apes. Whether man is or is not a member of the ape-order (or, if you prefer, the primate-order) in the phylogenetic sense, in any case his direct blood-relationship to the rest of the mammals, and especially the Placentals, is established. It is possible that the affinities of the various orders of mammals to each other are different from what we hypothetically assume to-day. But, in any case, the common descent of man and all the other mammals from one stem-form is beyond question. This long-extinct Protomammal was probably evolved from Proreptiles during the Triassic period, and must certainly be regarded as the monotreme and oviparous ancestor of all the mammals.

If we hold firmly to this fundamental and most important thesis, we shall see the "ape-question" in a very different light from that in which it is usually regarded. Little reflection is then needed to see that it is not nearly so important as it is said to be. The origin of the human race from a series of mammal ancestors, and the historic evolution of these from an earlier series of lower vertebrate ancestors, together with all the weighty conclusions that every thoughtful man deduces therefrom, remain untouched; so far as these are concerned, it is immaterial whether we regard true "apes" as our nearest ancestors or not. But as it has become the fashion to lay the chief stress in the whole question of man's origin on the "descent from the apes," I am compelled to return to it once more, and recall the facts of comparative anatomy and ontogeny that give a decisive answer to this "ape-question."

The shortest way to attain our purpose is that followed by Huxley in 1863 in his able work, which I have already often quoted, *Man's Place in Nature*—the way of comparative anatomy and ontogeny. We have to compare impartially all man's organs with the same organs in the higher apes, and then to examine if the differences between the two are greater

than the corresponding differences between the higher and the lower apes. The indubitable and incontrovertible result of this comparative-anatomical study, conducted with the greatest care and impartiality, was the *pithecometric*-principle, which we have called the Huxleyan law in honour of its formulator—namely, that the differences in organisation between man and the most advanced apes we know are much slighter than the corresponding differences in organisation between the higher and lower apes. We may even give a more precise formula to this law, by excluding the Platyrrhines or American apes as distant relatives, and restricting the comparison to the narrower family-circle of the Catarrhines, the apes of the Old World. Within the limits of this small group of mammals we found the structural differences between the lower and higher catarrhine apes—for instance, the baboon and the gorilla—to be much greater than the differences between the anthropoid apes and man. If we now turn to ontogeny, and find, according to our "law of the ontogenetic connection of systematically related forms," that the embryos of the anthropoid apes and man retain their resemblance for a longer time than the embryos of the highest and the lowest apes, we are forced, whether we like it or no, to recognise our descent from the order of apes. We can assuredly construct an approximate picture in the imagination of the form of our early Tertiary ancestors from the foregoing facts of comparative anatomy; however we may frame this in detail, it will be the picture of a true ape, and a distinct catarrhine ape. This has been shown so well by Huxley (1863) that the recent attacks of Klnasch, Varchow, and other anthropologists, have completely failed (cf. pp. 263-264). All the structural characters that distinguish the Catarrhines from the Platyrrhines are found in man. Hence in the genealogy of the mammals we must derive man immediately from the catarrhine group, and locate the origin of the human race in the Old World. Only the early root-form from which both descended was common to them.

It is, therefore, established beyond question for all impartial scientific inquiry that the human race comes directly from the apes of the Old World; but, at the same time, I repeat that this is not an important in connection with the main

question of the origin of man as is commonly supposed. Even if we entirely ignore it, all that we have learned from the zoological facts of comparative anatomy and ontogeny as to the placental character of man remains untouched. These prove beyond all doubt the common descent of man and all the rest of the mammals. Further, the main question is not in the least affected if it is said: "It is true that man is a mammal; but he has diverged at the very root of the class from all the other mammals, and has no closer relationship to any living group of mammals." The affinity is more or less close in any case, if we examine the relation of the mammal class to the sixty other classes of the animal world. Quite certainly the whole of the mammals, including man, have had a common origin; and it is equally certain that their common stem-forms were gradually evolved from a long series of lower Vertebrates.

The resistance to the theory of a descent from the apes is clearly due in most men to feeling rather than to reason. They shrink from the notion of such an origin just because they see in the ape organism a caricature of man, a distorted and unattractive image of themselves; because it hurts man's æsthetic complacency and self-embellishment. It is more flattering to think we have descended from some lofty and god-like being; and so, from the earliest times, human vanity has been pleased to believe in our origin from gods or demi-gods. The Church, with that sophist's reversal of ideas of which it is a master, has succeeded in representing this ridiculous piece of vanity as "Christian humility", and the very men who reject with horror the notion of an animal origin, and count themselves "children of God," love to prate of their "humble sense of servitude." In most of the sermons that have poured out from pulpit and altar against the doctrine of evolution human vanity and conceit have been a conspicuous element; and, although we have inherited this very characteristic weakness from the apes, we must admit that we have developed it to a higher degree, which is entirely repudiated by sound and normal intelligence. We are greatly amused at all the childish follies that the ridiculous pride of ancestry has maintained from the Middle Ages to our own times; yet there is a large amount of this empty feeling in

most men. Just as most people much prefer to trace their family back to some degenerate baron or some famous prince rather than to an unknown peasant, so most men would rather have as parent of the race a sinful and fallen Adam than an advancing and vigorous ape. It is a matter of taste, and to that extent we cannot quarrel over these genealogical tendencies. Personally, the notion of descent is more congenial to me than that of descent. It seems to me a finer thing to be the advanced offspring of a shrunken ancestor, that has developed progressively from the lower mammals in the struggle for life, than the degenerate descendant of a god-like being, made from a clod, and fallen for his sins, and an Eve created from one of his ribs. Speaking of the rib, I may add to what I have said about the development of the skeleton, that the number of ribs is just the same in man and woman. In both of them the ribs are formed from the middle germinal layer, and are, from the phylogenetic point of view, lower or ventral vertebral arches.

But it is said: "That is all very well, as far as the human body is concerned; on the facts quoted it is impossible to doubt that it has really and gradually been evolved from the long ancestral series of the Vertebrates. But it is quite another thing as regards man's mind, or soul; this cannot possibly have been developed from the vertebrate-soul." Let us see if we cannot meet this grave stricture from the well-known facts of comparative anatomy, physiology, and embryology. It will be best to begin with a comparative study of the souls of various groups of Vertebrates. Here we find such an enormous variety of vertebrate souls that, at first sight, it seems quite impossible to trace them all to a common "Primitive Vertebrate." Think of the tiny Amphioxus, with no real brain but a simple medullary tube, and its whole psychic life at the very lowest stage among the Vertebrates. The following group of the Cyclostomes are still very limited, though they have a brain. When we pass on to the fishes, we find their intelligence remaining at a very low level. We do not see any material advance in mental development until we go on to the Amphibia and Reptiles. There is

still greater advance when we come to the Mammals, though even here the minds of the Monotremes and of the stupid Marsupials remain at a low stage. But when we rise from these to the Placentals we find within this one vast group such a number of important stages of differentiation and progress that the psychic differences between the least intelligent (such as the sloths and armadillos) and the most intelligent Placentals (such as the dogs and apes) are much greater than the psychic differences between the lowest Placentals and the Marsupials or Monotremes. Most certainly the differences are far greater than the differences in mental power between the dog, the ape, and man. Yet all these animals are genetically-related members of a single natural class.

We see this to a still more astonishing extent in the comparative psychology of another class of animals, that is especially interesting for many reasons—the insect class. It is well known that we find in many insects a degree of intelligence that is found in man alone among the Vertebrates. Everybody knows of the famous communities and states of bees and ants, and of the very remarkable social arrangements in them, such as we find among the more advanced races of men, but among no other group of animals. I need only mention the social organization and government of the monarchic bees and the republican ants, and their division into different conditions—queen, drone-nobles, workers, educators, soldiers, etc. One of the most remarkable phenomena in this very interesting province is the cattle-keeping of the ants, which rear plant-lice as milk-cows and regularly extract their honied juice. Still more remarkable is the slave-holding of the large red ants, which steal the young of the small black ants and bring them up as slaves. It has long been known that these political and social arrangements of the ants are due to the deliberate co-operation of the countless citizens, and that they understand each other. A number of recent observers, especially Fritz Muller, Sir J. Lubbock (Lord Avebury), and August Forel, have put the astonishing degree of intelligence of these tiny Articulates beyond question.

Now, compare with these the mental life of many of the lower, especially the parasitic, insects, as Darwin did. There is, for instance, the cochineal insect

* The English reader will recognize from the German edition of Dr. Wallace and of the late Dr. Mearns—*Trans.*

(*Coccus*), which, in its adult state, has a motionless, shield-shaped body, attached to the leaves of plants. Its feet are atrophied. Its mouth is sunk in the tissue of the plants of which it absorbs the sap. The whole psychic life of these inert female parasites consists in the pleasure they experience from sucking the sap of the plant and in sexual intercourse with the males. It is the same with the maggot-like females of the fly (*Strep-siva*), which spend their lives parasitically and immovably, without wings or feet, in the abdomen of wasps. There is no question here of higher psychic action. If we compare these sluggish parasites with the intelligent and active ants, we must admit that the psychic differences between them are much greater than the psychic differences between the lowest and highest mammals, between the Monotremes, Marsupials, and Armadillos on the one hand, and the dog, ape, or man on the other. Yet all these insects belong to the same class of Articulata, just as all the mammals belong to one and the same class. And just as every consistent evolutionist must admit a common stem-form for all these insects, so he must also for all the mammals.

If we now turn from the comparative study of psychic life in different animals to the question of the organs of this function, we receive the answer that in all the higher animals they are always bound up with certain groups of cells, the ganglionic cells or neurons that compose the nervous system. All scientists without exception are agreed that the central nervous system is the organ of psychic life in the animal, and it is possible to prove this experimentally at any moment. When we partially or wholly destroy the central nervous system, we extinguish in the same proportion, partially or wholly, the "soul" or psychic activity of the animal. We have, therefore, to examine the features of the psychic organ in man. The reader already knows the incontestable answer to this question. Man's psychic organ is, in structure and origin, just the same organ as in all the other Vertebrates. It originates in the shape of a simple medullary tube from the outer membrane of the embryo—the skin-sense layer. The simple cerebral vesicle that is formed by the expansion of the head-part of this medullary tube divides by transverse constrictions into five, and these pass through more or less the same stages

of construction in the human embryo as in the rest of the mammals. As these are undoubtedly of a common origin, their brain and spinal cord must also have a common origin.

Physiology teaches us further, on the ground of observation and experiment, that the relation of the "soul" to its organ, the brain and spinal cord, is just the same in man as in the other mammals. The one cannot act at all without the other; it is just as much bound up with it as muscular movement is with the muscles. It can only develop in connection with it. If we are evolutionists at all, and grant the causal connection of ontogenesis and phylogenesis, we are forced to admit this thesis: The human soul or psyche, as a function of the medullary tube, has developed along with it, and just as brain and spinal cord now develop from the simple medullary tube in every human individual, so the human mind or the psychic life of the whole human race has been gradually evolved from the lower vertebrate soul. Just as to-day the intricate structure of the brain proceeds step by step from the same rudiment in every human individual—the same five cerebral vesicles—as in all the other Craniotes, so the human soul has been gradually developed in the course of millions of years from a long series of trinitate-souls. Finally, just as to-day in every human embryo the various parts of the brain differentiate after the special type of the ape-brain, so the human psyche has proceeded historically from the ape-soul.

It is true that this Monistic conception is rejected with horror by most men, and the Dualistic idea, which denies the inseparable connection of brain and mind, and regards body and soul as two totally different things, is still popular. But how can we reconcile this view with the known facts of evolution? It meets with difficulties equally great and insuperable in embryology and in phylogeny. If we suppose with the majority of men that the soul is an independent entity, which has nothing to do with the body originally, but merely inhabits it for a time, and gives expression to its experiences through the brain just as the pianist does through his instrument, we must assign a point in human embryology at which the soul enters into the brain; and at death again we must assign a moment at which it abandons the body. As, further, each human individual has inherited certain

personal features from each parent, we must suppose that in the act of conception pieces were detached from their souls and transferred to the embryo. A piece of the paternal soul goes with the spermatozoon, and a piece of the mother's soul remains in the ovum. At the moment of conception, when portions of the two nuclei of the copulating cells join together to form the nucleus of the stem-cell, the accompanying fragments of the immaterial souls must also be supposed to coalesce.

On this Dualistic view the phenomena of psychic development are totally incomprehensible. Everybody knows that the new-born child has no consciousness, no knowledge of itself and the surrounding world. Every parent who has impartially followed the mental development of his children will find it impossible to deny that it is a case of biological evolutionary processes. Just as all other functions of the body develop in connection with their organs, so the soul does in connection with the brain. This gradual unfolding of the soul of the child is, in fact, so wonderful and glorious a phenomenon that every mother or father who has eyes to observe is never tired of contemplating it. It is only our manuals of psychology that know nothing of this development; we are almost tempted to think sometimes that their authors can never have had children themselves. The human soul, as described in most of our psychological works, is merely the soul of a learned philosopher, who has read a good many books, but knows nothing of evolution, and never even reflects that his own soul has had a development.

When these Dualistic philosophers are consistent they must assign a moment in the phylogeny of the human soul at which it was first "introduced" into man's vertebrate body. Hence, at the time when the human body was evolved from the anthropoid body of the ape (probably in the Tertiary period), a specific human psychic element—or, as people love to say, "a spark of divinity"—must have been suddenly infused or breathed into the anthropoid brain, and been associated with the ape-soul already present in it. I need not insist on the enormous theoretical difficulties of this idea. I will only point out that this "spark of divinity," which is supposed to distinguish the soul of man from that of the other animals, must be itself capable of development,

and has, as a matter of fact, progressively developed in the course of human history. As a rule, reason is taken to be this "spark of divinity," and is supposed to be an exclusive possession of humanity. But comparative psychology shows us that it is quite impossible to set up this barrier between man and the bruta. Either we take the word "reason" in the wider sense, and then it is found in the higher mammals (ape, dog, elephant, horse) just as well as in most men; or else in the narrower sense, and then it is lacking in most men just as much as in the majority of animals. On the whole, we may still say of man's reason what Goethe's Mephistopheles said:—

Loft somewhat better might content him
But for the gleam of heavenly light that
Thou hast given him.
He calls it reason: thence his power's
Increased
To be still baselier than any beast.

If, then, we must reject these popular and, in some respects, agreeable Dualistic theories as untenable, because inconsistent with the genetic facts, there remains only the opposite or Monistic conception, according to which the human soul is, like any other animal soul, a function of the central nervous system, and develops in inseparable connection therewith. We see this ontogenetically in every child. The biogenetic law compels us to affirm it phylogenetically. Just as in every human embryo the skin-sense layer gives rise to the medullary tube, from the anterior end of which the five cerebral vesicles of the Cranium are developed, and from these the mammal brain (first with the characters of the lower, then with those of the higher mammals); and as the whole of this ontogenetic process is only a brief, hereditary reproduction of the same process in the phylogenesis of the Vertebrates; so the wonderful spiritual life of the human race through many thousands of years has been evolved step by step from the lowly psychic life of the lower Vertebrates, and the development of every child-soul is only a brief repetition of that long and complex phylogenetic process. From all these facts sound reason must conclude that the still prevalent belief in the immortality of the soul is an untenable superstition. I have shown its inconsistency with modern science in the eleventh chapter of *The Huddle of the Universe*.

Here it may also be well to point out

the great importance of anthropogeny, in the light of the biogenetic law, for the purposes of philosophy. The speculative philosophers who take cognizance of these ontogenetic facts, and explain them (in accordance with the law) phylogenetically, will advance the great questions of philosophy far more than the most distinguished thinkers of all ages have yet succeeded in doing. Most certainly every clear and consistent thinker must derive from the facts of comparative anatomy and ontogeny we have adduced a number of suggestive ideas that cannot fail to have an influence on the progress of philosophy. Nor can it be doubted that the candid statement and impartial appreciation of these facts will lead to the decisive triumph of the philosophic tendency that we call "Monistic" or "Mechanical," as opposed to the "Dualistic" or "Teleological," on which most of the ancient, medieval, and modern systems of philosophy are based. The Monistic or Mechanical philosophy affirms that all the phenomena of human life and of the rest of nature are ruled by fixed and unalterable laws; that there is everywhere a necessary causal connection of phenomena; and that, therefore, the whole knowable universe is a harmonious unity, a *cosmos*. It says, further, that all phenomena are due solely to mechanical or efficient causes, not to final causes. It does not admit free-will in the ordinary sense of the word. In the light of the Monistic philosophy the phenomena that we are wont to regard as the freest and most independent, the expressions of the human will, are subject just as much to rigid laws as any other natural phenomenon. As a matter of fact, impartial and thorough examination of our "free" volitions shows that they are never really free, but always determined by antecedent factors that can be traced to either heredity or adaptation. We cannot, therefore, admit the conventional distinction between nature and spirit. There is spirit everywhere in nature, and we know of no spirit outside of nature. Hence, also, the common antithesis of natural science and mental or moral science is untenable. Every science, as such, is both natural and mental. That is a firm principle of Monism, which, on its religious side, we may also denominate Pantheism. Man is not above, but in, nature.

It is true that the opponents of evolu-

tion love to misrepresent the Monistic philosophy based on it as "Materialism," and confuse the philosophic tendency of this name with a wholly unconnected and despicable moral materialism. Strictly speaking, it would be just as proper to call our system Spiritualism as Materialism. The real Materialistic philosophy affirms that the phenomena of life are, like all other phenomena, effects or products of matter. The opposite extreme, the Spiritualistic philosophy, says, on the contrary, that matter is a product of energy, and that all material forms are produced by free and independent forces. Thus, according to one-sided Materialism, the matter is antecedent to the living force; according to the equally one-sided view of the Spiritist, it is the reverse. Both views are Dualistic, and, in my opinion, both are false. For us the antithesis disappears in the Monistic philosophy, which knows neither matter without force nor force without matter. It is only necessary to reflect for some time over the question from the strictly scientific point of view to see that it is impossible to form a clear idea of either hypothesis. As Goethe said, "Matter can never exist or act without spirit, nor spirit without matter."

The human "spirit" or "soul" is merely a force or form of energy, inseparably bound up with the material substratum of the body. The thinking force of the mind is just as much connected with the structural elements of the brain as the motor force of the muscles with their structural elements. Our mental powers are functions of the brain as much as any other force is a function of a material body. We know of no matter that is devoid of force, and no forces that are not bound up with matter. When the forces enter into the phenomenon as movements we call them living or active forces; when they are in a state of rest or equilibrium we call them latent or potential. This applies equally to inorganic and organic bodies. The magnet that attracts iron filings, the powder that explodes, the steam that drives the locomotive, are living inorganics; they act by living force as much as the sensitive Mimosa does when it contracts its leaves at touch, or the venerable Amphioxus that buries itself in the sand of the sea, or man when he thinks. Only in the latter cases the combinations of the different forces that appear as "movement" in the

phenomenon are much more intricate and difficult to analyse than in the former.

Our study has led us to the conclusion that in the whole evolution of man, in his embryology and in his phylogeny, there are no living forces at work other than those of the rest of organic and inorganic nature. All the forces that are operative in it could be reduced in the ultimate analysis to growth, the fundamental evolutionary function that brings about the forms of both the organic and the inorganic. But growth itself depends on the attraction and repulsion of homogeneous and heterogeneous particles. seventy-five years ago Carl Ernst von Baer summed up the general result of his classic studies of animal development in the sentence "The evolution of the individual is the history of the growth of individuality in every respect." And if we go deeper to the root of this law of growth, we find that in the long run it can always be reduced to that attraction and repulsion of animated atoms which Empedocles called the "love and hatred" of the elements.

Thus the evolution of man is directed by the same "eternal, iron laws" as the development of any other body. These laws always lead us back to the same simple principles, the elementary principles of physics and chemistry. The various phenomena of nature only differ in the degree of complexity in which the different forces work together. Each single process of adaptation and heredity in the stem-history of our ancestors is in itself a very complex physiological phenomenon. Far more intricate are the processes of human embryology; in these are condensed and comprised thousands of the phylogenetic processes.

In my *General Morphology*, which appeared in 1866, I made the first attempt to apply the theory of evolution, as reformulated by Darwin, to the whole province of biology, and especially to provide with its assistance a mechanical foundation for the science of organic forms. The intimate relations that exist between all parts of organic science, especially the direct causal nexus between the two sections of evolution—ontogeny and phylogeny—were explained in that work for the first time by transformism, and were interpreted philosophically in the light of the theory of descent. The anthropological part of the *General Morphology* (Book vii.) contains the first attempt to

determine the series of man's ancestors (vol. ii., p. 448). However imperfect this attempt was, it provided a starting-point for further investigation. In the thirty-seven years that have since elapsed the biological horizon has been enormously widened, our empirical acquisitions in paleontology, comparative anatomy, and ontogeny have grown to an astonishing extent, thanks to the united efforts of a number of able workers and the employment of better methods. Many important biological questions that then appeared to be obscure enigmas seem to be entirely settled. Darwinism arose like the dawn of a new day of clear Monistic science after the dark night of mystic dogmatism, and we can say now, proudly and gladly, that there is daylight in our field of inquiry.

Philosophers and others, who are equally ignorant of the empirical sources of our evidence and the phylogenetic methods of utilizing it, have even lately claimed that in the matter of constructing our genealogical tree nothing more has been done than the discovery of a "gallery of ancestors," such as we find in the mansions of the nobility. This would be quite true if the genealogy given in the second part of this work were merely the juxtaposition of a series of animal forms, of which we gathered the genetic connection from their external physiognomic resemblances. As we have sufficiently proved already, it is for us a question of a totally different thing—of the morphological and historical proof of the phylogenetic connection of these ancestors on the basis of their identity in internal structure and embryonic development; and I think I have sufficiently shown in the first part of this work how far this is calculated to reveal to us their inner nature and its historical development. I see the essence of its significance precisely in the proof of historical connection. I am one of those scientists who believe in a real "natural history," and who think as much of an historical knowledge of the past as of an exact investigation of the present. The incalculable value of the historical consciousness cannot be sufficiently emphasised at a time when historical research is ignored and neglected, and when an "exact" school, as dogmatic as it is narrow, would substitute for it physical experiments and mathematical formulae. Historical knowledge cannot be replaced by any other branch of science.

It is clear that the prejudices that stand in the way of a general recognition of this "natural anthropogeny" are still very great; otherwise the long struggle of philosophic systems would have ended in favour of Monism. But we may confidently expect that a more general acquaintance with the genetic facts will gradually destroy these prejudices, and lead to the triumph of the natural conception of "man's place in nature." When we hear it said, in face of this expectation, that this would lead to retrogression in the intellectual and moral development of mankind, I cannot refrain from saying that, in my opinion, it will be just the reverse; that it will promote

to an enormous extent the advance of the human mind. All progress in our knowledge of truth means an advance in the higher cultivation of the human intelligence; and all progress in its application to practical life implies a corresponding improvement of morality. The worst enemies of the human race—ignorance and superstition—can only be vanquished by truth and reason. In any case, I hope and desire to have convinced the reader of these chapters that the true scientific comprehension of the human frame can only be attained in the way that we recognise to be the sole sound and effective one in organic science generally—namely, the way of Evolution.

INDEX

- ASTROCYTES**, 26
Astragalin, 234
 Abortive ova, 55
 Achromatin, 42
 Achroman, 42
 Acmla, 231
 Acoustic nerve, the, 269, 290
 Acquired characters, inheritance of, 349
 Acromia, the, 183, 229
 Auroganglion, the, 268, 275
 Adam's apple, the, 184
 Adapsida, 257
 Adaptation, 3, 5, 27
 After-birth, the, 167
 Agassiz, L., 34
 Age of life, 200
 Alimentary canal, evolution of the, 13, 14, 133, 208-17
 — structure of the, 169, 208-10
 Allantoic circulation, the, 171
 Allantois, development of the, 166
 Allmann, 20
Alphagomus, 243
 Anatomic cleavage, 40
 Anamniota, 217
Anamniotes, 217
 Anaxion, the, 115
 — formation of the, 131, 244
 Anaxion fluid, the, 134
 Anaxion, the, 47-9, 110
 Amphibia, the, 239
Amphibians, 231
 Amphigastria, 80
 Amphioxus, the, 105, 181-95
 — circulation of the, 184
 — colonization of the, 95
 — embryology of the, 191-95
 — structure of the, 183-88
 Amphioxus, 230
 Anaxion, the, 115
 Anatomy, comparative, 208
 Animalculists, 12
 Animal layer, the, 16
 Annelids, the, 121, 219
 Annelid theory, the, 143
 Anomodontia, 245
 Ant, intelligence of the, 353
Anthropanthus, 174, 262
 Anthropogeny, 1
 Anthropoid apes, the, 166, 173, 262
 Anthropology, 1, 35
 Anthropogenic period, 203
 Antism, 207
 Anura, 243
 Anura, the, 217
 Anus, formation of the, 199
 Aorta, the, 227
 — development of the, 170
 Age and man, 157, 164, 261, 307, 351
 Ape-man, the, 263, 264
 Apes, the, 257-60
Aphanogaster, 210
Aphanogaster, 221
 Appendiculus, 197
 Appendix vermiformis, the, 31
 Aquatic life, early prevalence of, 235
 Ararat, Mount, 24
 Archenteron, 64, 74
 Archeolithic age, 203
 Archicaryon, 55
 Archicarya, 230
 Archigastria, 64, 193
Archiparas, 263
Archiparas, 261
 Area, the germinal, 127
 Aristotle, 9
 Arms, structure of the, 205
 Arrow-worm, the, 191
 Articular arches, the, 245-26
 — cone, the, 244
 Arteries, evolution of the, 170, 223-24
 Articulation, the, 122, 219
 — skeleton of the, 294
 Articulation, 141-42
 Aryo - Romanic languages, the, 203
 Ascidia, the, 181, 185-90
 — embryology of the, 195-98
 Ascuta, 217
 Ascutal reproduction, 51
 Atlas, the, 247
 Atlas, the, 183, 185
 — (head), the, 246
 Auditory nerve, the, 289, 290
 Auricles of the heart, 325
Autolemnus, 257
 Axolotl, the, 243
 Bacteria, 38, 210
 Baer, K. E. von, 12-17
 Balanoglossus, 226
 Balour, F., 21
 Batrachia, 241
Batrachomys Stenot, 78
 Bec. generations of the, 9
 Bismarck, W., on evolution, 50
 Bilateral symmetry, 66
 — origin of, 221
 Bismarck, 258
 Biogenetic law, the, 2, 21, 23, 275, 299
 Biogeny, 2
 Biogeny, 33
 Bird, evolution of the, 248
 — ovum of the, 44-5, 80-1
 Biscoff, W., 17
 Bladder, evolution of the, 244, 339
 Blastema, the, 206, 213
 Blastocyst, the, 62, 74
 Blastocyst, the, 99
 Blastocyst, the, 62, 119, 120
 Blastoderm, the, 62
 Blastodermic vesicle, the, 219
 Blastopore, the, 62
 Blastopores, the, 62, 119
 Blastula, the, 62, 74
 — the mammal, 119
 Blood, importance of the, 318
 — recent experiments in mixture of, 173
 — structure of the, 319
 Blood-cells, the, 319
 Blood-vascular, the, 318-25
 — development of the, 168
 — of the vertebrate, 110
 — origin of the, 322-27
 Bomboe VILL., Bull of, 10
 Bonnet, 13
 Borneo wood-ape, the, 254
 Boveri, Theodor, 163
 Brachyura, 257
 Brain and mind, 275, 354-56
 — evolution of the, 8, 275-80
 — in the fish, 276
 — in the lower animals, 275
 — structure of the, 273-74
 Branchial arches, evolution of the, 293
 — cavity, the, 183, 189
 — system, the, 110
 Branchiostomes, 149
 Breasts, the, 113
 Bulbula, 184
 Calymniches, 234
Calymniches, 217
 Capillaries, the, 313
 Carcinoderm, the, 249
 Carboniferous strata, 202
 Carcinoma, 234
 Cardiac cavity, the, 170
 Cardiac, the, 328
 Catelecta, 213
 Caryobates, 34, 54
 Caryobates, 42
 Caryophylla, 34, 54
 Caryophylla, 42
 Caryon, 37
 Caryoplasm, 37

- Endoblast, 20, 64
Endothelium, 321
Enterococcus, 25, 223
Enteropneusta, 226
Entoderm, the, 20, 64
Eocene strata, 223
Epithelia, 229
Epiblast, 20, 64
Epidermis, the, 108, 268
Epididymus, the, 342
Epigastria, 20
Epiglottis, 11, 23
Epiglottis, the, 209
Epiphysis, the, 108
Episoma, 129
Episomata, 130, 194
Epispadia, 326
Epithelia, 37
Epithelia, 243, 251
Epovarium, the, 342
Equilibrium, sense of, 291
Euthenychuda, 257
Eustachian tube, the, 289
Euthenia, 253
Eve, 12
Evolution theory, the, 11, 208
— inductive nature of, 30
Eye, evolution of the, 284-288
— structure of the, 285
Eyelid, the third, 32
Eyelids, evolution of the, 288
FABRICIUS ab Aquapendente, 10
Face, embryonic development of the, 264
Fat glands in the skin, 269
Feathers, evolution of, 270
Fertilization, 51
— place of, 219
Fin, evolution of the, 239, 304
Fistula caecum, 6
Flagellate cells, 193
Floating bladder, the, 233, 241
— evolution of the, 314
Fetal circulation, 170-71
Food-yell, the, 67, 116-17
Foot, evolution of the, 241, 304-6
— of the ape and man, 251-59
Fore brain, the, 278
Fore kidneys, the, 256, 327
Fossiliferous strata, list of, 221
Fossils, 180
— scarcity of, 208
Free will, 295
Friedenthal, experiments of, 172
Frog, the, 241-42
— ova of the, 71-2
Fromina, 224
Function and structure, 7
Function of ova, 74
GARRYMAN'S duct, 341, 350
Ganglia, commencement of, 268
Ganglionic cell, the, 39
Gastrea, 253, 254
Gastrula, the, 3, 20, 206
— formation of the, 213
Germ theory, the, 20, 54, 59
Germinal disk, 214
Germinal vesicle, 214
Gastrocnemius, the, 62, 119, 120
Gastroepithelium, 213
Gastrotrichia, 214
Gastrula, the, 3, 20, 62
Gastrulation, 62
Gegenbauer, Carl, 270
— on evolution, 32
— on the skull, 300-1
Gemmation, 331
General Morphology, 8, 39
Genes, 23
Genital pore, the, 325
Geological evolution, length of, 200
— periods, 201
Geology, methods of, 180
— use of, 24
Germ-plans, theory of, 349
Germinal disk, 214, 215
— layers, the, 14, 16
— scheme of the, 32
— spot, the, 44
— vesicle, the, 43, 54
Germinal area, the, 121
Giant gorilla, the, 176
Gibbon, the, 173, 262
Gill-clefts and arches, 110
— formation of the, 151-2, 303
Gill-clefts, the, 183, 189
Gills, disappearance of the, 244
Glossopium, 210
Gnathostoma, 230, 232
Goethe as an evolutionist, 27, 299
Gonads, 110
Gonads, the, 111
— formation of the, 109-20
Gonads, 334
Gonochorism, beginning of, 322
Gonoducts, 322
Gonotomes, 145, 249
Gooder, 189
Gorilla, the, 174, 176, 262
Graafian follicles, the, 17, 179, 247
Gregarium, 211
Gillet-ganglion, the, 190
Gut, evolution of the, 310-17
Gyrus, 242
Gynecoclastism, 114
HAG-RISH, the, 188
Hair, evolution of the, 270
— on the human embryo and infant, 272
Hair, restriction of, by sexual selection, 271
Haliophanes, 215
Halmanna, 202
Hallier, Albrecht, 12
Haliophanes viridis, 213
Hand, evolution of the, 230, 304-6
— of the ape and man, 258
Hapalidae, 261
Harderian gland, the, 286
Hare-lip, 284
Harrison, Granville, 162
Hartmann, 262
Harvey, 10
Hatschek, 192
Hatters, 223, 265
Head-cavity, the, 138
Head-plates, the, 149
Heart, development of the, 7, 10, 117, 121, 170, 322, 324-27
— of the acedia, the, 190
— position of the, 327
Heimholtz, 207
Helminthes, 213
Hepatic gut, the, 103, 376
Hereditary, nature of, 3, 5, 27, 26-7, 349
Hermaphroditism, 2, 23, 114, 218, 322, 345
Hertwig, 21
Hesperophtheca, 259
Hu, W., 19
Histology, 18, 19
History of Creation, 6, 30
Holebiastic era, 67, 71, 77
Homocercus, 244, 245
Homology of the germinal layers, 20
Hook, evolution of the, 270
Hunterian ligament, the, 345
Huxley law, the, 171, 257, 262
Huxley, T. H., 7, 20, 29
Hydra, the, 69, 217
Hydrostatic apparatus in the fish, 325
Hydrates, 173, 262
Hydras Martinicensis, 241
Hynd house, the, 259
Hypernatism, 113
Hyperthelium, 113
Hypoblast, 20, 64
Hypoblastical groove, the, 110, 284, 286, 316
Hypodermis, the, 268
Hypopodermis, 257
Hypodermis, the, 129
Hypodermis, 170, 194
Hypopodia, 345
ICHTHYODONT, 224
Ichthyophis glaucus, 20
Ichthyofida, 257
Idea, the, 310
Immutability, Aristotle on, 10

- immortality of the soul, 56
 Impregnation-see, the, 53
 Indochina, 255
 Indo-Germanic languages, 373
 Induction and deduction, 37,
 206
 Inheritance of acquired char-
 acters, 349
 Insects, intelligence of, 353
 Intestine, the, 309, 326-33
 Intravaginal, 62
 Iris, the, 266
 Ischia, 261
 Iva, apocryphal, 263, 264
 Iva, evolution of the, 301
 Japanese strata, 202

 KANT, death of, 23
 Kelvin, Lord, on the origin
 of life, 207
 Kidneys, the, 111
 — formation of the, 139-51,
 336-42
 Kiesel, 262
 Köhler, 21
 Kowalevsky, 291

 LABIA, the, 346
 Labyrinth, the, 200
 Lacrymal glands, 269
 Landmark, J., 23, 25-7
 — theories of, 26, 349
 Lamprey, the, 130
 — ova of the, 73
 Lancelotti, the, 60, 181-95
 — description of the, 105
 Languegen, evolution of, 303
 Lanugo of the embryo, 271
 Larynx, the, 309
 — evolution of the, 314
 Lateral, the, 45
 Lateral plates, the, 129
 Laureolus strata, 201
 Leontothoma, the, 117
 Leg, evolution of the, 303
 — structure of the, 306
 Lemniscata, 137
 Lemniscata, 137
 Lemniscata, the, 237
 Lepidoptera, 137
 Leucocytes, 319
 Lila, age of, 200
 Lilia, evolution of the, 132,
 239, 304
 Limiting furrow, the, 123
 Linn, 42
 Liver, the, 309, 317
 Long-tongued ape, the, 164
 Love, importance of in nature,
 232
 Lungs, the, 110
 — evolution of the, 241,
 314-15
 Lyall, Sir C., 24
 Lymphatic vessels, the, 318
 Lymphocells, the, 319

 MACROPODIFORM, 331
 Macrospores, 331
 Macrophysa pleuralis, 213
 Male womb, the, 344, 393
 Mallochorion, the, 166
 Mallotheria, 237
 Malpighian capillaries, 339, 341
 Mammary characters of the, 112
 — gastrulation of the, 84
 Mammary, unity of the, 247-48
 Mammary glands, the, 113, 269
 Man and the ape, relation of,
 262, 357
 — origin of, 29
 Man's Place in Nature, 7, 29, 357
 Mantle, the, 189
 Mantle-folds, the, 185
 Marsupials, the, 290-52
 — ova of the, 85
 Materialism, 356
 Mathematical method, the, 30
 Mechanical causes, 6
 — embryology, 8, 19, 23
 Meckel's cartilage, 304
 Medulla capitis, the, 273
 — elongate, the, 274
 — spiracle, the, 273
 Medullary groove, the, 125
 — tube, the, 107, 228
 — formation of the,
 131, 133, 227, 267, 276
 Meisner, E., on the biogenetic
 law, 3
 Microbiotic era, 67, 71, 76
 Microcytes, 66, 321
 Mesentery, the, 98, 109, 310, 316
 Mesocardium, the, 317
 Mesoderm, the, 20, 64, 98, 99
 Mesogastria, 215
 Mesosporidia, the, 338
 Mesosporidia, the, 336
 Mesosporidia, the, 344
 Mesosporidia, the, 344
 Mesozoic period, the, 202
 Metagastria, the, 64
 Metagastria, the, 67
 Metaphorism, 143
 Metaphorism, the, 341
 Metanephros, the, 336
 Metanephros, 39
 Metanephros, 64, 67
 Metanephros, 248
 Metanephros, 20, 64
 Metanephros, the, 81
 Microgonidium, 331
 Microspores, 331
 Middle ear, the, 297
 Migration, effect of, 53
 Milk, secretion of the, 269
 Mind, evolution of, 253-54
 — in the lower animals, 353
 Muscops strata, 203
 Musculus, 40, 41
 Musculus, 40, 206, 209
 Musculus, 4, 356
 Monodelphism, 248
 Monodelphism, 223
 Monodelphism, 223
 Monodelphism, 118, 249
 — ova of the, 84
 Monodelphism, 248
 Monodelphism, 248
 Morphology, 2, 27
 Morula, the, 62, 213
 Motor-generative layer, the,
 19
 Mouth, development of the,
 124, 130
 — structure of the, 308
 Mouth layer, the, 16
 Müllerian duct, the, 347
 Muscle-layer, the, 16
 Muscles, evolution of the, 307
 — of the ear, rudimentary,
 292
 Myotomes, 108, 146
 Myonodes, the, 188, 290

 NARX, evolution of the, 270
 Naval pits, 283
 Natural philosophy, 25
 — selection, 26, 28, 349
 Navel, the, 117, 134
 Necrocytes, 357
 Necrocytes, the, 314
 Nerve, 224-26
 Nephroduct, evolution of the,
 358-59
 Nephrocytes, 149, 358
 Nerve-cell, the, 39
 Nerve, anatomy without, 267
 Nervous system, evolution of
 the, 7, 267
 Neurentic canal, the, 127
 Nodulating membrane, the,
 24, 265, 288
 Nose, the, in man and the
 ape, 164
 — development of the, 282-
 83
 — structure of the, 283
 Notochorda, the, 107
 Nuclei, 37
 Nucleolus, 44
 Nucleolus, the, 38, 44, 54
 Nucleus of the cell, 37

 OEGASIA, the, 309, 376
 Oen, 2, 27, 300
 Oen's bodies, 339
 Oligocene strata, 203
 Olfactory, 277
 On the generation of animals,
 Ontogeny, 2, 23
 — defective evidence of, 308
 Opaque area, the, 122

- Opossum, the, 152
 — ova of the, 85
 Optic nerve, the, 287
 Optic thalamus, 171
 — vesicles, the, 286
 Orang, the, 174, 268
 Ornithodorus, 248
 Ornithomyces, 85, 142
 Ornithoptera, 249
 Ossicles of the ear, 289
 Otoliths, 289
 Ova, number of, 347
 — of the lancelet, 192
 Ovarium, evolution of the, 153-
 34
 Oviduct, origin of the, 335-342
 Orobolus, the, 44
 Ovulus, 11
 Ovary, discovery of the, 76
 — nature of the, 40, 43-5
 — sets of the, 44

 PACHYLEMUS, the, 257
 Pacuan coracoides, 281
 Paleontology, 2
 — evolutionary evidence of,
 31
 — incompleteness of, 208
 — use of, 24
 Paleozoic age, the, 202
 Palaeozoic, 4
 Palaeogenetic structures, 4
 Palaeotherium, 244, 246
 Palaeozoic mammals, the, 250
 Paradidymus, the, 345
 Parietal zone, the, 129
 Parthenogenesis, 9, 13
 Parvula, Miss Julia, 154
 Pechana, 152
 Pelvic area, the, 123
 Pelvic cavity, the, 158
 Pemmophilus gmelinensis,
 215
 Penis-bone, the, 346
 Penis, variation of the, 345
 Pemmocle, 254
 Pemmocle, 68
 Pemmocle cavity, the, 185,
 196
 Pemmocle cavity, the, 338
 Pemmocle, the, 108, 163
 — formation of the, 136
 Pemmocle, 89
 Pemmocle, 202
 Pemmocle, the, 188, 190
 Pemmocle, 49, 320
 Pemmocle ganglion, the, 175
 Pemmocle, the, 309
 Pemmocle, comparison with,
 203
 Pemmocle Zoologica, 25
 Pemmocle and evolution, 6
 Pemmocle, 209
 Pemmocle, 2, 23
 Pemmocle, 114

 Physiology, backwardness of,
 7
 Phylogenesis, 209
 Pinnal eye, the, 108
 Pinnal, the, 291
 Pinnal, 263, 264
 Pinnal, principle, the,
 171
 Pinnal, the, 166, 153-54
 Pinnal, the, 166
 — characters of the, 259
 — gastrulation of the, 86
 Pinnal, 49, 320
 Pinnal, parthenogenesis
 of the, 13
 Pinnal, the, 89
 Pinnal, 38, 39
 Pinnal, 40, 59
 Pinnal, 36, 40, 209
 Pinnal, 59
 Pinnal, 221
 Pinnal, the, 221
 Pinnal, 201
 Pinnal, 234
 Pinnal, 238
 Pinnal, 203
 Pinnal, 54
 Pinnal, 55
 Pinnal, the, 11, 11
 Pinnal, the, 202
 Pinnal, the, 157, 157-60
 Pinnal, 263
 Pinnal, the, 69, 89,
 124, 125
 — gut, the, 20, 63, 214
 — kidneys, the, 111, 157
 — mouth, the, 20, 53
 — segments, 143
 — streak, the, 100, 102
 — vertebrae, 144, 196, 206,
 229
 Pinnal, the, 202
 Pinnal, 192
 Pinnal, the, 192, 218
 Pinnal, 253
 Pinnal, the, 44, 119
 Pinnal, the, 245
 Pinnal, 216
 Pinnal, 256
 Pinnal, the, 20, 53
 Pinnal, 335
 Pinnal, 247
 Pinnal, the, 157
 Pinnal, 54
 — maculae, 54
 Pinnal, 69
 Pinnal, the, 196
 — duct, the, 138, 159, 186
 — evolution of the, 338
 Pinnal, 254
 Pinnal, the, 257
 Pinnal, 333
 Pinnal, 203, 209
 Pinnal, 20, 63, 221

 Protamniotes, 249-44
 Protamniotes, 210
 Protamniotes, the, 202, 244
 Protamniotes, 36, 38
 Protamniotes, 111, 136
 Protamniotes, 210
 Protamniotes, 37, 209
 Protamniotes, 238
 Protamniotes, 248
 Protamniotes, 142, 144
 Protamniotes, 20, 210
 Protamniotes, the, 148
 — plates, the, 136, 144
 Protamniotes, 20, 211
 Protamniotes, 48
 Protamniotes, 13
 Protamniotes, evolution of the, 9
 Protamniotes, 8
 Protamniotes, 203
 Protamniotes, the, 209

 QUADRATUM, the, 247
 Quadratum, 258
 Quaternary period, 209

 RABBIT, ova of the, 86-7
 Radial, the, 203
 Radial's canal, 342
 Rectum, the, 317
 Regner de Graaf, 119
 Renal system, evolution of
 the, 335-42
 Reproduction, nature of, 130-
 31
 Reptiles, 245-47
 Respiratory organs, evolution
 of the, 314-15
 — pores, the, 183, 189
 Retina, the, 286
 Rhabdomyoma, 222
 Rhodocytes, 319
 Rhodocytes, 215
 Rhynchocephalus, 245
 Rite, the, 205
 — number of the, 332
 Ruchmentary maculae, 202
 — organs, 32
 — 141 of, 349-50
 — tone, 306

 SACCOLUTE, the, 289
 Sagitta, 64, 66, 191
 — conformation of, 93
 Salamander, the, 241
 — ova of the, 74
 Saddle-shape of embryo, 128-
 29
 Saffron, 174, 261
 Saffron, 245
 Saffron, the, 12
 Saffron, 12, 36
 Saffron, T., 18, 36
 Saffron, the, 206
 Saffron, 108, 243, 148

Spermatism, the, 344
 Spermatism, some of the, 349
 Squamoid, the, 180, 181-90
 Squamous period, the, 302
 Squamation, for, 141-42
 Squamatus-cells, 34
 Squamation-sphere, the, 17
 Sphacel, 213
 — skull of the, 301
 Selection, theory of, 28
 Selachia, 166, 168
 Selenophoric, 302
 Semi-organs, evolution of
 the, 351, 352
 — number of the, 281
 — origin of the, 281
 Sensory nerves, 279
 Serousness, the, 165
 Serous layer, the, 26
 Sex-organs, early vertebrate
 form of the, 173
 — evolution of the, 333-47
 Sexual modification, identical

Spinal cord, distribution of the,
 273
 Spina, the, 47
 Spiculism, 306
 Spines, the, 308
 Spodilyl, 142
 Sponges, classification of the,
 24
 — ova of the, 49
 Spontaneous generation, 26,
 206
 Stenoccephala, 239
 Stem-cell, the, 54
 Stem-norm, the, 129
 Stenarth, evolution of the,
 311-74, 316
 — structure of the human,
 309
 Stoma, thickness of, 200-201
 Struggle for life, the, 28
 Subcutis, the, 268
 Sweat glands, 269

Umbilical, character of all
 animals, 27
 — animals, 28, 47
 Uracine, the, 375, 341
 Urinary system, evolution of
 the, 315-42
 Urogenital ducts, 315
 Utricle, muscular, the, 344
 350
 Utricle, the, 289
 Vase deferens, 325
 Vascular layer, the, 26, 168
 — system, evolution of the,
 311-25
 — structure of the, 318
 Vegetative layer, the, 16
 Veins, evolution of the, 313-24
 Ventral podia, the, 166
 Ventricles of the heart, 325
 Vermis, 220, 223
 Vermiform appendage, the,
 22, 310, 317

Shark, the, 235
 — some of the, 283
 — ova of the, 25
 — placenta of the, 9
 — skull of the, 301
 Shoulder-blade, the, 306
 Sickle-groove, the, 82, 122
 Sieve-membrane, the, 267
 Splanchnic struts, 202
 Slinic, the, 257-60
 Sphenophore, embryology of
 the, 21
 Skeleton, structure of the, 294
 Splanchnic plate, the, 146
 Skin, the, 151
 — evolution of, 266-69
 — structure of the, 269
 Skin-layer, the, 26
 Skull, evolution of the, 249,
 299-303
 — structure of the, 299
 vertebral theory of the,
 300
 Skull, the sense of, 281
 Soul, evolution of the, 353-55
 — nature of the, 31, 330
 — phylogeny of the, 8
 — seat of the, 278
 Sound, sensations of, 289-30
 Sphenoclella, 242
 Spine, some of, 291
 Specimen, nature of the, 23, 34
 Speech, evolution of, 264
 Spermatids, 335, 342
 Spermatism, evolution of the,
 333-34
 Spermatosome, the, 33-3
 — discovery of the, 15, 21
 Spinal cord, development
 the, 8

Secondary, in man
 195, 205, 159
 Tailed men, 159-60
 Taste, the sense of, 282
 Teeth, evolution of the, 314
 — of the ape and man, 259
 Telocent, 224
 Telocentrical ova, 67, 68
 Temperature, sense of, 282
 Terrestrial life, beginning of,
 232
 Tertiary period, the, 203
 Thalamus generalis, the, 13
 Theories, value of, 181
 Theromorphs, 246
 Third eye, the, 286, 288
 Thyroid gland, the, 110, 124,
 315
 Time-variations in ontogeny, 5
 Timon, primary and second-
 ary, 37
 Tongue, the, 27
 Tocomaria, 245
 Toes, number of the, 240
 Toxigenesis, the, 246
 Touch, the sense of, 282
 Tracheata, 142, 215
 Trend, the, 25, 21
 Tree-drop, the, 241
 Triassic strata, 202
 Tridacina, 24
 Troglodytes, 174
 Turbates, the, 109
 Turbellaria, 223
 Turbated bones, the, 289
 Tympanic cavity, the, 288

Umbilical cord, the, 117
 — vesicle, the, 158

Uterine, the, 144
 — evolution of the, 296
 — structure of the, 294
 Vertebrates, character of the,
 104-10
 — descent of the, 3
 Vertebrae, 142
 Viscero-umbilical ligament
 the, 341
 Visceral protrusion, the,
 338
 Villi of the chorion, 165
 Vichow, R., 35
 — on the sperm, 303
 — on the evolution of,
 264
 Virgin-birth, 9, 13
 Vitalism, 6
 Vitelline duct, the, 138
 Valeriana, 213
 WALLACE, A. R., 29
 Water, organic properties
 of, 200
 Water vessels, 356
 Weismann's theories, 349
 Wolk, C. F., 13
 Wolfen bodies, 339
 Womb, evolution of the, 342-43

Yell, the, 43, 45, 67
 Yell, the, 242, 254



